

## Binomial Theorem

---

### Exercise 10A

**Q. 1. Using binomial theorem, expand each of the following:**

$$(1 - 2x)^5$$

**Answer :** To find: Expansion of  $(1 - 2x)^5$

Formula used: (i) 
$${}^nC_r = \frac{n!}{(n-r)!(r)!}$$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $(1 - 2x)^5$

$$\Rightarrow [{}^5C_0(1)^5] + [{}^5C_1(1)^{5-1}(-2x)^1] + [{}^5C_2(1)^{5-2}(-2x)^2] + [{}^5C_3(1)^{5-3}(-2x)^3] + [{}^5C_4(1)^{5-4}(-2x)^4] + [{}^5C_5(-2x)^5]$$

$$\Rightarrow \left[ \frac{5!}{0!(5-0)!} (1)^5 \right] - \left[ \frac{5!}{1!(5-1)!} (1)^4(2x) \right] + \left[ \frac{5!}{2!(5-2)!} (1)^3(4x^2) \right]$$

$$- \left[ \frac{5!}{3!(5-3)!} (1)^2(8x^3) \right] + \left[ \frac{5!}{4!(5-4)!} (1)^1(16x^4) \right] - \left[ \frac{5!}{5!(5-5)!} (32x^5) \right]$$

$$\Rightarrow 1 - 5(2x) + 10(4x^2) - 10(8x^3) + 5(16x^4) - 1(32x^5)$$

$$\Rightarrow 1 - 10x + 40x^2 - 80x^3 + 80x^4 - 32x^5$$

On rearranging

**Ans)**  $-32x^5 + 80x^4 - 80x^3 + 40x^2 - 10x + 1$

**Q. 2. Using binomial theorem, expand each of the following:**

$$(2x - 3)^6$$

**Answer :** To find: Expansion of  $(2x - 3)^6$

Formula used: (i) 
$${}^nC_r = \frac{n!}{(n-r)!(r)!}$$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $(2x - 3)^6$

$$\Rightarrow [{}^6C_0(2x)^6] + [{}^6C_1(2x)^{6-1}(-3)^1] + [{}^6C_2(2x)^{6-2}(-3)^2] + [{}^6C_3(2x)^{6-3}(-3)^3] + [{}^6C_4(2x)^{6-4}(-3)^4] + [{}^6C_5(2x)^{6-5}(-3)^5] + [{}^6C_6(-3)^6]$$

$$\Rightarrow \left[ \frac{6!}{0!(6-0)!} (2x)^6 \right] - \left[ \frac{6!}{1!(6-1)!} (2x)^5(3) \right] + \left[ \frac{6!}{2!(6-2)!} (2x)^4(9) \right]$$

$$- \left[ \frac{6!}{3!(6-3)!} (2x)^3(27) \right] + \left[ \frac{6!}{4!(6-4)!} (2x)^2(81) \right]$$

$$- \left[ \frac{6!}{5!(6-5)!} (2x)^1(243) \right] + \left[ \frac{6!}{6!(6-6)!} (729) \right]$$

$$\Rightarrow [(1)(64x^6)] - [(6)(32x^5)(3)] + [15(16x^4)(9)] - [20(8x^3)(27)] + [15(4x^2)(81)] - [(6)(2x)(243)] + [(1)(729)]$$

$$\Rightarrow 64x^6 - 576x^5 + 2160x^4 - 4320x^3 + 4860x^2 - 2916x + 729$$

$$\text{Ans) } 64x^6 - 576x^5 + 2160x^4 - 4320x^3 + 4860x^2 - 2916x + 729$$

**Q. 3. Using binomial theorem, expand each of the following:**

$$(3x + 2y)^5$$

**Answer :** To find: Expansion of  $(3x + 2y)^5$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

We have,  $(3x + 2y)^5$

$$\Rightarrow [{}^5C_0(3x)^{5-0}] + [{}^5C_1(3x)^{5-1}(2y)^1] + [{}^5C_2(3x)^{5-2}(2y)^2] + [{}^5C_3(3x)^{5-3}(2y)^3] + [{}^5C_4(3x)^{5-4}(2y)^4] + [{}^5C_5(2y)^5]$$

$$\Rightarrow \left[ \frac{5!}{0!(5-0)!} (243x^5) \right] + \left[ \frac{5!}{1!(5-1)!} (81x^4)(2y) \right] +$$

$$\left[ \frac{5!}{2!(5-2)!} (27x^3)(4y^2) \right] + \left[ \frac{5!}{3!(5-3)!} (9x^2)(8y^3) \right] +$$

$$\left[ \frac{5!}{4!(5-4)!} (3x)(16y^4) \right] + \left[ \frac{5!}{5!(5-5)!} (32y^5) \right]$$

$$\Rightarrow [1(243x^5)] + [5(81x^4)(2y)] + [10(27x^3)(4y^2)] + [10(9x^2)(8y^3)] + [5(3x)(16y^4)] + [1(32y^5)]$$

$$\Rightarrow 243x^5 + 810x^4y + 1080x^3y^2 + 720x^2y^3 + 240xy^4 + 32y^5$$

$$\text{Ans) } 243x^5 + 810x^4y + 1080x^3y^2 + 720x^2y^3 + 240xy^4 + 32y^5$$

**Q. 4. Using binomial theorem, expand each of the following:**

$$(2x - 3y)^4$$

**Answer :** To find: Expansion of  $(2x - 3y)^4$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

We have,  $(2x - 3y)^4$

$$\Rightarrow [{}^4C_0(2x)^{4-0}] + [{}^4C_1(2x)^{4-1}(-3y)^1] + [{}^4C_2(2x)^{4-2}(-3y)^2] + [{}^4C_3(2x)^{4-3}(-3y)^3] + [{}^4C_4(-3y)^4]$$

$$\left[ \frac{4!}{0!(4-0)!} (2x)^4 \right] - \left[ \frac{4!}{1!(4-1)!} (2x)^3(3y) \right] + \left[ \frac{4!}{2!(4-2)!} (2x)^2(9y^2) \right] -$$

$$\left[ \frac{4!}{3!(4-3)!} (2x)^1(27y^3) \right] + \left[ \frac{4!}{4!(4-4)!} (81y^4) \right]$$

$$\Rightarrow [1(16x^4)] - [4(8x^3)(3y)] + [6(4x^2)(9y^2)] - [4(2x)(27y^3)] + [1(81y^4)]$$

$$\Rightarrow 16x^4 - 96x^3y + 216x^2y^2 - 216xy^3 + 81y^4$$

$$\text{Ans) } 16x^4 - 96x^3y + 216x^2y^2 - 216xy^3 + 81y^4$$

**Q. 5. Using binomial theorem, expand each of the following:**

$$\left(\frac{2x}{3} - \frac{3}{2x}\right)^6$$

**Answer :** To find: Expansion of  $\left(\frac{2x}{3} - \frac{3}{2x}\right)^6$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $\left(\frac{2x}{3} - \frac{3}{2x}\right)^6$

$$\Rightarrow \left[ {}^6C_0 \left(\frac{2x}{3}\right)^{6-0} \right] + \left[ {}^6C_1 \left(\frac{2x}{3}\right)^{6-1} \left(-\frac{3}{2x}\right)^1 \right] + \left[ {}^6C_2 \left(\frac{2x}{3}\right)^{6-2} \left(-\frac{3}{2x}\right)^2 \right] +$$

$$\left[ {}^6C_3 \left(\frac{2x}{3}\right)^{6-3} \left(-\frac{3}{2x}\right)^3 \right] + \left[ {}^6C_4 \left(\frac{2x}{3}\right)^{6-4} \left(-\frac{3}{2x}\right)^4 \right]$$

$$+ \left[ {}^6C_5 \left(\frac{2x}{3}\right)^{6-5} \left(-\frac{3}{2x}\right)^5 \right] + \left[ {}^6C_6 \left(-\frac{3}{2x}\right)^6 \right]$$

$$\Rightarrow \left[ \frac{6!}{0!(6-0)!} \left(\frac{2x}{3}\right)^6 \right] - \left[ \frac{6!}{1!(6-1)!} \left(\frac{2x}{3}\right)^5 \left(\frac{3}{2x}\right) \right] +$$

$$\left[ \frac{6!}{2!(6-2)!} \left(\frac{2x}{3}\right)^4 \left(\frac{9}{4x^2}\right) \right] - \left[ \frac{6!}{3!(6-3)!} \left(\frac{2x}{3}\right)^3 \left(\frac{27}{8x^3}\right) \right] +$$

$$\left[ \frac{6!}{4!(6-4)!} \left(\frac{2x}{3}\right)^2 \left(\frac{81}{16x^4}\right) \right] - \left[ \frac{6!}{5!(6-5)!} \left(\frac{2x}{3}\right)^1 \left(\frac{243}{32x^5}\right) \right]$$

$$+ \left[ \frac{6!}{6!(6-6)!} \left(\frac{729}{64x^6}\right) \right]$$

$$\Rightarrow \left[ 1 \left( \frac{64x^6}{729} \right) \right] - \left[ 6 \left( \frac{32x^5}{243} \right) \left( \frac{3}{2x} \right) \right] + \left[ 15 \left( \frac{16x^4}{81} \right) \left( \frac{9}{4x^2} \right) \right] - \left[ 20 \left( \frac{8x^3}{27} \right) \left( \frac{27}{8x^3} \right) \right] + \left[ 15 \left( \frac{4x^2}{9} \right) \left( \frac{81}{16x^4} \right) \right] - \left[ 6 \left( \frac{2x}{3} \right) \left( \frac{243}{32x^5} \right) \right] + \left[ 1 \left( \frac{729}{64x^6} \right) \right]$$

$$\Rightarrow \frac{64}{729}x^6 - \frac{32}{27}x^4 + \frac{20}{3}x^2 - 20 + \frac{135}{4} \frac{1}{x^2} - \frac{243}{8} \frac{1}{x^4} + \frac{729}{64} \frac{1}{x^6}$$

$$\text{Ans) } \frac{64}{729}x^6 - \frac{32}{27}x^4 + \frac{20}{3}x^2 - 20 + \frac{135}{4} \frac{1}{x^2} - \frac{243}{8} \frac{1}{x^4} + \frac{729}{64} \frac{1}{x^6}$$

**Q. 6. Using binomial theorem, expand each of the following:**

$$\left( x^2 - \frac{3}{x} \right)^7$$

**Answer :** To find: Expansion of  $\left( x^2 - \frac{3x}{7} \right)^7$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $\left( x^2 - \frac{3x}{7} \right)^7$

$$\Rightarrow \left[ {}^7C_0(x^2)^{7-0} \right] + \left[ {}^7C_1(x^2)^{7-1} \left( -\frac{3x}{7} \right)^1 \right] + \left[ {}^7C_2(x^2)^{7-2} \left( -\frac{3x}{7} \right)^2 \right] +$$

$$\left[ {}^7C_3(x^2)^{7-3} \left( -\frac{3x}{7} \right)^3 \right] + \left[ {}^7C_4(x^2)^{7-4} \left( -\frac{3x}{7} \right)^4 \right] + \left[ {}^7C_5(x^2)^{7-5} \left( -\frac{3x}{7} \right)^5 \right] +$$

$$\left[ {}^7C_6(x^2)^{7-6} \left( -\frac{3x}{7} \right)^6 \right] + \left[ {}^7C_7 \left( -\frac{3x}{7} \right)^7 \right]$$

$$\Rightarrow \left[ \frac{7!}{0!(7-0)!} (x^2)^7 \right] - \left[ \frac{7!}{1!(7-1)!} (x^2)^6 \left( \frac{3x}{7} \right) \right] + \left[ \frac{7!}{2!(7-2)!} (x^2)^5 \left( \frac{9x^2}{49} \right) \right] -$$

$$\left[ \frac{7!}{3!(7-3)!} (x^2)^4 \left( \frac{27x^3}{343} \right) \right] + \left[ \frac{7!}{4!(7-4)!} (x^2)^3 \left( \frac{81x^4}{2401} \right) \right] - \left[ \frac{7!}{5!(7-5)!} \right.$$

$$\left. (x^2)^2 \left( \frac{243x^5}{16807} \right) \right] + \left[ \frac{7!}{6!(7-6)!} (x^2)^1 \left( \frac{729x^6}{117649} \right) \right] - \left[ \frac{7!}{7!(7-7)!} \left( \frac{2187x^7}{823543} \right) \right]$$

$$\Rightarrow [1(x^{14})] - \left[ 7(x^{12}) \left( \frac{3x}{7} \right) \right] + \left[ 21(x^{10}) \left( \frac{9x^2}{49} \right) \right] - \left[ 35(x^8) \left( \frac{27x^3}{343} \right) \right] +$$

$$\left[ 35(x^6) \left( \frac{81x^4}{2401} \right) \right] - \left[ 21(x^4) \left( \frac{243x^5}{16807} \right) \right] + \left[ 7(x^2) \left( \frac{729x^6}{117649} \right) \right] -$$

$$\left[ 1 \left( \frac{2187x^7}{823543} \right) \right]$$

$$\Rightarrow x^{14} - 3x^{13} + \left( \frac{27}{7} \right) x^{12} - \left( \frac{135}{49} \right) x^{11} + \left( \frac{405}{343} \right) x^{10} -$$

$$\left( \frac{729}{2401} \right) x^9 + \left( \frac{729}{16807} \right) x^8 - \left( \frac{2187}{823543} \right) x^7$$

**Ans)**

$$x^{14} - 3x^{13} + \left( \frac{27}{7} \right) x^{12} - \left( \frac{135}{49} \right) x^{11} + \left( \frac{405}{343} \right) x^{10} - \left( \frac{729}{2401} \right) x^9 + \left( \frac{729}{16807} \right) x^8 -$$

$$\left( \frac{2187}{823543} \right) x^7$$

**Q. 7. Using binomial theorem, expand each of the following:**

$$\left( x - \frac{1}{y} \right)^5$$

**Answer :** To find: Expansion of  $\left( x - \frac{1}{y} \right)^5$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $\left(x - \frac{1}{y}\right)^5$

$$\Rightarrow {}^5C_0(x)^{5-0} + {}^5C_1(x)^{5-1}\left(-\frac{1}{y}\right)^1 + {}^5C_2(x)^{5-2}\left(-\frac{1}{y}\right)^2 + {}^5C_3(x)^{5-3}\left(-\frac{1}{y}\right)^3 + {}^5C_4(x)^{5-4}\left(-\frac{1}{y}\right)^4 + {}^5C_5\left(-\frac{1}{y}\right)^5$$

$$\Rightarrow \left[ \frac{5!}{0!(5-0)!} (x^5) \right] - \left[ \frac{5!}{1!(5-1)!} (x^4) \left(\frac{1}{y}\right)^1 \right] + \left[ \frac{5!}{2!(5-2)!} (x^3) \left(\frac{1}{y^2}\right) \right]$$

$$- \left[ \frac{5!}{3!(5-3)!} (x^2) \left(\frac{1}{y^3}\right) \right] + \left[ \frac{5!}{4!(5-4)!} (x) \left(\frac{1}{y^4}\right) \right] - \left[ \frac{5!}{5!(5-5)!} \left(\frac{1}{y^5}\right) \right]$$

$\Rightarrow$

$$[1(x^5)] - \left[ 5 \left(\frac{x^4}{y}\right) \right] + \left[ 10 \left(\frac{x^3}{y^2}\right) \right] - \left[ 10 \left(\frac{x^2}{y^3}\right) \right] + \left[ 5 \left(\frac{x}{y^4}\right) \right] - [1(y^5)]$$

$$\Rightarrow x^5 - 5 \frac{x^4}{y} + 10 \frac{x^3}{y^2} - 10 \frac{x^2}{y^3} + 5 \frac{x}{y^4} - y^5$$

Ans)  $x^5 - 5 \frac{x^4}{y} + 10 \frac{x^3}{y^2} - 10 \frac{x^2}{y^3} + 5 \frac{x}{y^4} - y^5$

**Q. 8. Using binomial theorem, expand each of the following:**

$$\left(\sqrt{x} + \sqrt{y}\right)^8$$

**Answer :** To find: Expansion of  $\left(\sqrt{x} + \sqrt{y}\right)^8$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$$

We have,  $(\sqrt{x} + \sqrt{y})^8$

We can write  $\sqrt{x}$  as  $x^{\frac{1}{2}}$  and  $\sqrt{y}$  as  $y^{\frac{1}{2}}$

Now, we have to solve for  $(x^{\frac{1}{2}} + y^{\frac{1}{2}})^8$

$$\Rightarrow \left[ {}^8C_0 \left( x^{\frac{1}{2}} \right)^{8-0} \right] + \left[ {}^8C_1 \left( x^{\frac{1}{2}} \right)^{8-1} \left( y^{\frac{1}{2}} \right)^1 \right] + \left[ {}^8C_2 \left( x^{\frac{1}{2}} \right)^{8-2} \left( y^{\frac{1}{2}} \right)^2 \right] +$$

$$\left[ {}^8C_3 \left( x^{\frac{1}{2}} \right)^{8-3} \left( y^{\frac{1}{2}} \right)^3 \right] + \left[ {}^8C_4 \left( x^{\frac{1}{2}} \right)^{8-4} \left( y^{\frac{1}{2}} \right)^4 \right] + \left[ {}^8C_5 \left( x^{\frac{1}{2}} \right)^{8-5} \left( y^{\frac{1}{2}} \right)^5 \right] +$$

$$\left[ {}^8C_6 \left( x^{\frac{1}{2}} \right)^{8-6} \left( y^{\frac{1}{2}} \right)^6 \right] + \left[ {}^8C_7 \left( x^{\frac{1}{2}} \right)^{8-7} \left( y^{\frac{1}{2}} \right)^7 \right] + \left[ {}^8C_8 \left( y^{\frac{1}{2}} \right)^8 \right]$$

$$\Rightarrow \left[ \frac{8!}{0!(8-0)!} \left( x^{\frac{8}{2}} \right) \right] + \left[ \frac{8!}{1!(8-1)!} \left( x^{\frac{7}{2}} \right) \left( y^{\frac{1}{2}} \right) \right] + \left[ \frac{8!}{2!(8-2)!} \left( x^{\frac{6}{2}} \right) \left( y^{\frac{2}{2}} \right) \right] +$$

$$\left[ \frac{8!}{3!(8-3)!} \left( x^{\frac{5}{2}} \right) \left( y^{\frac{3}{2}} \right) \right] + \left[ \frac{8!}{4!(8-4)!} \left( x^{\frac{4}{2}} \right) \left( y^{\frac{4}{2}} \right) \right] + \left[ \frac{8!}{5!(8-5)!} \left( x^{\frac{3}{2}} \right) \left( y^{\frac{5}{2}} \right) \right] +$$

$$\left[ \frac{8!}{6!(8-6)!} \left( x^{\frac{2}{2}} \right) \left( y^{\frac{6}{2}} \right) \right] + \left[ \frac{8!}{7!(8-7)!} \left( x^{\frac{1}{2}} \right) \left( y^{\frac{7}{2}} \right) \right] + \left[ \frac{8!}{8!(8-8)!} \left( y^{\frac{8}{2}} \right) \right]$$

$$\Rightarrow [1(x^4)] + \left[ 8 \left( x^{\frac{7}{2}} \right) \left( y^{\frac{1}{2}} \right) \right] + [28(x^3)(y)] + \left[ 56 \left( x^{\frac{5}{2}} \right) \left( y^{\frac{3}{2}} \right) \right]$$

$$+ [70(x^2)(y^2)] + \left[ 56 \left( x^{\frac{3}{2}} \right) \left( y^{\frac{5}{2}} \right) \right] + [28(x^1)(y^3)] + \left[ 8 \left( x^{\frac{1}{2}} \right) \left( y^{\frac{7}{2}} \right) \right] + [1(y^4)]$$

**Ans)**  $(x^4) + 8(x^{7/2})(y^{1/2}) + 28(x^3)(y) + 56(x^{5/2})(y^{3/2}) + 70(x^2)(y^2) + 56(x^{3/2})(y^{5/2}) +$   
 $28(x^1)(y^3) + 8(x^{1/2})(y^{7/2}) + (y^4)$

**Q. 9. Using binomial theorem, expand each of the following:**

$$\left( \sqrt[3]{x} - \sqrt[3]{y} \right)^6$$



**Answer :** To find: Expansion of  $(\sqrt[3]{x} - \sqrt[3]{y})^6$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $(\sqrt[3]{x} - \sqrt[3]{y})^6$

We can write  $\sqrt[3]{x}$  as  $x^{\frac{1}{3}}$  and  $\sqrt[3]{y}$  as  $y^{\frac{1}{3}}$

Now, we have to solve for  $(x^{\frac{1}{3}} - y^{\frac{1}{3}})^6$

$$\Rightarrow \left[ {}^6C_0 \left( x^{\frac{1}{3}} \right)^{6-0} \right] + \left[ {}^6C_1 \left( x^{\frac{1}{3}} \right)^{6-1} \left( -y^{\frac{1}{3}} \right)^1 \right] + \left[ {}^6C_2 \left( x^{\frac{1}{3}} \right)^{6-2} \left( -y^{\frac{1}{3}} \right)^2 \right] +$$

$$\left[ {}^6C_3 \left( x^{\frac{1}{3}} \right)^{6-3} \left( -y^{\frac{1}{3}} \right)^3 \right] + \left[ {}^6C_4 \left( x^{\frac{1}{3}} \right)^{6-4} \left( -y^{\frac{1}{3}} \right)^4 \right] + \left[ {}^6C_5 \left( x^{\frac{1}{3}} \right)^{6-5} \left( -y^{\frac{1}{3}} \right)^5 \right] +$$

$$\left[ {}^6C_6 \left( -y^{\frac{1}{3}} \right)^6 \right]$$

$$\Rightarrow \left[ {}^6C_0 \left( x^{\frac{6}{3}} \right) \right] - \left[ {}^6C_1 \left( x^{\frac{5}{3}} \right) \left( y^{\frac{1}{3}} \right) \right] + \left[ {}^6C_2 \left( x^{\frac{4}{3}} \right) \left( y^{\frac{2}{3}} \right) \right] - \left[ {}^6C_3 \left( x^{\frac{3}{3}} \right) \left( y^{\frac{3}{3}} \right) \right] +$$

$$\left[ {}^6C_4 \left( x^{\frac{2}{3}} \right) \left( y^{\frac{4}{3}} \right) \right] - \left[ {}^6C_5 \left( x^{\frac{1}{3}} \right) \left( y^{\frac{5}{3}} \right) \right] + \left[ {}^6C_6 \left( y^{\frac{6}{3}} \right) \right]$$

$$\Rightarrow \left[ \frac{6!}{0!(6-0)!} (x^2) \right] - \left[ \frac{6!}{1!(6-1)!} \left( x^{\frac{5}{3}} \right) \left( y^{\frac{1}{3}} \right) \right] + \left[ \frac{6!}{2!(6-2)!} \left( x^{\frac{4}{3}} \right) \left( y^{\frac{2}{3}} \right) \right]$$

$$- \left[ \frac{6!}{3!(6-3)!} (x)(y) \right] + \left[ \frac{6!}{4!(6-4)!} \left( x^{\frac{2}{3}} \right) \left( y^{\frac{4}{3}} \right) \right] - \left[ \frac{6!}{5!(6-5)!} \left( x^{\frac{1}{3}} \right) \left( y^{\frac{5}{3}} \right) \right]$$

$$+ \left[ \frac{6!}{6!(6-6)!} (y^2) \right]$$

$$\Rightarrow [1(x^2)] - \left[6\left(x^{\frac{5}{3}}\right)\left(y^{\frac{1}{3}}\right)\right] + \left[15\left(x^{\frac{4}{3}}\right)\left(y^{\frac{2}{3}}\right)\right] - [20(x)(y)] + \left[15\left(x^{\frac{2}{3}}\right)\left(y^{\frac{4}{3}}\right)\right] - \left[6\left(x^{\frac{1}{3}}\right)\left(y^{\frac{5}{3}}\right)\right] + [1(y^2)]$$

$$\Rightarrow x^2 - 6x^{\frac{5}{3}}y^{\frac{1}{3}} + 15x^{\frac{4}{3}}y^{\frac{2}{3}} - 20xy + 15x^{\frac{2}{3}}y^{\frac{4}{3}} - 6x^{\frac{1}{3}}y^{\frac{5}{3}} + y^2$$

$$\text{Ans)} x^2 - 6x^{5/3}y^{1/3} + 15x^{4/3}y^{2/3} - 20xy + 15x^{2/3}y^{4/3} - 6x^{1/3}y^{5/3} + y^2$$

**Q. 10. Using binomial theorem, expand each of the following:**

$$(1 + 2x - 3x^2)^4$$

**Answer :** To find: Expansion of  $(1 + 2x - 3x^2)^4$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$\text{We have, } (1 + 2x - 3x^2)^4$$

$$\text{Let } (1+2x) = a \text{ and } (-3x^2) = b \dots (i)$$

$$\text{Now the equation becomes } (a + b)^4$$

$$\Rightarrow [{}^4C_0(a)^{4-0}] + [{}^4C_1(a)^{4-1}(b)^1] + [{}^4C_2(a)^{4-2}(b)^2] + [{}^4C_3(a)^{4-3}(b)^3] + [{}^4C_4(b)^4]$$

$$\Rightarrow [{}^4C_0(a)^4] + [{}^4C_1(a)^3(b)^1] + [{}^4C_2(a)^2(b)^2] + [{}^4C_3(a)(b)^3] + [{}^4C_4(b)^4]$$

(Substituting value of b from eqn. i)

$$\Rightarrow \left[ \frac{4!}{0!(4-0)!} (a)^4 \right] + \left[ \frac{4!}{1!(4-1)!} (a)^3(-3x^2)^1 \right] + \left[ \frac{4!}{2!(4-2)!} (a)^2(-3x^2)^2 \right]$$

$$+ \left[ \frac{4!}{3!(4-3)!} (a) (-3x^2)^3 \right] + \left[ \frac{4!}{4!(4-4)!} (-3x^2)^4 \right]$$

(Substituting value of b from eqn. i)

$$\Rightarrow [1(1+2x)^4] - [4(1+2x)^3(3x^2)] + [6(1+2x)^2(9x^4)] - [4(1+2x)(27x^6)^3] + [1(81x^8)^4] \quad \dots(ii)$$

We need the value of  $a^4, a^3$  and  $a^2$ , where  $a = (1+2x)$

For  $(1+2x)^4$ , Applying Binomial theorem

$$(1+2x)^4 \Rightarrow$$

$${}^4C_0(1)^{4-0} + {}^4C_1(1)^{4-1}(2x)^1 + {}^4C_2(1)^{4-2}(2x)^2 + {}^4C_3(1)^{4-3}(2x)^3 + {}^4C_4(2x)^4$$

$$\Rightarrow \frac{4!}{0!(4-0)!} (1)^4 + \frac{4!}{1!(4-1)!} (1)^3(2x)^1 + \frac{4!}{2!(4-2)!} (1)^2(2x)^2$$

$$+ \frac{4!}{3!(4-3)!} (1)(2x)^3 + \frac{4!}{4!(4-4)!} (2x)^4$$

$$\Rightarrow [1] + [4(1)(2x)] + [6(1)(4x^2)] + [4(1)(8x^3)] + [1(16x^4)]$$

$$\Rightarrow 1 + 8x + 24x^2 + 32x^3 + 16x^4$$

$$\text{We have } (1+2x)^4 = 1 + 8x + 24x^2 + 32x^3 + 16x^4 \dots (iii)$$

For  $(a+b)^3$ , we have formula  $a^3+b^3+3a^2b+3ab^2$

For,  $(1+2x)^3$ , substituting  $a = 1$  and  $b = 2x$  in the above formula

$$\Rightarrow 1^3 + (2x)^3 + 3(1)^2(2x) + 3(1)(2x)^2$$

$$\Rightarrow 1 + 8x^3 + 6x + 12x^2$$

$$\Rightarrow 8x^3 + 12x^2 + 6x + 1 \dots (iv)$$

For  $(a+b)^2$ , we have formula  $a^2+2ab+b^2$

For,  $(1+2x)^2$ , substituting  $a = 1$  and  $b = 2x$  in the above formula

$$\Rightarrow (1)^2 + 2(1)(2x) + (2x)^2$$

$$\Rightarrow 1 + 4x + 4x^2$$

$$\Rightarrow 4x^2 + 4x + 1 \dots (v)$$

Putting the value obtained from eqn. (iii), (iv) and (v) in eqn. (ii)

$$\begin{aligned}
&\Rightarrow 1(1 + 8x + 24x^2 + 32x^3 + 16x^4) - 4(8x^3 + 12x^2 + 6x + 1)(3x^2) \\
&+ 6(4x^2 + 4x + 1)(9x^4) - 4(1+2x)(27x^6)^3 + 1(81x^8) \\
&\Rightarrow 1(1 + 8x + 24x^2 + 32x^3 + 16x^4) - 4(24x^5 + 36x^4 + 18x^3 + 3x^2) \\
&+ 6(36x^6 + 36x^5 + 9x^4) - 4(27x^6 + 54x^7) + 1(81x^8) \\
&\Rightarrow 1 + 8x + 24x^2 + 32x^3 + 16x^4 - 96x^5 - 144x^4 - 72x^3 - 12x^2 + 216x^6 + 216x^5 + 54x^4 - \\
&108x^6 - 216x^7 + 81x^8
\end{aligned}$$

On rearranging

$$\text{Ans)} 81x^8 - 216x^7 + 108x^6 + 120x^5 - 74x^4 - 40x^3 + 12x^2 + 8x + 1$$

**Q. 11. Using binomial theorem, expand each of the following:**

$$\left(1 + \frac{x}{2} - \frac{2}{x}\right)^4, x \neq 0$$

**Answer :** To find: Expansion of  $\left(1 + \frac{x}{2} - \frac{2}{x}\right)^4, x \neq 0$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

We have,  $\left(1 + \frac{x}{2} - \frac{2}{x}\right)^4, x \neq 0$

Let  $\left(1 + \frac{x}{2}\right) = a$  and  $\left(-\frac{2}{x}\right) = b \dots (i)$

Now the equation becomes  $(a + b)^4$

$$\Rightarrow [{}^4C_0(a)^{4-0}] + [{}^4C_1(a)^{4-1}(b)^1] + [{}^4C_2(a)^{4-2}(b)^2] + [{}^4C_3(a)^{4-3}(b)^3] + [{}^4C_4(b)^4]$$

$$\Rightarrow [{}^4C_0(a)^4] + [{}^4C_1(a)^3(b)^1] + [{}^4C_2(a)^2(b)^2] + [{}^4C_3(a)(b)^3] + [{}^4C_4(b)^4]$$

(Substituting value of b from eqn. i)

$$\Rightarrow \left[ \frac{4!}{0!(4-0)!} (a)^4 \right] + \left[ \frac{4!}{1!(4-1)!} (a)^3 \left( -\frac{2}{x} \right)^1 \right] + \left[ \frac{4!}{2!(4-2)!} (a)^2 \left( -\frac{2}{x} \right)^2 \right] + \left[ \frac{4!}{3!(4-3)!} (a)^1 \left( -\frac{2}{x} \right)^3 \right] + \left[ \frac{4!}{4!(4-4)!} \left( -\frac{2}{x} \right)^4 \right]$$

(Substituting value of a from eqn. i)

$$\Rightarrow \left[ 1 \left( 1 + \frac{x}{2} \right)^4 \right] - \left[ 4 \left( 1 + \frac{x}{2} \right)^3 \left( \frac{2}{x} \right) \right] + \left[ 6 \left( 1 + \frac{x}{2} \right)^2 \left( \frac{4}{x^2} \right) \right] - \left[ 4 \left( 1 + \frac{x}{2} \right)^1 \left( \frac{8}{x^3} \right) \right] + \left[ 1 \left( \frac{16}{x^4} \right) \right] \dots (ii)$$

We need the value of  $a^4, a^3$  and  $a^2$ , where  $a = \left( 1 + \frac{x}{2} \right)$

For  $\left( 1 + \frac{x}{2} \right)^4$ , Applying Binomial theorem

$$\left( 1 + \frac{x}{2} \right)^4 = \left[ {}^4C_0(1)^{4-0} \right] + \left[ {}^4C_1(1)^4 - 1 \left( \frac{x}{2} \right)^1 \right] + \left[ {}^4C_2(1)^4 - 2 \left( \frac{x}{2} \right)^2 \right] + \left[ {}^4C_3(1)^4 - 3 \left( \frac{x}{2} \right)^3 \right] + \left[ {}^4C_4 \left( \frac{x}{2} \right)^4 \right]$$

$$\Rightarrow \left[ \frac{4!}{0!(4-0)!} (1)^4 \right] + \left[ \frac{4!}{1!(4-1)!} (1)^3 \left( \frac{x}{2} \right)^1 \right] + \left[ \frac{4!}{2!(4-2)!} (1)^2 \left( \frac{x}{2} \right)^2 \right]$$

$$+ \left[ \frac{4!}{3!(4-3)!} (1) \left( \frac{x}{2} \right)^3 \right] + \left[ \frac{4!}{4!(4-4)!} \left( \frac{x}{2} \right)^4 \right]$$

$$\Rightarrow [1] + \left[ 4(1) \left( \frac{x}{2} \right) \right] + \left[ 6(1) \left( \frac{x^2}{4} \right) \right] + \left[ 4(1) \left( \frac{x^3}{8} \right) \right] + \left[ 1 \left( \frac{x^4}{16} \right) \right]$$

$$\Rightarrow 1 + 2x + \frac{3}{2}x^2 + \frac{x^3}{2} + \frac{x^4}{16}$$

On rearranging the above eqn.

$$\Rightarrow \frac{1}{16}x^4 + \frac{1}{2}x^3 + \frac{3}{2}x^2 + 2x + 1 \dots \text{(iii)}$$

We have,  $\left(1 + \frac{x}{2}\right)^4 = \frac{1}{16}x^4 + \frac{1}{2}x^3 + \frac{3}{2}x^2 + 2x + 1$

For,  $(a+b)^3$ , we have formula  $a^3+b^3+3a^2b+3ab^2$

For,  $\left(1 + \frac{x}{2}\right)^3$ , substituting  $a = 1$  and  $b = \frac{x}{2}$  in the above formula

$$\Rightarrow 1^3 + \left(\frac{x}{2}\right)^3 + 3(1)^2\left(\frac{x}{2}\right) + 3(1)\left(\frac{x}{2}\right)^2$$

$$\Rightarrow 1 + \left(\frac{x^3}{8}\right) + \left(\frac{3x}{2}\right) + \left(\frac{3x^2}{4}\right)$$

$$\Rightarrow \left(\frac{x^3}{8}\right) + \left(\frac{3x^2}{4}\right) + \left(\frac{3x}{2}\right) + 1 \dots \text{(iv)}$$

For,  $(a+b)^2$ , we have formula  $a^2+2ab+b^2$

For,  $\left(1 + \frac{x}{2}\right)^2$ , substituting  $a = 1$  and  $b = \frac{x}{2}$  in the above formula

$$\Rightarrow (1)^2 + 2(1)\left(\frac{x}{2}\right) + \left(\frac{x}{2}\right)^2$$

$$\Rightarrow 1 + x + \left(\frac{x^2}{4}\right)$$

$$\Rightarrow \frac{x^2}{4} + x + 1 \dots \text{(v)}$$

Putting the value obtained from eqn. (iii),(iv) and (v) in eqn. (ii)

$$\begin{aligned}
&\Rightarrow \left[ 1 \left( \frac{1}{16}x^4 + \frac{1}{2}x^3 + \frac{3}{2}x^2 + 2x + 1 \right) \right] - \left[ 4 \left( \frac{x^3}{8} + \frac{3x^2}{4} + \frac{3x}{2} + 1 \right) \left( \frac{2}{x} \right) \right] \\
&\left[ 6 \left( \frac{x^2}{4} + x + 1 \right) \left( \frac{4}{x^2} \right) \right] - \left[ 4 \left( 1 + \frac{x}{2} \right) \left( \frac{8}{x^3} \right) \right] + \left[ 1 \left( \frac{16}{x^4} \right) \right] \\
&\Rightarrow \frac{1}{16}x^4 + \frac{1}{2}x^3 + \frac{3}{2}x^2 + 2x + 1 - x^2 - 6x - 12 - \frac{8}{x} + 6 + \frac{24}{x} + \frac{24}{x^2} \\
&- \frac{32}{x^3} - \frac{16}{x^2} + \frac{16}{x^4}
\end{aligned}$$

On rearranging

$$\text{Ans) } \frac{1}{16}x^4 + \frac{1}{2}x^3 + \frac{1}{2}x^2 - 4x - 5 + \frac{16}{x} + \frac{8}{x^2} - \frac{32}{x^3} + \frac{16}{x^4}$$

**Q. 12. Using binomial theorem, expand each of the following:**

$$(3x^2 - 2ax + 3a^2)^3$$

**Answer :** To find: Expansion of  $(3x^2 - 2ax + 3a^2)^3$

$$\text{Formula used: (i) } {}^nC_r = \frac{n!}{(n-r)!(r)!}$$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$\text{We have, } (3x^2 - 2ax + 3a^2)^3$$

$$\text{Let, } (3x^2 - 2ax) = p \dots (i)$$

$$\text{The equation becomes } (p + 3a^2)^3$$

$$\Rightarrow [{}^3C_0(p)^{3-0}] + [{}^3C_1(p)^{3-1}(3a^2)^1] + [{}^3C_2(p)^{3-2}(3a^2)^2] + [{}^3C_3(3a^2)^3]$$

$$\Rightarrow [{}^3C_0(p)^3] + [{}^3C_1(p)^2(3a^2)] + [{}^3C_2(p)(9a^4)] + [{}^3C_3(27a^6)]$$

Substituting the value of p from eqn. (i)

$$\Rightarrow \left[ \frac{3!}{0!(3-0)!} (3x^2 - 2ax)^3 \right] + \left[ \frac{3!}{1!(3-1)!} (3x^2 - 2ax)^2(3a^2) \right]$$

$$+ \left[ \frac{3!}{2!(3-2)!} (3x^2 - 2ax)(9a^4) \right] + \left[ \frac{3!}{3!(3-3)!} (27a^6) \right]$$

$$\Rightarrow [1(3x^2 - 2ax)^3] + [3(3x^2 - 2ax)^2(3a^2)] + [3(3x^2 - 2ax)(9a^4)] + [1(27a^6)^3]$$

...

(ii)

We need the value of  $p^3$  and  $p^2$ , where  $p = 3x^2 - 2ax$

For,  $(a+b)^3$ , we have formula  $a^3+b^3+3a^2b+3ab^2$

For,  $(3x^2 - 2ax)^3$ , substituting  $a = 3x^2$  and  $b = -2ax$  in the above formula

$$\Rightarrow [(3x^2)^3] + [(-2ax)^3] + [3(3x^2)^2(-2ax)] + [3(3x^2)(-2ax)^2]$$

$$\Rightarrow 27x^6 - 8a^3x^3 - 54ax^5 + 36a^2x^4 \dots \text{(iii)}$$

For,  $(a+b)^2$ , we have formula  $a^2+2ab+b^2$

For,  $(3x^2 - 2ax)^2$ , substituting  $a = 3x^2$  and  $b = -2ax$  in the above formula

$$\Rightarrow [(3x^2)^2] + [2(3x^2)(-2ax)] + [(-2ax)^2]$$

$$\Rightarrow 9x^4 - 12x^3a + 4a^2x^2 \dots \text{(iv)}$$

Putting the value obtained from eqn. (iii) and (iv) in eqn. (ii)

$$\Rightarrow [1(27x^6 - 8a^3x^3 - 54ax^5 + 36a^2x^4)] + [3(9x^4 - 12x^3a + 4a^2x^2)(3a^2)] + [3(3x^2 - 2ax)(9a^4)] + [1(27a^6)]$$

$$\Rightarrow 27x^6 - 8a^3x^3 - 54ax^5 + 36a^2x^4 + 81a^2x^4 - 108x^3a^3 + 36a^4x^2 + 81a^4x^2 - 54a^5x + 27a^6$$

On rearranging

$$\text{Ans) } 27x^6 - 54ax^5 + 117a^2x^4 - 116x^3a^3 + 117a^4x^2 - 54a^5x + 27a^6$$

**Q. 13. Evaluate :**

$$(\sqrt{2} + 1)^6 + (\sqrt{2} - 1)^6$$



**Answer :** To find: Value of  $(\sqrt{2}+1)^6 + (\sqrt{2}-1)^6$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$(a+1)^6 = [{}^6C_0a^6] + [{}^6C_1a^{6-1}1] + [{}^6C_2a^{6-2}1^2] + [{}^6C_3a^{6-3}1^3] + [{}^6C_4a^{6-4}1^4] + [{}^6C_5a^{6-5}1^5] + [{}^6C_61^6]$$

$$\Rightarrow {}^6C_0a^6 + {}^6C_1a^5 + {}^6C_2a^4 + {}^6C_3a^3 + {}^6C_4a^2 + {}^6C_5a + {}^6C_6 \dots (i)$$

$$(a-1)^6 =$$

$$[{}^6C_0a^6] + [{}^6C_1a^{6-1}(-1)^1] + [{}^6C_2a^{6-2}(-1)^2] + [{}^6C_3a^{6-3}(-1)^3] + [{}^6C_4a^{6-4}(-1)^4] + [{}^6C_5a^{6-5}(-1)^5] + [{}^6C_6(-1)^6]$$

$$\Rightarrow {}^6C_0a^6 - {}^6C_1a^5 + {}^6C_2a^4 - {}^6C_3a^3 + {}^6C_4a^2 - {}^6C_5a + {}^6C_6 \dots (ii)$$

Adding eqn. (i) and (ii)

$$(a+1)^6 + (a-1)^6 = [{}^6C_0a^6 + {}^6C_1a^5 + {}^6C_2a^4 + {}^6C_3a^3 + {}^6C_4a^2 + {}^6C_5a + {}^6C_6] + [{}^6C_0a^6 - {}^6C_1a^5 + {}^6C_2a^4 - {}^6C_3a^3 + {}^6C_4a^2 - {}^6C_5a + {}^6C_6]$$

$$\Rightarrow 2[{}^6C_0a^6 + {}^6C_2a^4 + {}^6C_4a^2 + {}^6C_6]$$

$$\Rightarrow 2\left[\left(\frac{6!}{0!(6-0)!} a^6\right) + \left(\frac{6!}{2!(6-2)!} a^4\right) + \left(\frac{6!}{4!(6-4)!} a^2\right) + \left(\frac{6!}{6!(6-6)!}\right)\right]$$

$$\Rightarrow 2[(1)a^6 + (15)a^4 + (15)a^2 + (1)]$$

$$\Rightarrow 2[a^6 + 15a^4 + 15a^2 + 1] = (a+1)^6 + (a-1)^6$$

Putting the value of  $a = \sqrt{2}$  in the above equation

$$(\sqrt{2}+1)^6 + (\sqrt{2}-1)^6 = 2[(\sqrt{2})^6 + 15(\sqrt{2})^4 + 15(\sqrt{2})^2 + 1]$$

$$\Rightarrow 2[8 + 15(4) + 15(2) + 1]$$

$$\Rightarrow 2[8 + 60 + 30 + 1]$$

$$\Rightarrow 2[99]$$

$$\Rightarrow 198$$

**Ans) 198**

**Q. 14. Evaluate :**

$$(\sqrt{3} + 1)^5 - (\sqrt{3} - 1)^5$$

**Answer :** To find: Value of  $(\sqrt{3} + 1)^5 - (\sqrt{3} - 1)^5$

Formula used: (I)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$(a+1)^5 = {}^5C_0a^5 + {}^5C_1a^{5-1}1 + {}^5C_2a^{5-2}1^2 + {}^5C_3a^{5-3}1^3 + {}^5C_4a^{5-4}1^4 + {}^5C_51^5$$

$$\Rightarrow {}^5C_0a^5 + {}^5C_1a^4 + {}^5C_2a^3 + {}^5C_3a^2 + {}^5C_4a + {}^5C_5 \dots (i)$$

$$(a-1)^5$$

$$= [{}^5C_0a^5] + [{}^5C_1a^{5-1}(-1)^1] + [{}^5C_2a^{5-2}(-1)^2] + [{}^5C_3a^{5-3}(-1)^3] + [{}^5C_4a^{5-4}(-1)^4] + [{}^5C_5(-1)^5]$$

$$\Rightarrow {}^5C_0a^5 - {}^5C_1a^4 + {}^5C_2a^3 - {}^5C_3a^2 + {}^5C_4a - {}^5C_5 \dots (ii)$$

Subtracting (ii) from (i)

$$(a+1)^5 - (a-1)^5 = [{}^5C_0a^5 + {}^5C_1a^4 + {}^5C_2a^3 + {}^5C_3a^2 + {}^5C_4a + {}^5C_5] - [{}^5C_0a^5 - {}^5C_1a^4 + {}^5C_2a^3 - {}^5C_3a^2 + {}^5C_4a - {}^5C_5]$$

$$\Rightarrow 2[{}^5C_1a^4 + {}^5C_3a^2 + {}^5C_5]$$

$$\Rightarrow 2 \left[ \left( \frac{5!}{1!(5-1)!} a^4 \right) + \left( \frac{5!}{3!(5-3)!} a^2 \right) + \left( \frac{5!}{5!(5-5)!} \right) \right]$$

$$\Rightarrow 2[(5)a^4 + (10)a^2 + (1)]$$

$$\Rightarrow 2[5a^4 + 10a^2 + 1] = (a+1)^5 - (a-1)^5$$

Putting the value of  $a = \sqrt{3}$  in the above equation

$$(\sqrt{3}+1)^5 - (\sqrt{3}-1)^5 = 2[5(\sqrt{3})^4 + 10(\sqrt{3})^2 + 1]$$

$$\Rightarrow 2[(5)(9) + (10)(3) + 1]$$

$$\Rightarrow 2[45+30+1]$$

$$\Rightarrow 152$$

**Ans) 152**

**Q. 15. Evaluate :**

$$(2+\sqrt{3})^7 + (2-\sqrt{3})^7$$

**Answer:** To find: Value of  $(2+\sqrt{3})^7 + (2-\sqrt{3})^7$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$(a+b)^7 = [{}^7C_0a^7] + [{}^7C_1a^{7-1}b] + [{}^7C_2a^{7-2}b^2] + [{}^7C_3a^{7-3}b^3] + [{}^7C_4a^{7-4}b^4] + [{}^7C_5a^{7-5}b^5] + [{}^7C_6a^{7-6}b^6] + [{}^7C_7b^7]$$

$$\Rightarrow {}^7C_0a^7 + {}^7C_1a^6b + {}^7C_2a^5b^2 + {}^7C_3a^4b^3 + {}^7C_4a^3b^4 + {}^7C_5a^2b^5 + {}^7C_6a^1b^6 + {}^7C_7b^7 \dots (i)$$

$$(a-b)^7 = [{}^7C_0a^7] + [{}^7C_1a^{7-1}(-b)] + [{}^7C_2a^{7-2}(-b)^2] + [{}^7C_3a^{7-3}(-b)^3] + [{}^7C_4a^{7-4}(-b)^4] + [{}^7C_5a^{7-5}(-b)^5] + [{}^7C_6a^{7-6}(-b)^6] + [{}^7C_7(-b)^7]$$

$$\Rightarrow {}^7C_0a^7 - {}^7C_1a^6b + {}^7C_2a^5b^2 - {}^7C_3a^4b^3 + {}^7C_4a^3b^4 - {}^7C_5a^2b^5 + {}^7C_6a^1b^6 - {}^7C_7b^7 \dots (ii)$$

Adding eqn. (i) and (ii)

$$(a+b)^7 + (a-b)^7 = [{}^7C_0a^7 + {}^7C_1a^6b + {}^7C_2a^5b^2 + {}^7C_3a^4b^3 + {}^7C_4a^3b^4 + {}^7C_5a^2b^5 + {}^7C_6a^1b^6 + {}^7C_7b^7] + [{}^7C_0a^7 - {}^7C_1a^6b + {}^7C_2a^5b^2 - {}^7C_3a^4b^3 + {}^7C_4a^3b^4 - {}^7C_5a^2b^5 + {}^7C_6a^1b^6 - {}^7C_7b^7]$$

$$\Rightarrow 2[{}^7C_0a^7 + {}^7C_2a^5b^2 + {}^7C_4a^3b^4 + {}^7C_6a^1b^6]$$

$$\Rightarrow 2\left[\left[\frac{7!}{0!(7-0)!}a^7\right] + \left[\frac{7!}{2!(7-2)!}a^5b^2\right] + \left[\frac{7!}{4!(7-4)!}a^3b^4\right] + \left[\frac{7!}{6!(7-6)!}a^1b^6\right]\right]$$

$$\Rightarrow 2[(1)a^7 + (21)a^5b^2 + (35)a^3b^4 + (7)ab^6]$$

$$\Rightarrow 2[a^7 + 21a^5b^2 + 35a^3b^4 + 7ab^6] = (a+b)^7 + (a-b)^7$$

Putting the value of  $a = 2$  and  $b = \sqrt{3}$  in the above equation

$$(2+\sqrt{3})^7 + (2-\sqrt{3})^7$$

$$= 2\left[\{2^7\} + \{21(2)^5(\sqrt{3})^2\} + \{35(2)^3(\sqrt{3})^4\} + \{7(2)(\sqrt{3})^6\}\right]$$

$$= 2[128 + 21(32)(3) + 35(8)(9) + 7(2)(27)]$$

$$= 2[128 + 2016 + 2520 + 378]$$

$$= 10084$$

**Ans)** 10084

**Q. 16. Evaluate :**

$$(\sqrt{3} + \sqrt{2})^6 - (\sqrt{3} - \sqrt{2})^6$$

**Answer :** To find: Value of  $(\sqrt{3} + \sqrt{2})^6 - (\sqrt{3} - \sqrt{2})^6$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$(a+b)^6 = {}^6C_0a^6 + {}^6C_1a^{6-1}b + {}^6C_2a^{6-2}b^2 + {}^6C_3a^{6-3}b^3 + {}^6C_4a^{6-4}b^4 + {}^6C_5a^{6-5}b^5 + {}^6C_6b^6$$

$$\Rightarrow {}^6C_0a^6 + {}^6C_1a^5b + {}^6C_2a^4b^2 + {}^6C_3a^3b^3 + {}^6C_4a^2b^4 + {}^6C_5ab^5 + {}^6C_6b^6 \dots (i)$$

$$(a-b)^6 =$$

$$= [{}^6C_0a^6] + [{}^6C_1a^{6-1}(-b)] + [{}^6C_2a^{6-2}(-b)^2] + [{}^6C_3a^{6-3}(-b)^3] + [{}^6C_4a^{6-4}(-b)^4] + [{}^6C_5a^{6-5}(-b)^5] + [{}^6C_6(-b)^6]$$

$$\Rightarrow {}^6C_0a^6 - {}^6C_1a^5b + {}^6C_2a^4b^2 - {}^6C_3a^3b^3 + {}^6C_4a^2b^4 - {}^6C_5ab^5 + {}^6C_6b^6 \dots (ii)$$

Subtracting (ii) from (i)

$$(a+b)^6 - (a-b)^6 = [{}^6C_0a^6 + {}^6C_1a^5b + {}^6C_2a^4b^2 + {}^6C_3a^3b^3 + {}^6C_4a^2b^4 + {}^6C_5ab^5 + {}^6C_6b^6] - [{}^6C_0a^6 - {}^6C_1a^5b + {}^6C_2a^4b^2 - {}^6C_3a^3b^3 + {}^6C_4a^2b^4 - {}^6C_5ab^5 + {}^6C_6b^6]$$

$$= 2[{}^6C_1a^5b + {}^6C_3a^3b^3 + {}^6C_5ab^5]$$

$$= 2 \left[ \left\{ \frac{6!}{1!(6-1)!} a^5 a \right\} + \left\{ \frac{6!}{3!(6-3)!} a^3 b^3 \right\} + \left\{ \frac{6!}{5!(6-5)!} ab^5 \right\} \right]$$

$$= 2[(6)a^5b + (20)a^3b^3 + (6)ab^5]$$

$$\Rightarrow (a+b)^6 - (a-b)^6 = 2[(6)a^5b + (20)a^3b^3 + (6)ab^5]$$

Putting the value of  $a = \sqrt{3}$  and  $b = \sqrt{2}$  in the above equation

$$(\sqrt{3} + \sqrt{2})^6 - (\sqrt{3} - \sqrt{2})^6$$

$$\Rightarrow 2[(6)(\sqrt{3})^5(\sqrt{2}) + (20)(\sqrt{3})^3(\sqrt{2})^3 + (6)(\sqrt{3})(\sqrt{2})^5]$$

$$\Rightarrow 2[54(\sqrt{6}) + 120(\sqrt{6}) + 24(\sqrt{6})]$$

$$\Rightarrow 396\sqrt{6}$$

**Ans)**  $396\sqrt{6}$

Q. 17. Prove that

$$\sum_{r=0}^n {}^nC_r \cdot 3^r = 4^n$$

**Answer :**

$$\sum_{r=0}^n {}^nC_r \cdot 3^r = 4^n$$

To prove:

$$\sum_{r=0}^n {}^nC_r \cdot a^{n-r} b^r = (a+b)^n$$

Formula used:

Proof: In the above formula if we put  $a = 1$  and  $b = 3$ , then we will get

$$\sum_{r=0}^n {}^nC_r \cdot 1^{n-r} 3^r = (1+3)^n$$

Therefore,

$$\sum_{r=0}^n {}^nC_r \cdot 3^r = (4)^n$$

Hence Proved.

**Q. 18. Using binomial theorem, evaluate each of the following :**

(i)  $(101)^4$  (ii)  $(98)^4$   
(iii)  $(1.2)^4$

**Answer :** (i)  $(101)^4$

To find: Value of  $(101)^4$

$$\text{Formula used: (i) } {}^nC_r = \frac{n!}{(n-r)!(r)!}$$

$$(ii) (a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1} b + {}^nC_2 a^{n-2} b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$$

$$101 = (100+1)$$

$$\text{Now } (101)^4 = (100+1)^4$$

$$\begin{aligned} (100+1)^4 = & [{}^4C_0 (100)^{4-0}] + [{}^4C_1 (100)^{4-1} (1)^1] + [{}^4C_2 (100)^{4-2} (1)^2] + \\ & [{}^4C_3 (100)^{4-3} (1)^3] + [{}^4C_4 (1)^4] \end{aligned}$$

$$\Rightarrow [{}^4C_0(100)^4] + [{}^4C_1(100)^3(1)^1] + [{}^4C_2(100)^2(1)^2] + [{}^4C_3(100)^1(1)^3] + [{}^4C_4(1)^4]$$

$$\Rightarrow \left[ \frac{4!}{0!(4-0)!} (100000000) \right] + \left[ \frac{4!}{1!(4-1)!} (1000000) \right] + \left[ \frac{4!}{2!(4-2)!} (10000) \right] + \left[ \frac{4!}{3!(4-3)!} (100) \right] + \left[ \frac{4!}{4!(4-4)!} (1) \right]$$

$$\Rightarrow [(1)(100000000)] + [(4)(1000000)] + [(6)(10000)] + [(4)(100)] + [(1)(1)]$$

$$= 104060401$$

**Ans)** 104060401

**(ii)**  $(98)^4$

To find: Value of  $(98)^4$

Formula used: (I)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$\text{(ii)} (a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$$

$$98 = (100-2)$$

$$\text{Now } (98)^4 = (100-2)^4$$

$$\begin{aligned} & (100-2)^4 \\ &= [{}^4C_0(100)^{4-0}] + [{}^4C_1(100)^{4-1}(-2)^1] + [{}^4C_2(100)^{4-2}(-2)^2] + [{}^4C_3(100)^{4-3}(-2)^3] + [{}^4C_4(-2)^4] \end{aligned}$$

$$\Rightarrow [{}^4C_0(100)^4] - [{}^4C_1(100)^3(2)] + [{}^4C_2(100)^2(4)] - [{}^4C_3(100)^1(8)] + [{}^4C_4(16)]$$

$$\Rightarrow \left[ \frac{4!}{0!(4-0)!} (100000000) \right] - \left[ \frac{4!}{1!(4-1)!} (1000000)(2) \right] + \left[ \frac{4!}{2!(4-2)!} (10000)(4) \right] - \left[ \frac{4!}{3!(4-3)!} (100)(8) \right] + \left[ \frac{4!}{4!(4-4)!} (16) \right]$$

$$\Rightarrow [(1)(100000000)] - [(4)(1000000)(2)] + [(6)(10000)(4)] - [(4)(100)(8)] + [(1)(16)]$$

$$= 92236816$$

**Ans)** 92236816

**(iii)**  $(1.2)^4$

To find: Value of  $(1.2)^4$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $(a+b)^n = {}^nC_0a^n + {}^nC_1a^{n-1}b + {}^nC_2a^{n-2}b^2 + \dots + {}^nC_{n-1}ab^{n-1} + {}^nC_nb^n$

$$1.2 = (1 + 0.2)$$

Now  $(1.2)^4 = (1 + 0.2)^4$

$$\begin{aligned} & (1+0.2)^4 \\ &= [{}^4C_0(1)^{4-0}] + [{}^4C_1(1)^{4-1}(0.2)^1] + [{}^4C_2(1)^{4-2}(0.2)^2] + \\ & [{}^4C_3(1)^{4-3}(0.2)^3] + [{}^4C_4(0.2)^4] \end{aligned}$$

$$\Rightarrow [{}^4C_0(1)^4] + [{}^4C_1(1)^3(0.2)^1] + [{}^4C_2(1)^2(0.2)^2] + [{}^4C_3(1)^1(0.2)^3] + [{}^4C_4(0.2)^4]$$

$$\begin{aligned} & \Rightarrow \left[ \frac{4!}{0!(4-0)!} (1) \right] + \left[ \frac{4!}{1!(4-1)!} (1)(0.2) \right] + \left[ \frac{4!}{2!(4-2)!} (1)(0.04) \right] + \\ & \left[ \frac{4!}{3!(4-3)!} (1)(0.008) \right] + \left[ \frac{4!}{4!(4-4)!} (0.0016) \right] \end{aligned}$$

$$\Rightarrow [(1)(1)] + [(4)(1)(0.2)] + [(6)(1)(0.04)] + [(4)(1)(0.008)] + [(1)(0.0016)]$$

$$= 2.0736$$

**Ans)** 2.0736

**Q. 19.** Using binomial theorem, prove that  $(2^{3n} - 7n - 1)$  is divisible by 49, where  $n \in \mathbb{N}$ .



**Answer :** To prove:  $(2^{3n} - 7n - 1)$  is divisible by 49, where  $n \in \mathbb{N}$

Formula used:  $(a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$

$$(2^{3n} - 7n - 1) = (2^3)^n - 7n - 1$$

$$\Rightarrow 8^n - 7n - 1$$

$$\Rightarrow (1+7)^n - 7n - 1$$

$$\Rightarrow {}^nC_0 1^n + {}^nC_1 1^{n-1}7 + {}^nC_2 1^{n-2}7^2 + \dots + {}^nC_{n-1} 1^{n-1}7 + {}^nC_n 7^n - 7n - 1$$

$$\Rightarrow {}^nC_0 + {}^nC_1 7 + {}^nC_2 7^2 + \dots + {}^nC_{n-1} 7^{n-1} + {}^nC_n 7^n - 7n - 1$$

$$\Rightarrow 1 + 7n + 7^2[{}^nC_2 + {}^nC_3 7 + \dots + {}^nC_{n-1} 7^{n-3} + {}^nC_n 7^{n-2}] - 7n - 1$$

$$\Rightarrow 7^2[{}^nC_2 + {}^nC_3 7 + \dots + {}^nC_{n-1} 7^{n-3} + {}^nC_n 7^{n-2}]$$

$$\Rightarrow 49[{}^nC_2 + {}^nC_3 7 + \dots + {}^nC_{n-1} 7^{n-3} + {}^nC_n 7^{n-2}]$$

$$\Rightarrow 49K, \text{ where } K = ({}^nC_2 + {}^nC_3 7 + \dots + {}^nC_{n-1} 7^{n-3} + {}^nC_n 7^{n-2})$$

$$\text{Now, } (2^{3n} - 7n - 1) = 49K$$

Therefore  $(2^{3n} - 7n - 1)$  is divisible by 49

**Q. 20. Prove that** 
$$(2 + \sqrt{x})^4 + (2 - \sqrt{x})^4 = 2(16 + 24x + x^2)$$

**Answer :** To prove: 
$$(2 + \sqrt{x})^4 + (2 - \sqrt{x})^4 = 2(16 + 24x + x^2)$$

Formula used: (i) 
$${}^nC_r = \frac{n!}{(n-r)!(r)!}$$

(ii) 
$$(a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$$

$$(a+b)^4 = {}^4C_0 a^4 + {}^4C_1 a^3 b + {}^4C_2 a^2 b^2 + {}^4C_3 a b^3 + {}^4C_4 b^4$$

$$\Rightarrow {}^4C_0 a^4 + {}^4C_1 a^3 b + {}^4C_2 a^2 b^2 + {}^4C_3 a b^3 + {}^4C_4 b^4 \dots \text{ (i)}$$

$$(a-b)^4 = {}^4C_0 a^4 + {}^4C_1 a^3 (-b) + {}^4C_2 a^2 (-b)^2 + {}^4C_3 a (-b)^3 + {}^4C_4 (-b)^4$$

$$\Rightarrow {}^4C_0 a^4 - {}^4C_1 a^3 b + {}^4C_2 a^2 b^2 - {}^4C_3 a b^3 + {}^4C_4 b^4 \dots \text{ (ii)}$$

Adding (i) and (ii)

$$(a+b)^4 + (a-b)^4 = [{}^4C_0a^4 + {}^4C_1a^3b + {}^4C_2a^2b^2 + {}^4C_3a^1b^3 + {}^4C_4b^4] + [{}^4C_0a^4 - {}^4C_1a^3b + {}^4C_2a^2b^2 - {}^4C_3ab^3 + {}^4C_4b^4]$$

$$\Rightarrow 2[{}^4C_0a^4 + {}^4C_2a^2b^2 + {}^4C_4b^4]$$

$$\Rightarrow 2\left[\left(\frac{4!}{0!(4-0)!}a^4\right) + \left(\frac{4!}{2!(4-2)!}a^2b^2\right) + \left(\frac{4!}{4!(4-4)!}b^4\right)\right]$$

$$\Rightarrow 2[(1)a^4 + (6)a^2b^2 + (1)b^4]$$

$$\Rightarrow 2[a^4 + 6a^2b^2 + b^4]$$

$$\text{Therefore, } (a+b)^4 + (a-b)^4 = 2[a^4 + 6a^2b^2 + b^4]$$

Now, putting  $a = 2$  and  $b = (\sqrt{x})$  in the above equation.

$$(2+\sqrt{x})^4 + (2-\sqrt{x})^4 = 2[(2)^4 + 6(2)^2(\sqrt{x})^2 + (\sqrt{x})^4]$$

$$= 2(16+24x+x^2)$$

Hence proved.

**Q. 21. Find the 7<sup>th</sup> term in the expansion of  $\left(\frac{4x}{5} + \frac{5}{2x}\right)^8$ .**

**Answer :** To find: 7<sup>th</sup> term in the expansion of  $\left(\frac{4x}{5} + \frac{5}{2x}\right)^8$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $T_{r+1} = {}^nC_r a^{n-r} b^r$

For 7<sup>th</sup> term,  $r+1=7$

$$\Rightarrow r = 6$$

$$\text{In, } \left(\frac{4x}{5} + \frac{5}{2x}\right)^8$$

$$7^{\text{th}} \text{ term} = T_{6+1}$$

$$\Rightarrow {}^8C_6 \left(\frac{4x}{5}\right)^{8-6} \left(\frac{5}{2x}\right)^6$$

$$\Rightarrow \frac{8!}{6!(8-6)!} \left(\frac{4x}{5}\right)^2 \left(\frac{5}{2x}\right)^6$$

$$\Rightarrow (28) \left(\frac{16x^2}{25}\right) \left(\frac{15625}{64x^6}\right)$$

$$\Rightarrow \frac{4375}{x^4}$$

$$\text{Ans) } \frac{4375}{x^4}$$

**Q. 22. Find the 9<sup>th</sup> term in the expansion of  $\left(\frac{a}{b} - \frac{b}{2a^2}\right)^{12}$ .**

**Answer :** To find: 9<sup>th</sup> term in the expansion of  $\left(\frac{a}{b} - \frac{b}{2a^2}\right)^{12}$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

$$(ii) T_{r+1} = {}^nC_r a^{n-r} b^r$$

For 9<sup>th</sup> term,  $r+1=9$

$$\Rightarrow r = 8$$

$$\text{In, } \left(\frac{a}{b} - \frac{b}{2a^2}\right)^{12}$$

$$9^{\text{th}} \text{ term} = T_{8+1}$$

$$\Rightarrow {}^{12}C_8 \left(\frac{a}{b}\right)^{12-8} \left(\frac{-b}{2a^2}\right)^8$$

$$\Rightarrow \frac{12!}{8!(12-8)!} \left(\frac{a}{b}\right)^4 \left(\frac{-b}{2a^2}\right)^8$$

$$\Rightarrow 495 \left( \frac{a^4}{b^4} \right) \left( \frac{b^8}{256a^{16}} \right)$$

$$\Rightarrow \left( \frac{495b^4}{256a^{12}} \right)$$

$$\text{Ans) } \left( \frac{495b^4}{256a^{12}} \right)$$

**Q. 23. Find the 16<sup>th</sup> term in the expansion of  $(\sqrt{x} - \sqrt{y})^{17}$ .**

**Answer :** To find: 16<sup>th</sup> term in the expansion of  $(\sqrt{x} - \sqrt{y})^{17}$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $T_{r+1} = {}^nC_r a^{n-r} b^r$

For 16<sup>th</sup> term,  $r+1=16$

$$\Rightarrow r = 15$$

$$\text{In, } (\sqrt{x} - \sqrt{y})^{17}$$

$$16^{\text{th}} \text{ term} = T_{15+1}$$

$$\Rightarrow {}^{17}C_{15} (\sqrt{x})^{17-15} (-\sqrt{y})^{15}$$

$$\Rightarrow \frac{17!}{15!(17-15)!} (\sqrt{x})^2 (-\sqrt{y})^{15}$$

$$\Rightarrow 136(x)(-y)^{\frac{15}{2}}$$

$$\Rightarrow -136x y^{\frac{15}{2}}$$

$$\text{Ans) } -136 y^{\frac{15}{2}}$$

**Q. 24. Find the 13<sup>th</sup> term in the expansion of  $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}$ ,  $x \neq 0$ .**

**Answer :** To find: 13<sup>th</sup> term in the expansion of  $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}$

Formula used: (i)  ${}^nC_r = \frac{n!}{(n-r)!(r)!}$

(ii)  $T_{r+1} = {}^nC_r a^{n-r} b^r$

For 13<sup>th</sup> term,  $r+1=13$

$$\Rightarrow r = 12$$

$$\text{In, } \left(9x - \frac{1}{3\sqrt{x}}\right)^{18}$$

$$13^{\text{th}} \text{ term} = T_{12+1}$$

$$\Rightarrow {}^{18}C_{12} (9x)^{18-12} \left(-\frac{1}{3\sqrt{x}}\right)^{12}$$

$$\Rightarrow \frac{18!}{12!(18-12)!} (9x)^6 \left(-\frac{1}{3\sqrt{x}}\right)^{12}$$

$$\Rightarrow 18564 (531441 x^6) \left(\frac{1}{531441 x^6}\right)$$

$$\Rightarrow 18564$$

**Q. 25. Find the coefficients of  $x^7$  and  $x^8$  in the expansion of  $\left(2 + \frac{x}{3}\right)^n$ .**

**Answer :** To find : coefficients of  $x^7$  and  $x^8$

Formula :  $t_{r+1} = {}^nC_r a^{n-r} b^r$

Here,  $a=2$ ,  $b = \frac{x}{3}$

We have,  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

$$\begin{aligned}\therefore t_{r+1} &= \binom{n}{r} (2)^{n-r} \left(\frac{x}{3}\right)^r \\ &= \binom{n}{r} \frac{2^{n-r}}{3^r} x^r\end{aligned}$$

To get a coefficient of  $x^7$ , we must have,

$$x^7 = x^r$$

$$\bullet r = 7$$

Therefore, the coefficient of  $x^7 = \binom{n}{7} \frac{2^{n-7}}{3^7}$

And to get the coefficient of  $x^8$  we must have,

$$x^8 = x^r$$

$$\bullet r = 8$$

Therefore, the coefficient of  $x^8 = \binom{n}{8} \frac{2^{n-8}}{3^8}$

Conclusion :

$$\bullet \text{Coefficient of } x^7 = \binom{n}{7} \frac{2^{n-7}}{3^7}$$

$$\bullet \text{Coefficient of } x^8 = \binom{n}{8} \frac{2^{n-8}}{3^8}$$

**Q. 26. Find the ratio of the coefficient of  $x^{15}$  to the term independent of  $x$  in the**

**expansion of  $\left(x^2 + \frac{2}{x}\right)^{15}$ .**

**Answer :** To Find: the ratio of the coefficient of  $x^{15}$  to the term independent of  $x$

Formula :  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

Here,  $a=x^2$ ,  $b = \frac{2}{x}$  and  $n=15$

We have a formula,

$$\begin{aligned}t_{r+1} &= \binom{n}{r} a^{n-r} b^r \\&= \binom{15}{r} (x^2)^{15-r} \left(\frac{2}{x}\right)^r \\&= \binom{15}{r} (x)^{30-2r} (2)^r (x)^{-r} \\&= \binom{15}{r} (x)^{30-2r-r} (2)^r \\&= \binom{15}{r} (2)^r (x)^{30-3r}\end{aligned}$$

To get coefficient of  $x^{15}$  we must have,

$$(x)^{30-3r} = x^{15}$$

$$\bullet 30 - 3r = 15$$

$$\bullet 3r = 15$$

$$\bullet r = 5$$

Therefore, coefficient of  $x^{15} = \binom{15}{5} (2)^5$

Now, to get coefficient of term independent of  $x$  that is coefficient of  $x^0$  we must have,

$$(x)^{30-3r} = x^0$$

$$\bullet 30 - 3r = 0$$

$$\bullet 3r = 30$$

$$\bullet r = 10$$

Therefore, coefficient of  $x^0 = \binom{15}{10} (2)^{10}$

But  $\binom{15}{10} = \binom{15}{5}$  .....  $[\because \binom{n}{r} = \binom{n}{n-r}]$

Therefore, the coefficient of  $x^0 = \binom{15}{5} (2)^{10}$

Therefore,

$$\frac{\text{coefficient of } x^{15}}{\text{coefficient of } x^0} = \frac{\binom{15}{5} (2)^5}{\binom{15}{5} (2)^{10}}$$

$$= \frac{1}{(2)^5}$$

$$= \frac{1}{32}$$

Hence, coefficient of  $x^{15}$  : coefficient of  $x^0 = 1:32$

Conclusion : The ratio of coefficient of  $x^{15}$  to coefficient of  $x^0 = 1:32$

**Q. 27. Show that the ratio of the coefficient of  $x^{10}$  in the expansion of  $(1 - x^2)^{10}$  and**

**the term independent of  $x$  in the expansion of  $\left(x - \frac{2}{x}\right)^{10}$  is 1 : 32.**

**Answer : To Prove** : coefficient of  $x^{10}$  in  $(1-x^2)^{10}$  : coefficient of  $x^0$  in  $\left(x - \frac{2}{x}\right)^{10} = 1:32$

For  $(1-x^2)^{10}$ ,

Here,  $a=1$ ,  $b=-x^2$  and  $n=15$

We have formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{10}{r} (1)^{10-r} (-x^2)^r$$



$$= - \binom{10}{r} (1) (x)^{2r}$$

To get coefficient of  $x^{10}$  we must have,

$$(x)^{2r} = x^{10}$$

$$\bullet 2r = 10$$

$$\bullet r = 5$$

Therefore, coefficient of  $x^{10} = - \binom{10}{5}$

$$\text{For } \left(x - \frac{2}{x}\right)^{10},$$

Here,  $a=x$ ,  $b = \frac{-2}{x}$  and  $n=10$

We have a formula,

$$\begin{aligned} t_{r+1} &= \binom{n}{r} a^{n-r} b^r \\ &= \binom{10}{r} (x)^{10-r} \left(\frac{-2}{x}\right)^r \\ &= \binom{10}{r} (x)^{10-r} (-2)^r (x)^{-r} \\ &= \binom{10}{r} (x)^{10-r-r} (-2)^r \\ &= \binom{10}{r} (-2)^r (x)^{10-2r} \end{aligned}$$

Now, to get coefficient of term independent of  $x$  that is coefficient of  $x^0$  we must have,

$$(x)^{10-2r} = x^0$$

$$\bullet 10 - 2r = 0$$

$$\bullet 2r = 10$$

- $r = 5$

Therefore, coefficient of  $x^0 = -\binom{10}{5} (2)^5$

Therefore,

$$\frac{\text{coefficient of } x^{10} \text{ in } (1 - x^2)^{10}}{\text{coefficient of } x^0 \text{ in } \left(x - \frac{2}{x}\right)^{10}} = \frac{-\binom{15}{5}}{-\binom{15}{5} (2)^5}$$

$$= \frac{1}{(2)^5}$$

$$= \frac{1}{32}$$

Hence,

Coefficient of  $x^{10}$  in  $(1-x^2)^{10}$ : coefficient of  $x^0$  in  $\left(x - \frac{2}{x}\right)^{10} = 1:32$

**Q. 28. Find the term independent of  $x$  in the expansion of  $(91 + x + 2x^3)$**

$$\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9.$$

**Answer:** To Find : term independent of  $x$ , i.e. coefficient of  $x^0$

Formula:  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, the expansion of  $\left(x - \frac{2}{x}\right)^{10}$  is given by,

$$\begin{aligned}
\left(x - \frac{2}{x}\right)^{10} &= \sum_{r=0}^{10} \binom{10}{r} (x)^{10-r} \left(\frac{-2}{x}\right)^r \\
&= \binom{10}{0} (x)^{10} \left(\frac{-2}{x}\right)^0 + \binom{10}{1} (x)^9 \left(\frac{-2}{x}\right)^1 + \binom{10}{2} (x)^8 \left(\frac{-2}{x}\right)^2 + \dots \dots \dots \\
&\quad + \binom{10}{10} (x)^0 \left(\frac{-2}{x}\right)^{10} \\
&= x^{10} + \binom{10}{1} (x)^9 (-2) \frac{1}{x} + \binom{10}{2} (x)^8 (-2)^2 \frac{1}{x^2} + \dots + \binom{10}{10} (x)^0 (-2)^{10} \frac{1}{x^{10}} \\
&= x^{10} - (2) \binom{10}{1} (x)^8 + (2)^2 \binom{10}{2} (x)^6 + \dots \dots \dots + (2)^{10} \binom{10}{10} \frac{1}{x^{10}}
\end{aligned}$$

Now,

$$\begin{aligned}
(91 + x + 2x^3) \left(x - \frac{2}{x}\right)^{10} \\
= (91 + x + 2x^3) \left( x^{10} - (2) \binom{10}{1} (x)^8 + (2)^2 \binom{10}{2} (x)^6 + \dots \dots \dots \right. \\
\quad \left. + (2)^{10} \binom{10}{10} \frac{1}{x^{10}} \right)
\end{aligned}$$

Multiplying the second bracket by 91, x and 2x<sup>3</sup>

$$\begin{aligned}
&= \left\{ 91x^{10} - 91(2) \binom{10}{1} (x)^8 + 91(2)^2 \binom{10}{2} (x)^6 + \dots + 91(2)^{10} \binom{10}{10} \frac{1}{x^{10}} \right\} \\
&\quad + \left\{ x \cdot x^{10} - x \cdot (2) \binom{10}{1} (x)^8 + x \cdot (2)^2 \binom{10}{2} (x)^6 + \dots \dots \dots \right. \\
&\quad \left. + x \cdot (2)^{10} \binom{10}{10} \frac{1}{x^{10}} \right\} \\
&\quad + \left\{ 2x^3 \cdot x^{10} - 2x^3 \cdot (2) \binom{10}{1} (x)^8 + 2x^3 \cdot (2)^2 \binom{10}{2} (x)^6 + \dots \dots \dots \right. \\
&\quad \left. + 2x^3 \cdot (2)^{10} \binom{10}{10} \frac{1}{x^{10}} \right\}
\end{aligned}$$

In the first bracket, there will be a 6<sup>th</sup> term of x<sup>0</sup> having coefficient  $91(-2)^5 \binom{10}{5}$

While in the second and third bracket, the constant term is absent.

Therefore, the coefficient of term independent of x, i.e. constant term in the above expansion

$$= 91(-2)^5 \binom{10}{5}$$

$$= -91 \cdot (2)^5 \frac{10 \times 9 \times 8 \times 7 \times 6}{5 \times 4 \times 3 \times 2 \times 1}$$

$$= -91(2)^5 (252)$$

Conclusion: coefficient of term independent of x =  $-91(2)^5 (252)$

**Q. 29. Find the coefficient of x in the expansion of  $(1 - 3x + 7x^2)(1 - x)^{16}$ .**

**Answer :** To Find : coefficient of x

Formula :  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, expansion of  $(1-x)^{16}$  is given by,

$$\begin{aligned} (1-x)^{16} &= \sum_{r=0}^{16} \binom{16}{r} (1)^{16-r} (-x)^r \\ &= \binom{16}{0} (1)^{16} (-x)^0 + \binom{16}{1} (1)^{15} (-x)^1 + \binom{16}{2} (1)^{14} (-x)^2 + \dots \dots \dots \\ &\quad + \binom{16}{16} (1)^0 (-x)^{16} \\ &= 1 - \binom{16}{1} x + \binom{16}{2} x^2 + \dots \dots \dots + \binom{16}{16} x^{16} \end{aligned}$$

Now,

$$(1 - 3x + 7x^2)(1 - x)^{16}$$

$$= (1 - 3x + 7x^2) \left( 1 - \binom{16}{1}x + \binom{16}{2}x^2 + \dots + \binom{16}{16}x^{16} \right)$$

Multiplying the second bracket by 1, (-3x) and 7x<sup>2</sup>

$$= \left( 1 - \binom{16}{1}x + \binom{16}{2}x^2 + \dots + \binom{16}{16}x^{16} \right)$$

$$+ \left( -3x + 3x \binom{16}{1}x - 3x \binom{16}{2}x^2 + \dots - 3x \binom{16}{16}x^{16} \right)$$

$$+ \left( 7x^2 - 7x^2 \binom{16}{1}x + 7x^2 \binom{16}{2}x^2 + \dots + 7x^2 \binom{16}{16}x^{16} \right)$$

In the above equation terms containing x are

$$-\binom{16}{1}x \text{ and } -3x$$

Therefore, the coefficient of x in the above expansion

$$= -\binom{16}{1} - 3$$

$$= -16 - 3$$

$$= -19$$

Conclusion: coefficient of x = -19

**Q. 30. Find the coefficient of**

**(i) x<sup>5</sup> in the expansion of (x + 3)<sup>8</sup>**

**(ii) x<sup>6</sup> in the expansion of  $\left( 3x^2 - \frac{1}{3x} \right)^9$ .**

**(iii) x<sup>-15</sup> in the expansion of  $\left( 3x^2 - \frac{a}{3x^3} \right)^{10}$ .**

**(iv) a<sup>7</sup>b<sup>5</sup> in the expansion of (a - 2b)<sup>12</sup>.**

**Answer : (i)** Here, a=x, b=3 and n=8

We have a formula,

$$\begin{aligned} t_{r+1} &= \binom{n}{r} a^{n-r} b^r \\ &= \binom{8}{r} (x)^{8-r} (3)^r \\ &= \binom{8}{r} (3)^r (x)^{8-r} \end{aligned}$$

To get coefficient of  $x^5$  we must have,

$$(x)^{8-r} = x^5$$

$$\bullet 8 - r = 5$$

$$\bullet r = 3$$

Therefore, coefficient of  $x^5 = \binom{8}{3} (3)^3$

$$= \frac{8 \times 7 \times 6}{3 \times 2 \times 1} \cdot (27)$$

$$= 1512$$

(ii) Here,  $a=3x^2$ ,  $b = \frac{-1}{3x}$  and  $n=9$

We have a formula,

$$\begin{aligned} t_{r+1} &= \binom{n}{r} a^{n-r} b^r \\ &= \binom{9}{r} (3x^2)^{9-r} \left(\frac{-1}{3x}\right)^r \\ &= \binom{9}{r} (3)^{9-r} (x^2)^{9-r} \left(\frac{-1}{3}\right)^r (x)^{-r} \\ &= \binom{9}{r} (3)^{9-r} (x)^{18-2r} \left(\frac{-1}{3}\right)^r (x)^{-r} \end{aligned}$$

$$= \binom{9}{r} (3)^{9-r} (x)^{18-2r-r} \left(\frac{-1}{3}\right)^r$$

$$= \binom{9}{r} (3)^{9-r} \left(\frac{-1}{3}\right)^r (x)^{18-3r}$$

To get coefficient of  $x^6$  we must have,

$$(x)^{18-3r} = x^6$$

$$\bullet 18 - 3r = 6$$

$$\bullet 3r = 12$$

$$\bullet r = 4$$

$$\text{Therefore, coefficient of } x^6 = \binom{9}{4} (3)^{9-4} \left(\frac{-1}{3}\right)^4$$

$$= \frac{9 \times 8 \times 7 \times 6}{4 \times 3 \times 2 \times 1} \cdot (3)^5 \left(\frac{1}{3}\right)^4$$

$$= 126 \times 3$$

$$= 378$$

$$\text{(iii) Here, } a=3x^2, b = \frac{-a}{3x^3} \text{ and } n=10$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{10}{r} (3x^2)^{10-r} \left(\frac{-a}{3x^3}\right)^r$$

$$= \binom{10}{r} (3)^{10-r} (x^2)^{10-r} \left(\frac{-a}{3}\right)^r (x)^{-3r}$$

$$= \binom{10}{r} (3)^{10-r} (x)^{20-2r} \left(\frac{-a}{3}\right)^r (x)^{-3r}$$

$$= \binom{10}{r} (3)^{10-r} (x)^{20-2r-3r} \left(\frac{-a}{3}\right)^r$$

$$= \binom{10}{r} (3)^{10-r} \left(\frac{-a}{3}\right)^r (x)^{20-5r}$$

To get coefficient of  $x^{-15}$  we must have,

$$(x)^{20-5r} = x^{-15}$$

$$\bullet 20 - 5r = -15$$

$$\bullet 5r = 35$$

$$\bullet r = 7$$

$$\text{Therefore, coefficient of } x^{-15} = \binom{10}{7} (3)^{10-7} \left(\frac{-a}{3}\right)^7$$

$$\text{But } \binom{10}{7} = \binom{10}{3} \dots\dots\dots [\because \binom{n}{r} = \binom{n}{n-r}]$$

$$\text{Therefore, the coefficient of } x^{-15} = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} \cdot (3)^3 \left(\frac{-a}{3}\right)^7$$

$$= 120 \cdot (-a)^7 \left(\frac{1}{3}\right)^4$$

$$= (-a)^7 \frac{120}{3^4}$$

$$= (-a)^7 \frac{40}{27}$$

**(iv)** Here,  $a=a$ ,  $b=-2b$  and  $n=12$

We have formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{12}{r} (a)^{12-r} (-2b)^r$$



$$= \binom{12}{r} (-2)^r (a)^{12-r} (b)^r$$

To get coefficient of  $a^7b^5$  we must have,

$$(a)^{12-r} (b)^r = a^7b^5$$

$$\bullet r = 5$$

Therefore, coefficient of  $a^7b^5 = \binom{12}{5} (-2)^5$

$$= \frac{12 \times 11 \times 10 \times 9 \times 8}{5 \times 4 \times 3 \times 2 \times 1} \cdot (-32)$$

$$= 792 \cdot (-32)$$

$$= -25344$$

**Q. 31. Show that the term containing  $x^3$  does not exist in the expansion**

of  $\left(3x - \frac{1}{2x}\right)^8$ .

**Answer :** For  $\left(3x - \frac{1}{2x}\right)^8$ ,

$$a=3x, b = \frac{-1}{2x} \text{ and } n=8$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{8}{r} (3x)^{8-r} \left(\frac{-1}{2x}\right)^r$$

$$= \binom{8}{r} (3)^{8-r} (x)^{8-r} \left(\frac{-1}{2}\right)^r (x)^{-r}$$

$$= \binom{8}{r} (3)^{8-r} (x)^{8-r-r} \left(\frac{-1}{2}\right)^r$$

$$= \binom{8}{r} (3)^{8-r} \left(\frac{-1}{2}\right)^r (x)^{8-2r}$$

To get coefficient of  $x^3$  we must have,

$$(x)^{8-2r} = (x)^3$$

$$\bullet 8 - 2r = 3$$

$$\bullet 2r = 5$$

$$\bullet r = 2.5$$

As  $\binom{8}{r} = \binom{8}{2.5}$  is not possible

Therefore, the term containing  $x^3$  does not exist in the expansion of  $\left(3x - \frac{1}{2x}\right)^8$

**Q. 32. Show that the expansion of  $\left(2x^2 - \frac{1}{x}\right)^{20}$  does not contain any term involving  $x^9$ .**

**Answer :** For  $\left(2x^2 - \frac{1}{x}\right)^{20}$ ,

$$a=2x^2, \quad b = \frac{-1}{x} \quad \text{and } n=20$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{20}{r} (2x^2)^{20-r} \left(\frac{-1}{x}\right)^r$$

$$= \binom{20}{r} (2)^{20-r} (x^2)^{20-r} (-1)^r (x)^{-r}$$

$$= \binom{20}{r} (3)^{20-r} (x)^{40-2r} (-1)^r (x)^{-r}$$

$$= \binom{20}{r} (3)^{20-r} (x)^{40-2r-r} (-1)^r$$

$$= \binom{20}{r} (3)^{20-r} (-1)^r (x)^{40-3r}$$

To get coefficient of  $x^9$  we must have,

$$(x)^{40-3r} = (x)^9$$

$$\bullet 40 - 3r = 9$$

$$\bullet 3r = 31$$

$$\bullet r = 10.3333$$

As  $\binom{20}{r} = \binom{20}{10.3333}$  is not possible

Therefore, the term containing  $x^9$  does not exist in the expansion of  $\left(2x^2 - \frac{1}{x}\right)^{20}$

**Q. 33. Show that the expansion of  $\left(x^2 + \frac{1}{x}\right)^{12}$  does not contain any term involving  $x^{-1}$ .**

**Answer :** For  $\left(x^2 + \frac{1}{x}\right)^{12}$ ,

$$a=x^2, \quad b = \frac{1}{x} \text{ and } n=12$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{12}{r} (x^2)^{12-r} \left(\frac{1}{x}\right)^r$$

$$= \binom{12}{r} (x)^{24-2r} (x)^{-r}$$

$$= \binom{12}{r} (x)^{24-2r-r}$$

$$= \binom{12}{r} (x)^{24-3r}$$

To get coefficient of  $x^{-1}$  we must have,

$$(x)^{24-3r} = (x)^{-1}$$

$$\bullet 24 - 3r = -1$$

$$\bullet 3r = 25$$

$$\bullet r = 8.3333$$

As  $\binom{20}{r} = \binom{20}{8.3333}$  is not possible

Therefore, the term containing  $x^{-1}$  does not exist in the expansion of  $\left(x^2 + \frac{1}{x}\right)^{12}$

**Q. 34. Write the general term in the expansion of**

$$(x^2 - y)^6$$

**Answer : To Find :** General term, i.e.  $t_{r+1}$

For  $(x^2 - y)^6$

$a=x^2$ ,  $b=-y$  and  $n=6$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{6}{r} (x^2)^{6-r} (-y)^r$$

**Conclusion :** General term  $= \binom{6}{r} (x^2)^{6-r} (-y)^r$

**Q. 35. Find the 5<sup>th</sup> term from the end in the expansion of  $\left(x - \frac{1}{x}\right)^{12}$ .**

**Answer :** To Find : 5<sup>th</sup> term from the end

Formulae :

$$\bullet \quad t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\bullet \quad \binom{n}{r} = \binom{n}{n-r}$$

For  $\left(x - \frac{1}{x}\right)^{12}$ ,

$$a=x, \quad b = \frac{-1}{x} \text{ and } n=12$$

As  $n=12$ , therefore there will be total  $(12+1)=13$  terms in the expansion

Therefore,

5<sup>th</sup> term from the end =  $(13-5+1)^{\text{th}}$  i.e. 9<sup>th</sup> term from the starting.

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

For  $t_9$ ,  $r=8$

$$\therefore t_9 = t_{8+1}$$

$$= \binom{12}{8} (x)^{12-8} \left(\frac{-1}{x}\right)^8$$

$$= \binom{12}{4} (x)^4 (x)^{-8} \dots\dots\dots \left[ \because \binom{n}{r} = \binom{n}{n-r} \right]$$

$$= \frac{12 \times 11 \times 10 \times 9}{4 \times 3 \times 2 \times 1} (x)^{4-8}$$

$$= 495 (x)^{-4}$$

Therefore, a 5<sup>th</sup> term from the end  $= 495 (x)^{-4}$

Conclusion : 5<sup>th</sup> term from the end  $= 495 (x)^{-4}$

**Q. 36. Find the 4<sup>th</sup> term from the end in the expansion of  $\left(\frac{4x}{5} - \frac{5}{2x}\right)^9$ .**

**Answer :** To Find : 4<sup>th</sup> term from the end

Formulae :

$$\bullet \quad t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\bullet \quad \binom{n}{r} = \binom{n}{n-r}$$

$$\text{For } \left(\frac{4x}{5} - \frac{5}{2x}\right)^9,$$

$$a = \frac{4x}{5}, \quad b = \frac{-5}{2x} \text{ and } n=9$$

As  $n=9$ , therefore there will be total  $(9+1)=10$  terms in the expansion

Therefore,

4<sup>th</sup> term from the end  $= (10-4+1)^{\text{th}}$ , i.e. 7<sup>th</sup> term from the starting.

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

For  $t_7$ ,  $r=6$

$$\therefore t_7 = t_{6+1}$$

$$= \binom{10}{6} \left(\frac{4x}{5}\right)^{10-6} \left(\frac{-5}{2x}\right)^6$$

$$= \binom{10}{4} \left(\frac{4x}{5}\right)^4 \left(\frac{-5}{2x}\right)^6 \dots \dots \dots [\because \binom{n}{r} = \binom{n}{n-r}]$$

$$= \binom{10}{4} \frac{(4)^4}{(5)^4} (x)^4 \frac{(-5)^6}{(2)^6} (x)^{-6}$$

$$= \frac{10 \times 9 \times 8 \times 7}{4 \times 3 \times 2 \times 1} (100) (x)^{-2}$$

$$= 21000 (x)^{-2}$$

Therefore, a 4<sup>th</sup> term from the end =  $21000 (x)^{-2}$

Conclusion : 4<sup>th</sup> term from the end =  $21000 (x)^{-2}$

**Q. 37. Find the 4<sup>th</sup> term from the beginning and end in the expansion**

**of**  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$  .

**Answer : To Find :**

I. 4<sup>th</sup> term from the beginning

II. 4<sup>th</sup> term from the end

**Formulae :**

•  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

•  $\binom{n}{r} = \binom{n}{n-r}$

For  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$  ,

$a = \sqrt[3]{2}$  ,  $b = \frac{1}{\sqrt[3]{3}}$  and  $n=9$

As  $n=9$  , therefore there will be total  $(n+1)$  terms in the expansion

Therefore,

I. For the 4<sup>th</sup> term from the starting.

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

For  $t_4$ ,  $r=3$

$$\therefore t_4 = t_{3+1}$$

$$= \binom{n}{3} (\sqrt[3]{2})^{n-3} \left(\frac{1}{\sqrt[3]{3}}\right)^3$$

$$= \binom{n}{3} (2)^{\frac{n-3}{3}} \frac{1}{3}$$

$$= \binom{n}{3} \cdot \frac{(2)^{\frac{n-3}{3}}}{3}$$

$$= \frac{n!}{(n-3)! \times 3!} \cdot \frac{(2)^{\frac{n-3}{3}}}{3}$$

$$\text{Therefore, a 4}^{\text{th}} \text{ term from the starting} = \frac{n!}{(n-3)! \times 3!} \cdot \frac{(2)^{\frac{n-3}{3}}}{3}$$

Now,

II. For the 4<sup>th</sup> term from the end

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

For  $t_{(n-2)}$ ,  $r = (n-2)-1 = (n-3)$

$$\therefore t_{(n-2)} = t_{(n-3)+1}$$



$$\begin{aligned}
&= \binom{n}{n-3} (\sqrt[3]{2})^{n-(n-3)} \left(\frac{1}{\sqrt[3]{3}}\right)^{(n-3)} \\
&= \binom{n}{3} (\sqrt[3]{2})^3 (3)^{\frac{-(n-3)}{3}} \dots \dots \dots \left[ \because \binom{n}{r} = \binom{n}{n-r} \right] \\
&= \binom{n}{4} (2) (3)^{\frac{3-n}{3}} \\
&= \frac{n!}{(n-4)! \times 4!} (2) (3)^{\frac{3-n}{3}}
\end{aligned}$$

Therefore, a 4<sup>th</sup> term from the end  $= \frac{n!}{(n-4)! \times 4!} (2) (3)^{\frac{3-n}{3}}$

Conclusion :

I. 4<sup>th</sup> term from the beginning  $= \frac{n!}{(n-3)! \times 3!} \cdot \frac{(2)^{\frac{n-3}{3}}}{3}$

II. 4<sup>th</sup> term from the end  $= \frac{n!}{(n-4)! \times 4!} (2) (3)^{\frac{3-n}{3}}$

**Q. 38. Find the middle term in the expansion of :**

(i)  $(3 + x)^6$

(ii)  $\left(\frac{x}{3} + 3y\right)^8$

(iii)  $\left(\frac{x}{a} - \frac{a}{x}\right)^{10}$

(iv)  $\left(x^2 - \frac{2}{x}\right)^{10}$

**Answer : (i)** For  $(3 + x)^6$ ,

$a=3$ ,  $b=x$  and  $n=6$

As n is even,  $\left(\frac{n+2}{2}\right)^{\text{th}}$  is the middle term

Therefore, the middle term  $= \left(\frac{6+2}{2}\right)^{\text{th}} = \left(\frac{8}{2}\right)^{\text{th}} = (4)^{\text{th}}$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, for  $4^{\text{th}}$ ,  $r=3$

Therefore, the middle term is

$$t_4 = t_{3+1}$$

$$= \binom{6}{3} (3)^{6-3} (x)^3$$

$$= \frac{6 \times 5 \times 4}{3 \times 2 \times 1} \cdot (3)^3 (x)^3$$

$$= (20) \cdot (27) x^3$$

$$= 540 x^3$$

(ii) For  $\left(\frac{x}{3} + 3y\right)^8$ ,

$$a = \frac{x}{3}, b=3y \text{ and } n=8$$

As n is even,  $\left(\frac{n+2}{2}\right)^{\text{th}}$  is the middle term

Therefore, the middle term  $= \left(\frac{8+2}{2}\right)^{\text{th}} = \left(\frac{10}{2}\right)^{\text{th}} = (5)^{\text{th}}$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, for 5<sup>th</sup>, r=4

Therefore, the middle term is

$$\begin{aligned}t_5 &= t_{4+1} \\&= \binom{8}{4} \left(\frac{x}{3}\right)^{8-4} (3y)^4 \\&= \binom{8}{4} \left(\frac{x}{3}\right)^4 (3)^4 (y)^4 \\&= \binom{8}{4} \frac{(x)^4}{(3)^4} (3)^4 (y)^4 \\&= \frac{8 \times 7 \times 6 \times 5}{4 \times 3 \times 2 \times 1} \cdot (x)^4 (y)^4 \\&= (70) \cdot x^4 y^4\end{aligned}$$

(iii) For  $\left(\frac{x}{a} - \frac{a}{x}\right)^{10}$ ,

$$a = \frac{x}{a}, \quad b = \frac{-a}{x} \text{ and } n=10$$

As n is even,  $\left(\frac{n+2}{2}\right)^{\text{th}}$  is the middle term

$$\text{Therefore, the middle term} = \left(\frac{10+2}{2}\right)^{\text{th}} = \left(\frac{12}{2}\right)^{\text{th}} = (6)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, for 6<sup>th</sup>, r=5

Therefore, the middle term is

$$t_6 = t_{5+1}$$

$$\begin{aligned}
&= \binom{10}{5} \left(\frac{x}{a}\right)^{10-5} \left(\frac{-a}{x}\right)^5 \\
&= \binom{10}{5} \left(\frac{x}{a}\right)^5 (-a)^5 \left(\frac{1}{x}\right)^5 \\
&= \binom{10}{5} \frac{(x)^5}{(a)^5} (-a)^5 \left(\frac{1}{x}\right)^5 \\
&= \frac{10 \times 9 \times 8 \times 7 \times 6}{5 \times 4 \times 3 \times 2 \times 1} \cdot (-1) \\
&= -252
\end{aligned}$$

(iv) For  $\left(x^2 - \frac{2}{x}\right)^{10}$ ,

$$a=x^2, \quad b = \frac{-2}{x} \text{ and } n=10$$

As n is even,  $\left(\frac{n+2}{2}\right)^{\text{th}}$  is the middle term

$$\text{Therefore, the middle term} = \left(\frac{10+2}{2}\right)^{\text{th}} = \left(\frac{12}{2}\right)^{\text{th}} = (6)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Therefore, for the 6<sup>th</sup> middle term,  $r=5$

Therefore, the middle term is

$$\begin{aligned}
t_6 &= t_{5+1} \\
&= \binom{10}{5} (x^2)^{10-5} \left(\frac{-2}{x}\right)^5
\end{aligned}$$

$$\begin{aligned}
&= \binom{10}{5} (x^2)^5 (-2)^5 \left(\frac{1}{x}\right)^5 \\
&= \binom{10}{5} \frac{(x)^{10}}{(x)^5} (-32) \\
&= \frac{10 \times 9 \times 8 \times 7 \times 6}{5 \times 4 \times 3 \times 2 \times 1} \cdot (-32) (x)^5 \\
&= -252 (32) x^5 \\
&= -8064 x^5
\end{aligned}$$

**Q. 39. A. Find the two middle terms in the expansion of :**

$$(x^2 + a^2)^5$$

**Answer :** For  $(x^2 + a^2)^5$ ,

$$a = x^2, b = a^2 \text{ and } n = 5$$

As  $n$  is odd, there are two middle terms i.e.

$$\text{I. } \left(\frac{n+1}{2}\right)^{\text{th}} \text{ and II. } \left(\frac{n+3}{2}\right)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\text{I. The first, middle term is } \left(\frac{n+1}{2}\right)^{\text{th}} = \left(\frac{5+1}{2}\right)^{\text{th}} = \left(\frac{6}{2}\right)^{\text{th}} = (3)^{\text{rd}}$$

Therefore, for the 3<sup>rd</sup> middle term,  $r=2$

Therefore, the first middle term is

$$\begin{aligned}
t_3 &= t_{2+1} \\
&= \binom{5}{2} (x^2)^{5-2} (a^2)^2
\end{aligned}$$

$$= \binom{5}{2} (x^2)^3 (a)^4$$

$$= \binom{5}{2} (x)^6 (a)^4$$

$$= \frac{5 \times 4}{2 \times 1} \cdot (x)^6 (a)^4$$

$$= 10 \cdot a^4 \cdot x^6$$

II. The second middle term is  $\binom{\frac{n+3}{2}}{2}^{\text{th}} = \binom{\frac{5+3}{2}}{2}^{\text{th}} = \binom{\frac{8}{2}}{2}^{\text{th}} = (4)^{\text{th}}$

Therefore, for the 4<sup>th</sup> middle term, r=3

Therefore, the second middle term is

$$t_4 = t_{3+1}$$

$$= \binom{5}{3} (x^2)^{5-3} (a^2)^3$$

$$= \binom{5}{3} (x^2)^2 (a)^6$$

$$= \binom{5}{2} (x)^4 (a)^6 \dots\dots\dots [\because \binom{n}{r} = \binom{n}{n-r}]$$

$$= \frac{5 \times 4}{2 \times 1} \cdot (x)^4 (a)^6$$

$$= 10 \cdot a^6 \cdot x^4$$

Q. 39. B. Find the two middle terms in the expansion of:

$$\left(x^4 - \frac{1}{x^3}\right)^{11}$$

**Answer :** For  $\left(x^4 - \frac{1}{x^3}\right)^{11}$ ,

$$a = x^4, b = \frac{-1}{x^3} \text{ and } n=11$$

As n is odd, there are two middle terms i.e.

$$\text{II. } \left(\frac{n+1}{2}\right)^{\text{th}} \text{ and II. } \left(\frac{n+3}{2}\right)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\text{I. The first middle term is } \left(\frac{n+1}{2}\right)^{\text{th}} = \left(\frac{11+1}{2}\right)^{\text{th}} = \left(\frac{12}{2}\right)^{\text{th}} = (6)^{\text{th}}$$

Therefore, for the 6<sup>th</sup> middle term,  $r=5$

Therefore, the first middle term is

$$\begin{aligned} t_6 &= t_{5+1} \\ &= \binom{11}{5} (x^4)^{11-5} \left(\frac{-1}{x^3}\right)^5 \\ &= \binom{11}{5} (x^4)^6 (-1)^5 \left(\frac{1}{x^3}\right)^5 \\ &= \binom{11}{5} (x)^{24} (-1) \frac{1}{x^{15}} \\ &= \frac{11 \times 10 \times 9 \times 8 \times 7}{5 \times 4 \times 3 \times 2 \times 1} \cdot (x)^9 (-1) \\ &= -462 \cdot x^9 \end{aligned}$$

$$\text{II. The second middle term is } \left(\frac{n+3}{2}\right)^{\text{th}} = \left(\frac{11+3}{2}\right)^{\text{th}} = \left(\frac{14}{2}\right)^{\text{th}} = (7)^{\text{th}}$$

Therefore, for the 7<sup>th</sup> middle term,  $r=6$

Therefore, the second middle term is

$$\begin{aligned}
t_7 &= t_{6+1} \\
&= \binom{11}{6} (x^4)^{11-6} \left(\frac{-1}{x^3}\right)^6 \\
&= \binom{11}{5} (x^4)^5 (-1)^6 \left(\frac{1}{x^3}\right)^6 \dots\dots\dots [\because \binom{n}{r} = \binom{n}{n-r}] \\
&= \binom{11}{5} (x)^{20} (1) \frac{1}{x^{18}} \\
&= \frac{11 \times 10 \times 9 \times 8 \times 7}{5 \times 4 \times 3 \times 2 \times 1} \cdot (x)^2 \\
&= 462 \cdot x^2
\end{aligned}$$

**Q. 39. C. Find the two middle terms in the expansion of :**

$$\left(\frac{p}{x} + \frac{x}{p}\right)^9$$

**Answer :** For  $\left(\frac{p}{x} + \frac{x}{p}\right)^9$ ,

$$a = \frac{p}{x}, \quad b = \frac{x}{p} \text{ and } n=9$$

As n is odd, there are two middle terms i.e.

$$\text{I. } \left(\frac{n+1}{2}\right)^{\text{th}} \text{ and II. } \left(\frac{n+3}{2}\right)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\text{I. The first middle term is } \left(\frac{n+1}{2}\right)^{\text{th}} = \left(\frac{9+1}{2}\right)^{\text{th}} = \left(\frac{10}{2}\right)^{\text{th}} = (5)^{\text{th}}$$

Therefore, for 5<sup>th</sup> middle term, r=4



Therefore, the first middle term is

$$\begin{aligned}
 t_5 &= t_{4+1} \\
 &= \binom{9}{4} \left(\frac{p}{x}\right)^{9-4} \left(\frac{x}{p}\right)^4 \\
 &= \binom{9}{4} \left(\frac{p}{x}\right)^5 (x)^4 \left(\frac{1}{p}\right)^4 \\
 &= \binom{9}{4} \left(\frac{p}{x}\right) \\
 &= \frac{9 \times 8 \times 7 \times 6}{4 \times 3 \times 2 \times 1} \cdot (p) \cdot (x)^{-1} \\
 &= 126p \cdot x^{-1}
 \end{aligned}$$

II. The second middle term is  $\left(\frac{n+3}{2}\right)^{\text{th}} = \left(\frac{9+3}{2}\right)^{\text{th}} = \left(\frac{12}{2}\right)^{\text{th}} = (6)^{\text{th}}$

Therefore, for the 6<sup>th</sup> middle term, r=5

Therefore, the second middle term is

$$\begin{aligned}
 t_6 &= t_{5+1} \\
 &= \binom{9}{5} \left(\frac{p}{x}\right)^{9-5} \left(\frac{x}{p}\right)^5 \\
 &= \binom{9}{4} \left(\frac{p}{x}\right)^4 (x)^5 \left(\frac{1}{p}\right)^5 \dots\dots\dots \left[\because \binom{n}{r} = \binom{n}{n-r}\right] \\
 &= \binom{9}{4} \left(\frac{x}{p}\right) \\
 &= \frac{9 \times 8 \times 7 \times 6}{4 \times 3 \times 2 \times 1} \cdot \left(\frac{1}{p}\right) \cdot (x)
 \end{aligned}$$

$$= 126 \left( \frac{1}{p} \right) \cdot (x)$$

**Q. 39. D. Find the two middle terms in the expansion of :**

$$\left( 3x - \frac{x^3}{6} \right)^9$$

**Answer :** For  $\left( 3x - \frac{x^3}{6} \right)^9$ ,

$$a=3x, \quad b = \frac{-x^3}{6} \quad \text{and } n=9$$

As  $n$  is odd, there are two middle terms i.e.

$$\text{I. } \left( \frac{n+1}{2} \right)^{\text{th}} \quad \text{and II. } \left( \frac{n+3}{2} \right)^{\text{th}}$$

General term  $t_{r+1}$  is given by,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$\text{I. The first middle term is } \left( \frac{n+1}{2} \right)^{\text{th}} = \left( \frac{9+1}{2} \right)^{\text{th}} = \left( \frac{10}{2} \right)^{\text{th}} = (5)^{\text{th}}$$

Therefore, for 5<sup>th</sup> middle term,  $r=4$

Therefore, the first middle term is

$$t_5 = t_{4+1}$$

$$= \binom{9}{4} (3x)^{9-4} \left( \frac{-x^3}{6} \right)^4$$

$$= \binom{9}{4} (3x)^5 (x^3)^4 \left( \frac{1}{6} \right)^4$$

$$\begin{aligned}
&= \binom{9}{4} (3)^5 (x)^5 (x)^{12} \left(\frac{1}{6}\right)^4 \\
&= \frac{9 \times 8 \times 7 \times 6}{4 \times 3 \times 2 \times 1} \cdot \frac{243}{1296} (x)^{17} \\
&= \frac{189}{8} (x)^{17}
\end{aligned}$$

II. The second middle term is  $\binom{\frac{n+3}{2}}{2}^{\text{th}} = \binom{\frac{9+3}{2}}{2}^{\text{th}} = \binom{\frac{12}{2}}{2}^{\text{th}} = (6)^{\text{th}}$

Therefore, for the 6<sup>th</sup> middle term,  $r=5$

Therefore, the second middle term is

$$\begin{aligned}
t_6 &= t_{5+1} \\
&= \binom{9}{5} (3x)^{9-5} \left(\frac{-x^3}{6}\right)^5 \\
&= \binom{9}{4} (3x)^4 (-x^3)^5 \left(\frac{1}{6}\right)^5 \dots\dots\dots [\because \binom{n}{r} = \binom{n}{n-r}] \\
&= \binom{9}{4} (3)^4 (x)^4 (-x)^{15} \left(\frac{1}{6}\right)^5 \\
&= \frac{9 \times 8 \times 7 \times 6}{4 \times 3 \times 2 \times 1} \cdot \frac{81}{7776} (-x)^{19} \\
&= -\frac{21}{16} (x)^{19}
\end{aligned}$$

**Q. 40. A. Find the term independent of x in the expansion of :**

$$\left(2x + \frac{1}{3x^2}\right)^9$$

**Answer : To Find :** term independent of x, i.e.  $x^0$

For  $\left(2x + \frac{1}{3x^2}\right)^9$

$a=2x$ ,  $b = \frac{1}{3x^2}$  and  $n=9$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{9}{r} (2x)^{9-r} \left(\frac{1}{3x^2}\right)^r$$

$$= \binom{9}{r} (x)^{9-r} (2)^{9-r} \left(\frac{1}{3}\right)^r \left(\frac{1}{x^2}\right)^r$$

$$= \binom{9}{r} (x)^{9-r} \frac{(2)^{9-r}}{(3)^r} (x)^{-2r}$$

$$= \binom{9}{r} \frac{(2)^{9-r}}{(3)^r} (x)^{9-r-2r}$$

$$= \binom{9}{r} \frac{(2)^{9-r}}{(3)^r} (x)^{9-3r}$$

Now, to get coefficient of term independent of  $x$  that is coefficient of  $x^0$  we must have,

$$(x)^{9-3r} = x^0$$

$$\bullet 9 - 3r = 0$$

$$\bullet 3r = 9$$

$$\bullet r = 3$$

Therefore, coefficient of  $x^0 = \binom{9}{3} \frac{(2)^{9-3}}{(3)^3}$

$$= \frac{9 \times 8 \times 7 (2)^6}{3 \times 2 \times 1 (3)^3}$$

$$= \frac{1792}{3}$$

Conclusion : coefficient of  $x^0 = \frac{1792}{3}$

**Q. 40. B. Find the term independent of x in the expansion of :**

$$\left( \frac{3x^2}{2} - \frac{1}{3x} \right)^6$$

**Answer :** To Find : term independent of x, i.e.  $x^0$

$$\text{For } \left( \frac{3x^2}{2} - \frac{1}{3x} \right)^6$$

$$a = \frac{3x^2}{2}, \quad b = -\frac{1}{3x} \text{ and } n=6$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{6}{r} \left( \frac{3x^2}{2} \right)^{6-r} \left( -\frac{1}{3x} \right)^r$$

$$= \binom{6}{r} \left( \frac{3}{2} \right)^{6-r} (x^2)^{6-r} \left( \frac{-1}{3} \right)^r \left( \frac{1}{x} \right)^r$$

$$= \binom{6}{r} \left( \frac{3}{2} \right)^{6-r} \left( \frac{-1}{3} \right)^r (x)^{12-2r} (x)^{-r}$$

$$= \binom{6}{r} \left( \frac{3}{2} \right)^{6-r} \left( \frac{-1}{3} \right)^r (x)^{12-2r-r}$$

$$= \binom{6}{r} \left(\frac{3}{2}\right)^{6-r} \left(\frac{-1}{3}\right)^r (x)^{12-3r}$$

Now, to get coefficient of term independent of x that is coefficient of  $x^0$  we must have,

$$(x)^{12-3r} = x^0$$

$$\bullet 12 - 3r = 0$$

$$\bullet 3r = 12$$

$$\bullet r = 4$$

$$\text{Therefore, coefficient of } x^0 = \binom{6}{4} \left(\frac{3}{2}\right)^{6-4} \left(\frac{-1}{3}\right)^4$$

$$= \binom{6}{2} \left(\frac{3}{2}\right)^2 \frac{1}{81} \dots\dots\dots [\because \binom{n}{r} = \binom{n}{n-r}]$$

$$= \frac{6 \times 5}{2 \times 1} \cdot \frac{9}{4} \cdot \frac{1}{81}$$

$$= \frac{15}{36}$$

$$\text{Conclusion : coefficient of } x^0 = \frac{15}{36}$$

**Q. 40. C. Find the term independent of x in the expansion of :**

$$\left(x - \frac{1}{x^2}\right)^{3n}$$

**Answer : To Find :** term independent of x, i.e.  $x^0$

$$\text{For } \left(x - \frac{1}{x^2}\right)^{3n}$$

$$a=x, b = -\frac{1}{x^2} \text{ and } N=3n$$

We have a formula,

$$\begin{aligned}
t_{r+1} &= \binom{N}{r} a^{N-r} b^r \\
&= \binom{3n}{r} (x)^{3n-r} \left(-\frac{1}{x^2}\right)^r \\
&= \binom{3n}{r} (x)^{3n-r} (-1)^r \left(\frac{1}{x^2}\right)^r \\
&= \binom{3n}{r} (x)^{3n-r} (-1)^r (x)^{-2r} \\
&= \binom{3n}{r} (-1)^r (x)^{3n-r-2r} \\
&= \binom{3n}{r} (-1)^r (x)^{3n-3r}
\end{aligned}$$

Now, to get coefficient of term independent of  $x$  that is coefficient of  $x^0$  we must have,

$$(x)^{3n-3r} = x^0$$

$$\bullet 3n - 3r = 0$$

$$\bullet 3r = 3n$$

$$\bullet r = n$$

$$\text{Therefore, coefficient of } x^0 = \binom{3n}{n} (-1)^n$$

$$\text{Conclusion : coefficient of } x^0 = \binom{3n}{n} (-1)^n$$

**Q. 40. D. Find the term independent of  $x$  in the expansion of :**

$$\left(3x - \frac{2}{x^2}\right)^{15}$$

**Answer : To Find :** term independent of  $x$ , i.e.  $x^0$

$$\text{For } \left(3x - \frac{2}{x^2}\right)^{15}$$

$$a=3x, \quad b = \frac{-2}{x^2} \text{ and } n=15$$

We have a formula,

$$\begin{aligned} t_{r+1} &= \binom{n}{r} a^{n-r} b^r \\ &= \binom{15}{r} (3x)^{15-r} \left(\frac{-2}{x^2}\right)^r \\ &= \binom{15}{r} (3)^{15-r} (x)^{15-r} (-2)^r \left(\frac{1}{x^2}\right)^r \\ &= \binom{15}{r} (3)^{15-r} (x)^{15-r} (-2)^r (x)^{-2r} \\ &= \binom{15}{r} (3)^{15-r} (-2)^r (x)^{15-r-2r} \\ &= \binom{15}{r} (3)^{15-r} (-2)^r (x)^{15-3r} \end{aligned}$$

Now, to get coefficient of term independent of  $x$  that is coefficient of  $x^0$  we must have,

$$(x)^{15-3r} = x^0$$

$$\bullet 15 - 3r = 0$$

$$\bullet 3r = 15$$

$$\bullet r = 5$$

$$\text{Therefore, coefficient of } x^0 = \binom{15}{5} (3)^{15-5} (-2)^5$$

$$= \frac{15 \times 14 \times 13 \times 12 \times 11}{5 \times 4 \times 3 \times 2 \times 1} \cdot (3)^{10} \cdot (-32)$$

$$= -3003 \cdot (3)^{10} \cdot (32)$$



Conclusion : coefficient of  $x^0 = -3003 \cdot (3)^{10} \cdot (32)$

**Q. 41. Find the coefficient of  $x^5$  in the expansion of  $(1 + x)^3 (1 - x)^6$ .**

**Answer :** To Find : coefficient of  $x^5$

For  $(1+x)^3$

$a=1$ ,  $b=x$  and  $n=3$

We have a formula,

$$\begin{aligned}(1+x)^3 &= \sum_{r=0}^3 \binom{3}{r} (1)^{3-r} x^r \\&= \binom{3}{0} (1)^3 x^0 + \binom{3}{1} (1)^2 x^1 + \binom{3}{2} (1)^1 x^2 + \binom{3}{3} (1)^0 x^3 \\&= 1 + 3x + 3x^2 + x^3\end{aligned}$$

For  $(1-x)^6$

$a=1$ ,  $b=-x$  and  $n=6$

We have formula,

$$\begin{aligned}(1-x)^6 &= \sum_{r=0}^6 \binom{6}{r} (1)^{6-r} (-x)^r \\&= \binom{6}{0} (1)^6 (-x)^0 + \binom{6}{1} (1)^5 (-x)^1 + \binom{6}{2} (1)^4 (-x)^2 + \binom{6}{3} (1)^3 (-x)^3 \\&\quad + \binom{6}{4} (1)^2 (-x)^4 + \binom{6}{5} (1)^1 (-x)^5 + \binom{6}{6} (1)^0 (-x)^6\end{aligned}$$

We have a formula,

$$\binom{n}{r} = \frac{n!}{(n-r)! \times r!}$$

By using this formula, we get,  $\times$

$$(1-x)^6 = 1 - 6x + 15x^2 - 20x^3 + 15x^4 - 6x^5 + x^6$$

$$\therefore (1+x)^3(1-x)^6$$

$$= (1+3x+3x^2+x^3)(1-6x+15x^2-20x^3+15x^4-6x^5+x^6)$$

Coefficients of  $x^5$  are

$$x^0 \cdot x^5 = 1 \times (-6) = -6$$

$$x^1 \cdot x^4 = 3 \times 15 = 45$$

$$x^2 \cdot x^3 = 3 \times (-20) = -60$$

$$x^3 \cdot x^2 = 1 \times 15 = 15$$

Therefore, Coefficients of  $x^5 = -6 + 45 - 60 + 15 = -6$

Conclusion : Coefficients of  $x^5 = -6$

**Q. 42. Find numerically the greatest term in the expansion of  $(2 + 3x)^9$ ,**

**where**  $x = \frac{3}{2}$ .

**Answer : To Find** : numerically greatest term

For  $(2+3x)^9$ ,

$a=2$ ,  $b=3x$  and  $n=9$

We have relation,

$$t_{r+1} \geq t_r \text{ or } \frac{t_{r+1}}{t_r} \geq 1$$

We have a formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$= \binom{9}{r} 2^{9-r} (3x)^r$$

$$= \frac{9!}{(9-r)! \times r!} 2^{9-r} (3)^r (x)^r$$

$$\therefore t_r = \binom{n}{r-1} a^{n-r+1} b^{r-1}$$

$$= \binom{9}{r-1} 2^{9-r+1} (3x)^{r-1}$$

$$= \frac{9!}{(9-r+1)! \times (r-1)!} 2^{10-r} (3)^{r-1} (x)^{r-1}$$

$$= \frac{9!}{(10-r)! \times (r-1)!} 2^{10-r} (3)^{r-1} (x)^{r-1}$$

$$\therefore \frac{t_{r+1}}{t_r} \geq 1$$

$$\therefore \frac{\frac{9!}{(9-r)! \times r!} 2^{9-r} (3)^r (x)^r}{\frac{9!}{(10-r)! \times (r-1)!} 2^{10-r} (3)^{r-1} (x)^{r-1}} \geq 1$$

$$\therefore \frac{9!}{(9-r)! \times r!} 2^{9-r} (3)^r (x)^r \geq \frac{9!}{(10-r)! \times (r-1)!} 2^{10-r} (3)^{r-1} (x)^{r-1}$$

$$\begin{aligned} \therefore \frac{9!}{(9-r)! \times r(r-1)!} 2^{9-r} (3)(3)^{r-1} (x)(x)^{r-1} \\ \geq \frac{9!}{(10-r)(9-r)! \times (r-1)!} (2)2^{9-r} (3)^{r-1} (x)^{r-1} \end{aligned}$$

$$\therefore \frac{1}{r} (3)(x) \geq \frac{1}{(10-r)} (2)$$

At  $x = 3/2$

$$\therefore \frac{1}{r} (3) \frac{3}{2} \geq \frac{1}{(10-r)} (2)$$

$$\therefore \frac{9}{4} \geq \frac{r}{(10-r)}$$

$$\therefore 9(10-r) \geq 4r$$

$$\therefore 90 - 9r \geq 4r$$

$$\bullet 90 \geq 13r$$

$$\bullet r \leq 6.923$$

Therefore,  $r=6$  and hence the 7<sup>th</sup> term is numerically greater.

By using formula,

$$t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

$$t_7 = \binom{9}{7} 2^{9-7} (3x)^7$$

$$= \binom{9}{2} 2^2 (3)^7 (x)^7$$

Conclusion : the 7<sup>th</sup> term is numerically greater with value  $\binom{9}{2} 2^2 (3)^7 (x)^7$

**Q. 43.** If the coefficients of 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> terms in the expansion of  $(1+x)^{2n}$  are in AP, show that  $2n^2 - 9n + 7 = 0$ .

**Answer** : For  $(1+x)^{2n}$

$a=1$ ,  $b=x$  and  $N=2n$

We have,  $t_{r+1} = \binom{N}{r} a^{N-r} b^r$

For the 2<sup>nd</sup> term,  $r=1$

$$\therefore t_2 = t_{1+1}$$

$$= \binom{2n}{1} (1)^{2n-1} (x)^1$$

$$= (2n) \times \dots \left[ \because \binom{n}{1} = n \right]$$

Therefore, the coefficient of 2<sup>nd</sup> term = (2n)

For the 3<sup>rd</sup> term, r=2

$$\therefore t_3 = t_{2+1}$$

$$= \binom{2n}{2} (1)^{2n-2} (x)^2$$

$$= \frac{(2n)!}{(2n-2)! \times 2!} x^2$$

$$= \frac{(2n)(2n-1)(2n-2)!}{(2n-2)! \times 2} x^2 \dots \dots \dots (n! = n \cdot (n-1)!)$$

$$= (n)(2n-1) x^2$$

Therefore, the coefficient of 3<sup>rd</sup> term = (n)(2n-1)

For the 4<sup>th</sup> term, r=3

$$\therefore t_4 = t_{3+1}$$

$$= \binom{2n}{3} (1)^{2n-3} (x)^3$$

$$= \frac{(2n)!}{(2n-3)! \times 3!} x^3$$

$$= \frac{(2n)(2n-1)(2n-2)(2n-3)!}{(2n-3)! \times 6} x^3 \dots \dots \dots (n! = n \cdot (n-1)!)$$

$$= \frac{(n)(2n-1) \cdot 2(n-1)}{3} x^3$$

$$= \frac{2(n)(2n-1) \cdot (n-1)}{3} x^3$$

Therefore, the coefficient of 3<sup>rd</sup> term  $= \frac{2(n)(2n-1).(n-1)}{3}$

As the coefficients of 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> terms are in A.P.

Therefore,

2 × coefficient of 3<sup>rd</sup> term = coefficient of 2<sup>nd</sup> term + coefficient of the 4<sup>th</sup> term

$$\therefore 2 \times (n)(2n-1) = (2n) + \frac{2(n)(2n-1).(n-1)}{3}$$

Dividing throughout by (2n),

$$\therefore 2n - 1 = 1 + \frac{(2n-1)(n-1)}{3}$$

$$\therefore 2n - 1 = \frac{3 + (2n-1)(n-1)}{3}$$

$$\bullet 3(2n-1) = 3 + (2n-1)(n-1)$$

$$\bullet 6n - 3 = 3 + (2n^2 - 2n - n + 1)$$

$$\bullet 6n - 3 = 3 + 2n^2 - 3n + 1$$

$$\bullet 3 + 2n^2 - 3n + 1 - 6n + 3 = 0$$

$$\bullet 2n^2 - 9n + 7 = 0$$

Conclusion : If the coefficients of 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> terms of  $(1+x)^{2n}$  are in A.P. then  $2n^2 - 9n + 7 = 0$

**Q. 44. Find the 6<sup>th</sup> term of the expansion  $(y^{1/2} + x^{1/3})^n$ , if the binomial coefficient of the 3<sup>rd</sup> term from the end is 45.**

**Answer :** Given : 3<sup>rd</sup> term from the end = 45

To Find : 6<sup>th</sup> term

For  $(y^{1/2} + x^{1/3})^n$ ,

$$a = y^{1/2}, b = x^{1/3}$$

We have,  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

As  $n=n$ , therefore there will be total  $(n+1)$  terms in the expansion.

3<sup>rd</sup> term from the end =  $(n+1-3+1)^{\text{th}}$  i.e.  $(n-1)^{\text{th}}$  term from the starting

For  $(n-1)^{\text{th}}$  term,  $r = (n-1-1) = (n-2)$

$$t_{(n-1)} = t_{(n-2)+1}$$

$$= \binom{n}{n-2} \left(y^{\frac{1}{2}}\right)^{n-(n-2)} \left(x^{\frac{1}{3}}\right)^{(n-2)}$$

$$= \binom{n}{2} \left(y^{\frac{1}{2}}\right)^2 (x)^{\frac{n-2}{3}} \dots \because \binom{n}{n-r} = \binom{n}{r}$$

$$= \frac{n(n-1)}{2} (y) (x)^{\frac{n-2}{3}}$$

$$\text{Therefore 3}^{\text{rd}} \text{ term from the end} = \frac{n(n-1)}{2} (y) (x)^{\frac{n-2}{3}}$$

$$\text{Therefore coefficient 3}^{\text{rd}} \text{ term from the end} = \frac{n(n-1)}{2}$$

$$\therefore 45 = \frac{n(n-1)}{2}$$

$$\bullet 90 = n(n-1)$$

$$\bullet 10(9) = n(n-1)$$

Comparing both sides,  $n=10$

For 6<sup>th</sup> term,  $r=5$

$$t_6 = t_{5+1}$$

$$= \binom{10}{5} \left(y^{\frac{1}{2}}\right)^{10-5} \left(x^{\frac{1}{3}}\right)^5$$

$$\begin{aligned}
&= \binom{10}{5} (y)^{\frac{5}{2}} (x)^{\frac{5}{3}} \\
&= \frac{10 \times 9 \times 8 \times 7 \times 6}{5 \times 4 \times 3 \times 2 \times 1} (y)^{\frac{5}{2}} (x)^{\frac{5}{3}} \\
&= 252 (y)^{\frac{5}{2}} (x)^{\frac{5}{3}}
\end{aligned}$$

Conclusion : 6<sup>th</sup> term =  $252 (y)^{\frac{5}{2}} (x)^{\frac{5}{3}}$

**Q. 45.** If the 17<sup>th</sup> and 18<sup>th</sup> terms in the expansion of  $(2 + a)^{50}$  are equal, find the value of a.

**Answer :** Given :  $t_{17} = t_{18}$

To Find : value of a

For  $(2 + a)^{50}$

A=2, b=a and n=50

We have,  $t_{r+1} = \binom{n}{r} A^{n-r} b^r$

For the 17<sup>th</sup> term, r=16

$$\begin{aligned}
\therefore t_{17} &= t_{16+1} \\
&= \binom{50}{16} (2)^{50-16} (a)^{16} \\
&= \binom{50}{16} (2)^{34} (a)^{16}
\end{aligned}$$

For the 18<sup>th</sup> term, r=17

$$\begin{aligned}
\therefore t_{18} &= t_{17+1} \\
&= \binom{50}{17} (2)^{50-17} (a)^{17}
\end{aligned}$$



$$= \binom{50}{17} (2)^{33} (a)^{17}$$

As 17<sup>th</sup> and 18<sup>th</sup> terms are equal

$$\therefore t_{18} = t_{17}$$

$$\therefore \binom{50}{17} (2)^{33} (a)^{17} = \binom{50}{16} (2)^{34} (a)^{16}$$

$$\therefore \binom{50}{17} (2)^{33} (a)^{17} = \binom{50}{16} (2)^{34} (a)^{16}$$

$$\therefore \frac{50!}{(50-17)! \times (17)!} (2)^{33} (a)^{17} = \frac{50!}{(50-16)! \times (16)!} (2)^{34} (a)^{16}$$

$$\dots\dots \left[ \because \binom{n}{r} = \frac{n!}{(n-r)! \times (r)!} \right]$$

$$\therefore \frac{(a)^{17}}{(a)^{16}} = \frac{50!}{(50-16)! \times (16)!} \cdot \frac{(50-17)! \times (17)!}{50!} \cdot \frac{(2)^{34}}{(2)^{33}}$$

$$\therefore a = \frac{(50-17) \times (50-16)! \times 17 \times (16)!}{(50-16)! \times (16)!} \cdot (2)$$

$$\dots\dots [\because n! = n(n-1)!]$$

$$\therefore a = (50-17) \times 17 \cdot (2)$$

$$\bullet a = 1122$$

Conclusion : value of a = 1122

**Q. 46. Find the coefficient of  $x^4$  in the expansion of  $(1+x)^n (1-x)^n$ . Deduce that  $C_2 = C_0C_4 - C_1C_3 + C_2C_2 - C_3C_1 + C_4C_0$ , where  $C_r$  stands for  ${}^nC_r$ .**

**Answer :** To Find : Coefficients of  $x^4$

For  $(1+x)^n$

a=1, b=x

We have a formula,

$$\begin{aligned}
 (1+x)^n &= \sum_{r=0}^n \binom{n}{r} (1)^{n-r} x^r \\
 &= \binom{n}{0} (1)^n x^0 + \binom{n}{1} (1)^{n-1} x^1 + \binom{n}{2} (1)^{n-2} x^2 + \dots + \binom{n}{n} (1)^{n-n} x^n \\
 &= \binom{n}{0} x^0 + \binom{n}{1} x + \binom{n}{2} x^2 + \dots + \binom{n}{n} x^n
 \end{aligned}$$

For  $(1-x)^n$

$a=1$ ,  $b=-x$  and  $n=n$

We have formula,

$$\begin{aligned}
 (1-x)^n &= \sum_{r=0}^n \binom{n}{r} (1)^{n-r} (-x)^r \\
 &= \binom{n}{0} (1)^n (-x)^0 + \binom{n}{1} (1)^{n-1} (-x)^1 + \binom{n}{2} (1)^{n-2} (-x)^2 + \dots \\
 &\quad + \binom{n}{n} (1)^{n-n} (-x)^n \\
 &= \binom{n}{0} (-x)^0 - \binom{n}{1} (x)^1 + \binom{n}{2} (x)^2 + \dots + \binom{n}{n} (-x)^n \\
 \therefore (1+x)^3(1-x)^6 \\
 &= \left\{ \binom{n}{0} x^0 + \binom{n}{1} x + \binom{n}{2} x^2 + \dots + \binom{n}{n} x^n \right\} \left\{ \binom{n}{0} (-x)^0 - \binom{n}{1} (x)^1 + \binom{n}{2} (x)^2 \right. \\
 &\quad \left. + \dots + \binom{n}{n} (-x)^n \right\}
 \end{aligned}$$

Coefficients of  $x^4$  are

$$x^0 \cdot x^4 = \binom{n}{0} \times \binom{n}{4} = C_0 C_4$$

$$x^1 \cdot x^3 = \binom{n}{1} \times (-1) \binom{n}{3} = -\binom{n}{1} \binom{n}{3} = -C_1 C_3$$

$$x^2 \cdot x^2 = \binom{n}{2} \times \binom{n}{2} = C_2 C_2$$

$$x^3 \cdot x^1 = \binom{n}{3} \times (-1) \binom{n}{1} = -\binom{n}{3} \binom{n}{1} = -C_3 C_1$$

$$x^4 \cdot x^0 = \binom{n}{4} \times \binom{n}{0} = C_4 C_0$$

Therefore, Coefficient of  $x^4$

$$= C_4 C_0 - C_1 C_3 + C_2 C_2 - C_3 C_1 + C_4 C_0$$

Let us assume,  $n=4$ , it becomes

$${}^4C_4 {}^4C_0 - {}^4C_1 {}^4C_3 + {}^4C_2 {}^4C_2 - {}^4C_3 {}^4C_1 + {}^4C_4 {}^4C_0$$

We know that,

$$\binom{n}{r} = \frac{n!}{(n-r)! \times r!}$$

By using above formula, we get,

$${}^4C_4 {}^4C_0 - {}^4C_1 {}^4C_3 + {}^4C_2 {}^4C_2 - {}^4C_3 {}^4C_1 + {}^4C_4 {}^4C_0$$

$$= (1)(1) - (4)(4) + (6)(6) - (4)(4) + (1)(1)$$

$$= 1 - 16 + 36 - 16 + 1$$

$$= 6$$

$$= {}^4C_2$$

Therefore, in general,

$$C_4 C_0 - C_1 C_3 + C_2 C_2 - C_3 C_1 + C_4 C_0 = C_2$$

Therefore, Coefficient of  $x^4 = C_2$

Conclusion :

- Coefficient of  $x^4 = C_2$
- $C_4 C_0 - C_1 C_3 + C_2 C_2 - C_3 C_1 + C_4 C_0 = C_2$

**Q. 47. Prove that the coefficient of  $x^n$  in the binomial expansion of  $(1 + x)^{2n}$  is twice the coefficient of  $x^n$  in the binomial expansion of  $(1 + x)^{2n-1}$ .**

**Answer :** To Prove : coefficient of  $x^n$  in  $(1+x)^{2n} = 2 \times$  coefficient of  $x^n$  in  $(1+x)^{2n-1}$

For  $(1+x)^{2n}$ ,

$a=1$ ,  $b=x$  and  $m=2n$

We have a formula,

$$t_{r+1} = \binom{m}{r} a^{m-r} b^r$$

$$= \binom{2n}{r} (1)^{2n-r} (x)^r$$

$$= \binom{2n}{r} (x)^r$$

To get the coefficient of  $x^n$ , we must have,

$$x^n = x^r$$

$$\bullet r = n$$

Therefore, the coefficient of  $x^n = \binom{2n}{n}$

$$= \frac{(2n)!}{n! \times (2n-n)!} \dots\dots\dots \left( \because \binom{n}{r} = \frac{n!}{r! \times (n-r)!} \right)$$

$$= \frac{(2n)!}{n! \times n!}$$

$$= \frac{2n \times (2n-1)!}{n! \times n(n-1)!} \dots\dots\dots (\because n! = n(n-1)!)$$

$$= \frac{2 \times (2n-1)!}{n! \times (n-1)!} \dots\dots\dots \text{cancelling } n$$

$$\text{Therefore, the coefficient of } x^n \text{ in } (1+x)^{2n} = \frac{2 \times (2n-1)!}{n! \times (n-1)!} \dots\dots\dots \text{eq(1)}$$

Now for  $(1+x)^{2n-1}$ ,

$a=1$ ,  $b=x$  and  $m=2n-1$

We have formula,

$$\begin{aligned}t_{r+1} &= \binom{m}{r} a^{m-r} b^r \\&= \binom{2n-1}{r} (1)^{2n-1-r} (x)^r \\&= \binom{2n-1}{r} (x)^r\end{aligned}$$

To get the coefficient of  $x^n$ , we must have,

$$x^n = x^r$$

$$\bullet r = n$$

Therefore, the coefficient of  $x^n$  in  $(1+x)^{2n-1} = \binom{2n-1}{n}$

$$= \frac{(2n-1)!}{n! \times (2n-1-n)!}$$

$$= \frac{1}{2} \times \frac{2 \times (2n-1)!}{n! \times (n-1)!}$$

.....multiplying and dividing by 2

Therefore,

Coefficient of  $x^n$  in  $(1+x)^{2n-1} = \frac{1}{2} \times$  coefficient of  $x^n$  in  $(1+x)^{2n}$

Or coefficient of  $x^n$  in  $(1+x)^{2n} = 2 \times$  coefficient of  $x^n$  in  $(1+x)^{2n-1}$

Hence proved.

**Q. 48. Find the middle term in the expansion of  $\left(\frac{p}{2} + 2\right)^8$**

**Answer : Given :**  $a = \frac{p}{2}$ ,  $b=2$  and  $n=8$

To find : middle term

Formula :

• The middle term  $= \binom{n+2}{2}$

$$\therefore t_{r+1} = \binom{n}{r} a^{n-r} b^r$$

Here,  $n$  is even.

Hence,

$$\binom{n+2}{2} = \binom{8+2}{2} = 5$$

Therefore,  $5^{\text{th}}$  the term is the middle term.

For  $t_5$ ,  $r=4$

We have,  $t_{r+1} = \binom{n}{r} a^{n-r} b^r$

$$\therefore t_5 = \binom{8}{4} \left(\frac{p}{2}\right)^{8-4} 2^4$$

$$\therefore t_5 = \frac{8 \times 7 \times 6 \times 5}{4 \times 3 \times 2 \times 1} \cdot \left(\frac{p}{2}\right)^4 \cdot (16)$$

$$\therefore t_5 = 70 \cdot \left(\frac{p^4}{16}\right) \cdot (16)$$

$$\therefore t_5 = 70 p^4$$

Conclusion : The middle term is  $70 p^4$ .

### Exercise 10B

**Q. 1. Show that the term independent of x in the expansion of  $\left(x - \frac{1}{x}\right)^{10}$  is -252.**

**Answer :** To show: the term independent of x in the expansion of  $\left(x - \frac{1}{x}\right)^{10}$  is -252.

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$T_{r+1} = {}^nC_r x^{n-r} y^r$  where

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $\left(x - \frac{1}{x}\right)^{10}$ , we get

$$T_{r+1} = {}^{10}C_r x^{10-r} \times \left(\frac{-1}{x}\right)^r$$

For finding the term which is independent of x,

$$10 - 2r = 5$$

$$r = 5$$

Thus, the term which would be independent of x is  $T_6$

$$T_6 = {}^{10}C_5 \times x^{10-5} \times \left(\frac{-1}{x}\right)^5$$

$$T_6 = {}^{10}C_5 \times x^{10-5} \times \left(\frac{-1}{x}\right)^5$$

$$T_6 = - {}^{10}C_5$$

$$T_6 = - \frac{10!}{5!(10-5)!}$$

$$T_6 = - \frac{10!}{5! \times 5!}$$

$$T_6 = - \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2}$$

$$T_6 = - \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2}$$

$$T_6 = -252$$

Thus, the term independent of  $x$  in the expansion of  $\left(x - \frac{1}{x}\right)^{10}$  is -252.

**Q. 2. If the coefficients of  $x^2$  and  $x^3$  in the expansion of  $(3 + px)^9$  are the same then prove that  $P = \frac{9}{7}$ .**

**Answer :** To prove: that. If the coefficients of  $x^2$  and  $x^3$  in the expansion of  $(3 + px)^9$  are the same then  $P = \frac{9}{7}$ .

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,



$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $(3 + px)^9$ , we get

$$T_{r+1} = {}^9C_r \times 3^{9-r} \times (px)^r$$

For finding the term which has  $x^2$  in it, is given by

$$r=2$$

Thus, the coefficients of  $x^2$  are given by,

$$T_3 = {}^9C_2 \times 3^{9-2} \times (px)^2$$

$$T_3 = {}^9C_2 \times 3^7 \times p^2 \times x^2$$

For finding the term which has  $x^2$  in it, is given by

$$r=3$$

Thus, the coefficients of  $x^3$  are given by,

$$T_3 = {}^9C_3 \times 3^{9-3} \times (px)^3$$

$$T_3 = {}^9C_3 \times 3^6 \times p^3 \times x^3$$

As the coefficients of  $x^2$  and  $x^3$  in the expansion of  $(3 + px)^9$  are the same.

$${}^9C_3 \times 3^6 \times p^3 = {}^9C_2 \times 3^7 \times p^2$$

$${}^9C_3 \times p = {}^9C_2 \times 3$$

$$\frac{9!}{3! \times 6!} \times p = \frac{9!}{2! \times 7!} \times 3$$

$$\frac{9!}{3 \times 2! \times 6!} \times p = \frac{9!}{2! \times 7 \times 6!} \times 3$$

$$p = \frac{9}{7}$$

Thus, the value of p for which coefficients of  $x^2$  and  $x^3$  in the expansion of  $(3 + px)^9$  are the same is  $\frac{9}{7}$

**Q. 3. Show that the coefficient of  $x^{-3}$  in the expansion of  $\left(x - \frac{1}{x}\right)^{11}$  is -330.**

**Answer :** To show: that the coefficient of  $x^{-3}$  in the expansion of  $\left(x - \frac{1}{x}\right)^{11}$  is -330.

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $\left(x - \frac{1}{x}\right)^{11}$ , we get

$$T_{r+1} = {}^{11}C_r x^{11-r} \times \left(\frac{-1}{x}\right)^r$$

For finding the term which has  $x^{-3}$  in it, is given by

$$11 - 2r = 3$$

$$2r = 14$$

$$r = 7$$

Thus, the term which has  $x^{-3}$  in it is  $T_8$

$$T_8 = {}^{11}C_7 \times x^{11-7} \times \left(\frac{-1}{x}\right)^7$$

$$T_8 = -{}^{11}C_7 \times x^{-3}$$

$$T_8 = -\frac{11!}{7!(11-7)!}$$

$$T_8 = -\frac{11 \times 10 \times 9 \times 8 \times 7!}{7! \times 4 \times 3 \times 2}$$

$$T_8 = -330$$

Thus, the coefficient of  $x^{-3}$  in the expansion of  $\left(x - \frac{1}{x}\right)^{11}$  is -330.

**Q. 4. Show that the middle term in the expansion of  $\left(\frac{2x^2}{3} + \frac{3}{2x^2}\right)^{10}$  is 252.**

**Answer :** To show: that the middle term in the expansion of  $\left(\frac{2x^2}{3} + \frac{3}{2x^2}\right)^{10}$  is 252.

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Total number of terms in the expansion is 11

Thus, the middle term of the expansion is  $T_6$  and is given by,

$$T_6 = {}^{10}C_5 \times \left(\frac{2x^2}{3}\right)^5 \times \left(\frac{3}{2x^2}\right)^5$$

$$T_6 = {}^{10}C_5$$

$$T_6 = \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2}$$

$$T_6 = 252$$

Thus, the middle term in the expansion of  $\left(\frac{2x^2}{3} + \frac{3}{2x^2}\right)^{10}$  is 252.

**Q. 5. Show that the coefficient of  $x^4$  in the expansion of  $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$  is  $\frac{405}{256}$ .**

**Answer :** To show: that the coefficient of  $x^4$  in the expansion of  $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$  is -330.

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$ , we get

$$T_{r+1} = {}^{10}C_r \times \left(\frac{x}{2}\right)^{10-r} \times \left(\frac{-3}{x^2}\right)^r$$

For finding the term which has  $x^4$  in it, is given by

$$10 - 3r = 4$$

$$3r = 6$$

$$R = 2$$

Thus, the term which has  $x^4$  in it is  $T_3$

$$T_3 = {}^{10}C_2 \times \left(\frac{x}{2}\right)^8 \times \left(\frac{-3}{x^2}\right)^2$$

$$T_3 = \frac{10! \times 9}{2! \times 8! \times 2^8}$$

$$T_3 = \frac{10 \times 9 \times 8! \times 9}{2 \times 8! \times 2^8}$$

$$T_3 = \frac{405}{256}$$

Thus, the coefficient of  $x^4$  in the expansion of  $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$  is  $\frac{405}{256}$

**Q. 6. Prove that there is no term involving  $x^6$  in the expansion of  $\left(2x^2 - \frac{3}{x}\right)^{11}$ .**

**Answer :** To prove: that there is no term involving  $x^6$  in the expansion of  $\left(2x^2 - \frac{3}{x}\right)^{11}$

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $\left(2x^2 - \frac{3}{x}\right)^{11}$ , we get

$$T_{r+1} = {}^{11}C_r \times (2x^2)^{11-r} \times \left(\frac{-3}{x}\right)^r$$

For finding the term which has  $x^6$  in it, is given by

$$22 - 2r - r = 6$$

$$3r = 16$$

$$r = \frac{16}{3}$$

Since,  $r = \frac{16}{3}$  is not possible as r needs to be a whole number

Thus, there is no term involving  $x^6$  in the expansion of  $\left(2x^2 - \frac{3}{x}\right)^{11}$ .

**Q. 7. Show that the coefficient of  $x^4$  in the expansion of  $(1 + 2x + x^2)^5$  is 212.**

**Answer :** To show: that the coefficient of  $x^4$  in the expansion of  $(1 + 2x + x^2)^5$  is 212.

Formula Used:

We have,

$$(1 + 2x + x^2)^5 = (1 + x + x + x^2)^5$$

$$= (1 + x + x(1+x))^5$$

$$= (1 + x)^5(1 + x)^5$$

$$= (1 + x)^{10}$$

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where } s$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term,

$$T_{r+1} = {}^{10}C_r \times x^{10-r} \times (1)^r$$

$$10-r=4$$

$$r=6$$

Thus, the coefficient of  $x^4$  in the expansion of  $(1 + 2x + x^2)^5$  is given by,

$${}^{10}C_4 = \frac{10!}{4!6!}$$

$${}^{10}C_4 = \frac{10 \times 9 \times 8 \times 7 \times 6!}{24 \times 6!}$$

$${}^{10}C_4 = 210$$

Thus, the coefficient of  $x^4$  in the expansion of  $(1 + 2x + x^2)^5$  is 210

**Q. 8. Write the number of terms in the expansion of  $(\sqrt{2} + 1)^5 + (\sqrt{2} - 1)^5$**

**Answer :** To find: the number of terms in the expansion of  $(\sqrt{2} + 1)^5 + (\sqrt{2} - 1)^5$

Formula Used:

Binomial expansion of  $(x + y)^n$  is given by,

$$(x + y)^n = \sum_{r=0}^n {}^nC_r x^{n-r} \times y^r$$

Thus,

$$\begin{aligned}
 & (\sqrt{2} + 1)^5 + (\sqrt{2} - 1)^5 \\
 &= \left( (\sqrt{2})^5 + (\sqrt{2})^4 \binom{5}{1} + \dots + \binom{5}{5} \right) \\
 &+ \left( (\sqrt{2})^5 - (\sqrt{2})^4 \binom{5}{1} + \dots - \binom{5}{5} \right)
 \end{aligned}$$

So, the no. of terms left would be 6

Thus, the number of terms in the expansion of  $(\sqrt{2} + 1)^5 + (\sqrt{2} - 1)^5$  is 6

**Q. 9. Which term is independent of x in the expansion of  $\left(x - \frac{1}{3x^2}\right)^9$  ?**

**Answer :** To find: the term independent of x in the expansion of  $\left(x - \frac{1}{3x^2}\right)^9$  ?

Formula Used:

A general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $\left(x - \frac{1}{3x^2}\right)^9$ , we get



$$T_{r+1} = {}^9C_r \times x^{9-r} \times \left(\frac{-1}{3x^2}\right)^r$$

$$T_{r+1} = {}^9C_r \times x^{9-r} \times (-1)^r \times 3x^{-2r}$$

$$T_{r+1} = {}^9C_r \times (-1)^r \times 3x^{9-3r}$$

For finding the term which is independent of x,

$$9-3r=0$$

$$r=3$$

Thus, the term which would be independent of x is  $T_4$

Thus, the term independent of x in the expansion of  $\left(x - \frac{1}{x}\right)^{10}$  is  $T_4$  i.e 4<sup>th</sup> term

**Q. 10. Write the coefficient of the middle term in the expansion of  $(1 + x)^{2n}$ .**

**Answer :** To find: that the middle term in the expansion of  $\left(\frac{2x^2}{3} + \frac{3}{2x^2}\right)^{10}$  is 252.

Formula Used:

A general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Total number of terms in the expansion is 11

Thus, the middle term of the expansion is  $T_6$  and is given by,

$$T_6 = {}^{10}C_5 \times \left(\frac{2x^2}{3}\right)^5 \times \left(\frac{3}{2x^2}\right)^5$$

$$T_6 = {}^{10}C_5$$

$$T_6 = \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2}$$

$$T_6 = 252$$

Thus, the middle term in the expansion of  $\left(\frac{2x^2}{3} + \frac{3}{2x^2}\right)^{10}$  is 252.

**Q. 11. Write the coefficient of  $x^7y^2$  in the expansion of  $(x + 2y)^9$**

**Answer :** To find: the coefficient of  $x^7y^2$  in the expansion of  $(x + 2y)^9$

Formula Used:

A general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $(x + 2y)^9$ , we get

$$T_{r+1} = {}^9C_r x^{9-r} \times (2y)^r$$

The value of  $r$  for which coefficient of  $x^7y^2$  is defined

$$R = 2$$

Hence, the coefficient of  $x^7y^2$  in the expansion of  $(x + 2y)^9$  is given by:

$$T_3 = {}^9C_3 \times x^{9-2} \times (2y)^2$$

$$T_3 = {}^9C_3 \times 4 \times x^7 \times (y)^2$$

$$T_3 = \frac{9!}{3! \times 6!} \times 4 \times x^7 \times (y)^2$$

$$T_3 = \frac{9 \times 8 \times 7 \times 6!}{6 \times 6!} \times 4 \times x^7 \times (y)^2$$

$$T_3 = 336$$

Thus, the coefficient of  $x^7y^2$  in the expansion of  $(x + 2y)^9$  is 336.

**Q. 12. If the coefficients of  $(r - 5)$ th and  $(2r - 1)$ th terms in the expansion of  $(1 + x)^{34}$  are equal, find the value of  $r$ .**

**Answer :** To find: the value of  $r$  with respect to the binomial expansion of  $(1 + x)^{34}$  where the coefficients of the  $(r - 5)$ th and  $(2r - 1)$ th terms are equal to each other

Formula Used:

The general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the  $(r - 5)$ th term, we get

$$T_{r-5} = {}^{34}C_{r-6} \times x^{r-6}$$

Thus, the coefficient of  $(r - 5)$ th term is  ${}^{34}C_{r-6}$

Now, finding the  $(2r - 1)$ th term, we get

$$T_{2r-1} = {}^{34}C_{2r-2} \times (x)^{2r-2}$$

Thus, coefficient of  $(2r - 1)$ th term is  ${}^{34}C_{2r-2}$

As the coefficients are equal, we get

$${}^{34}C_{2r-2} = {}^{34}C_{r-6}$$

$$2r - 2 = r - 6$$

$$R = -4$$

Value of  $r = -4$  is not possible

$$2r - 2 + r - 6 = 34$$

$$3r = 42$$

$$R = 14$$

Thus, value of  $r$  is 14

**Q. 13. Write the 4<sup>th</sup> term from the end in the expansion of**  $\left(\frac{3}{x^2} - \frac{x^3}{6}\right)^7$

**Answer :** To find: 4<sup>th</sup> term from the end in the expansion of  $\left(\frac{3}{x^2} - \frac{x^3}{6}\right)^7$

Formula Used:

A general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Total number of terms in the expansion is 8

Thus, the 4<sup>th</sup> term of the expansion is  $T_5$  and is given by,

$$T_5 = {}^7C_5 \times \left(\frac{3}{x^2}\right)^3 \times \left(\frac{-x^3}{6}\right)^4$$

$$T_5 = \frac{7 \times 6 \times 5!}{2 \times 5!} \times \frac{3 \times 3 \times 3}{6 \times 6 \times 6 \times 6} \times X^{-18}$$

$$T_5 = \frac{7 \times 6 \times 5!}{2 \times 5!} \times \frac{3 \times 3 \times 3}{6 \times 6 \times 6 \times 6} \times X^{-18}$$

$$T_5 = \frac{7}{16} X^{-18}$$

Thus, a 4<sup>th</sup> term from the end in the expansion of  $\left(\frac{3}{x^2} - \frac{x^3}{6}\right)^7$  is  $T_5 = \frac{7}{16} X^{-18}$

**Q. 14. Find the coefficient of  $x^n$  in the expansion of  $(1 + x)(1 - x)^n$ .**

**Answer :** To find: the coefficient of  $x^n$  in the expansion of  $(1 + x)(1 - x)^n$ .

Formula Used:

Binomial expansion of  $(x + y)^n$  is given by,

$$(x + y)^n = \sum_{r=0}^n \binom{n}{r} x^{n-r} \times y^r$$

Thus,

$$\begin{aligned} (1 + x)(1 - x)^n &= (1 + x) \left( \binom{n}{0} (-x) + \binom{n}{1} (-x)^1 \right. \\ &\quad \left. + \binom{n}{2} (-x)^2 + \dots + \binom{n}{n-1} (-x)^{n-1} + \binom{n}{n} (-x)^n \right) \end{aligned}$$

Thus, the coefficient of  $(x)^n$  is,

${}^nC_n - {}^nC_{n-1}$  (If n is even)

$-{}^nC_n + {}^nC_{n-1}$  (If n is odd)

Thus, the coefficient of  $(x)^n$  is,  ${}^nC_n - {}^nC_{n-1}$  (If n is even) and  $-{}^nC_n + {}^nC_{n-1}$  (If n is odd)

**Q. 15. In the binomial expansion of  $(a + b)^n$ , the coefficients of the 4<sup>th</sup> and 13<sup>th</sup> terms are equal to each other. Find the value of n.**

**Answer :** To find: the value of n with respect to the binomial expansion of  $(a + b)^n$  where the coefficients of the 4<sup>th</sup> and 13<sup>th</sup> terms are equal to each other

Formula Used:

A general term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$T_{r+1} = {}^nC_r x^{n-r} y^r$  where

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the 4<sup>th</sup> term, we get

$$T_4 = {}^nC_3 \times a^{n-3} \times (b)^3$$

Thus, the coefficient of a 4<sup>th</sup> term is  ${}^nC_3$

Now, finding the 13<sup>th</sup> term, we get

$$T_{13} = {}^nC_{12} \times a^{n-12} \times (b)^{12}$$

Thus, coefficient of 4<sup>th</sup> term is  ${}^nC_{12}$

As the coefficients are equal, we get

$${}^nC_{12} = {}^nC_3$$

$$\text{Also, } {}^nC_r = {}^nC_{n-r}$$

$${}^nC_{n-12} = {}^nC_3$$

$$n-12=3$$

$$n=15$$

Thus, value of n is 15

**Q. 16. Find the positive value of m for which the coefficient of  $x^2$  in the expansion of  $(1 + x)^m$  is 6.**

**Answer :** To find: the positive value of m for which the coefficient of  $x^2$  in the expansion of  $(1 + x)^m$  is 6.

Formula Used:

General term,  $T_{r+1}$  of binomial expansion  $(x + y)^n$  is given by,

$$T_{r+1} = {}^nC_r x^{n-r} y^r \text{ where}$$

$${}^nC_r = \frac{n!}{r!(n-r)!}$$

Now, finding the general term of the expression,  $(1 + x)^m$ , we get

$$T_{r+1} = {}^mC_r \times 1^{m-r} \times (x)^r$$

$$T_{r+1} = {}^mC_r \times (x)^r$$

The coefficient of  $(x)^2$  is  ${}^mC_2$

$${}^mC_2 = 6$$

$$\frac{m!}{2(m-2)!} = 6$$

$$\frac{m(m-1)(m-2)!}{2(m-2)!} = 6$$

$$m^2 - m - 6 = 0$$

$$(m-3)(m+2) = 0$$

$$m=3, -2$$

Since m cannot be negative. Therefore,

$$m=3$$

Thus, positive value of  $m$  is 3 for which the coefficient of  $x^2$  in the expansion of  $(1 + x)^m$  is 6