

CBSE Test Paper 01
Chapter 10 Vector Algebra

1. Find the angle between two vectors \vec{a} and \vec{b} with magnitudes $\sqrt{3}$ and 2, respectively, having $\vec{a} \cdot \vec{b} = \sqrt{6}$.
 - a. $\frac{\pi}{5}$
 - b. $\frac{\pi}{3}$
 - c. $\frac{\pi}{2}$
 - d. $\frac{\pi}{4}$
2. Find the angle between two vectors $\hat{i} - 2\hat{j} + 3\hat{k}$ and $3\hat{i} - 2\hat{j} + \hat{k}$.
 - a. $\cos^{-1}\left(\frac{4}{7}\right)$
 - b. $\cos^{-1}\left(\frac{6}{7}\right)$
 - c. $\cos^{-1}\left(\frac{5}{9}\right)$
 - d. $\cos^{-1}\left(\frac{5}{7}\right)$
3. Vector has
 - a. direction
 - b. None of these
 - c. magnitude
 - d. magnitude as well as direction
4. Find the sum of the vectors $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$ and $\vec{c} = \hat{i} - 6\hat{j} - 7\hat{k}$.
 - a. $-\hat{i} + 4\hat{j} - \hat{k}$
 - b. $-4\hat{j} - \hat{k}$
 - c. $-\hat{i} - 4\hat{j} - \hat{k}$
 - d. $\hat{i} - 4\hat{j} - \hat{k}$
5. Find the direction cosines of the vector $\hat{i} + 2\hat{j} + 3\hat{k}$.

- a. $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$
- b. $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, -\frac{3}{\sqrt{14}}$
- c. $\frac{1}{\sqrt{14}}, -\frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$
- d. $-\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$

6. The values of k which $|k\vec{a}| < |\vec{a}|$ and $k\vec{a} + \frac{1}{2}\vec{a}$ is parallel to \vec{a} holds true are _____.
7. If $\vec{r} \cdot \vec{a} = 0, \vec{r} \cdot \vec{b} = 0$, and $\vec{r} \cdot \vec{c} = 0$ for some non-zero vector \vec{r} , then the value of $\vec{a}(\vec{b} \times \vec{c})$ is _____.
8. The angle between two vectors \vec{a} and \vec{b} with magnitudes $\sqrt{3}$ and 4, respectively, $\vec{a} \cdot \vec{b} = 2\sqrt{3}$ is _____.
9. Find $\vec{a} \times \vec{b}$ if $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} + 5\hat{j} - 2\hat{k}$.
10. Find the projection of \vec{a} on \vec{b} , if $\vec{a} \cdot \vec{b} = 8$ and $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$.
11. \vec{a} Is unit vector and $(\vec{x} - \vec{a})(\vec{x} + \vec{a}) = 8$, Then find $|\vec{x}|$.
12. Find the position vector of the mid-point of the vector joining the points P (2, 3, 4) and Q(4,1, - 2)
13. Find sine of the angle between the vectors. $\vec{a} = 2\hat{i} - \hat{j} + 3\hat{k}, \vec{b} = \hat{i} + 3\hat{j} + 2\hat{k}$.
14. Find the projection of the vector $\hat{i} + 3\hat{j} + 7\hat{k}$ on the vector $7\hat{i} - \hat{j} + 8\hat{k}$
15. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$ and $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$. Find a vector of magnitude 6 units, which is parallel to the vector $2\vec{a} - \vec{b} + 3\vec{c}$.
16. Let $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$. Find a vector \vec{d} which is perpendicular to both \vec{a} and \vec{b} and $\vec{c} \cdot \vec{d} = 15$.
17. A girl walks 4 km towards west, then she walks 3 km in a direction 30° east of north and stops. Determine the girl's displacement from her initial point of departure.
18. Find a vector \vec{d} which is \perp to both \vec{a} and \vec{b} and $\vec{c} \cdot \vec{d} = 15$ Let $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$.

CBSE Test Paper 01
Chapter 10 Vector Algebra

Solution

1. d. $\frac{\pi}{4}$, **Explanation:** $|\vec{a}| = \sqrt{3}, |\vec{b}| = 2, \vec{a} \cdot \vec{b} = \sqrt{6}$
 $\Rightarrow \vec{a} \cdot \vec{b} = |\vec{a}| \cdot |\vec{b}| \cos \theta \Rightarrow \sqrt{6}$
 $= 2\sqrt{3} \cos \theta$
 $\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4}$
2. d. $\cos^{-1}\left(\frac{5}{7}\right)$, **Explanation:** $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j} + \hat{k}$
 $\Rightarrow |\vec{a}| = \sqrt{14}, |\vec{b}| = \sqrt{14}, \vec{a} \cdot \vec{b} = 10$
 $\Rightarrow \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \cos \theta \Rightarrow \frac{10}{14} = \cos \theta$
 $\Rightarrow \cos \theta = \frac{5}{7} \Rightarrow \theta = \cos^{-1} \frac{5}{7}$
3. d. magnitude as well as direction, **Explanation:** A vector has both magnitude as well as direction.
4. b. $-4\hat{j} - \hat{k}$, **Explanation:** We have: vectors $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$,
 $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$ and
5. a. $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$, **Explanation:** Let $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$,
Then, $\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{1^2 + 2^2 + 3^2}} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}}$
Therefore, the D.C.'s of vector a are :
 $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$.
6. $k \in]-1, 1 [$ $k \neq -\frac{1}{2}$
7. 0
8. $\frac{\pi}{3}$

$$\begin{aligned}
 9. \quad \vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 3 \\ 3 & 5 & -2 \end{vmatrix} \\
 &= \hat{i}(-2-15) - \hat{j}(-4-9) + \hat{k}(10-3) \\
 &= -17\hat{i} + 13\hat{j} + 7\hat{k}
 \end{aligned}$$

10. We are given that, $\vec{a} \cdot \vec{b} = 8$ and $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$

\therefore The projection of \vec{a} on \vec{b} is given as $= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

$$\begin{aligned}
 &= \frac{8}{\sqrt{2^2+6^2+3^2}} \\
 &= \frac{8}{\sqrt{4+36+9}} \\
 &= \frac{8}{\sqrt{49}} = \frac{8}{7}
 \end{aligned}$$

11. $|\vec{a}| = 1$

$$(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 8$$

$$|\vec{x}|^2 - |\vec{a}|^2 = 8$$

$$|\vec{x}|^2 - 1 = 8$$

$$|\vec{x}|^2 = 9$$

$$|\vec{x}| = 3$$

12. Given: Point P (2, 3, 4) and Q(4,1, - 2)

\therefore Position vector of point P is $\vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$

And Position vector of point Q is $\vec{b} = 4\hat{i} + \hat{j} - 2\hat{k}$

And Position vector of mid-point R of PQ is $\frac{\vec{a} + \vec{b}}{2} = \frac{2\hat{i} + 3\hat{j} + 4\hat{k} + 4\hat{i} + \hat{j} - 2\hat{k}}{2}$

$$= \frac{6\hat{i} + 4\hat{j} + 2\hat{k}}{2} = 3\hat{i} + 2\hat{j} + \hat{k}$$

$$13. \quad \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 3 \\ 1 & 3 & 2 \end{vmatrix}$$

$$= -11\hat{i} - \hat{j} + 7\hat{k}$$

$$|\vec{a} \times \vec{b}| = \sqrt{(-11)^2 + (-1)^2 + (7)^2}$$

$$= \sqrt{171} = 3\sqrt{19}$$

$$\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}||\vec{b}|} = \frac{3\sqrt{19}}{\sqrt{14}\sqrt{14}} = \frac{3}{14}\sqrt{19}$$

14. Let $\vec{a} = \hat{i} + 3\hat{j} + 7\hat{k}$ and $\vec{b} = 7\hat{i} - \hat{j} + 8\hat{k}$

$$\begin{aligned}
 \text{Projection of vector } \vec{a} \text{ on } \vec{b} &= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \\
 &= \frac{(1)(7) + (3)(-1) + 7(8)}{\sqrt{(7)^2 + (-1)^2 + (8)^2}} \\
 &= \frac{7 - 3 + 56}{\sqrt{49 + 61 + 64}} = \frac{60}{\sqrt{114}}
 \end{aligned}$$

15. According to the question ,

$$\vec{a} = \hat{i} + \hat{j} + \hat{k},$$

$$\vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k} \text{ and}$$

$$\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$$

$$\text{Now, } 2\vec{a} - \vec{b} + 3\vec{c}$$

$$= 2(\hat{i} + \hat{j} + \hat{k}) - (4\hat{i} - 2\hat{j} + 3\hat{k}) + 3(\hat{i} - 2\hat{j} + \hat{k})$$

$$= 2\hat{i} + 2\hat{j} + 2\hat{k} - 4\hat{i} + 2\hat{j} - 3\hat{k} + 3\hat{i} - 6\hat{j} + 3\hat{k}$$

$$= \hat{i} - 2\hat{j} + 2\hat{k}$$

$$\Rightarrow 2\vec{a} - \vec{b} + 3\vec{c} = \hat{i} - 2\hat{j} + 2\hat{k}$$

$$\text{Now, a unit vector in the direction of vector is } 2\vec{a} - \vec{b} + 3\vec{c} = \frac{2\vec{a} - \vec{b} + 3\vec{c}}{|2\vec{a} - \vec{b} + 3\vec{c}|}$$

$$= \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{(1)^2 + (-2)^2 + (2)^2}}$$

$$= \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{9}}$$

$$= \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3}$$

$$= \frac{1}{3}\hat{i} - \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k}$$

Vector of magnitude 6 units parallel to the vector is ,

$$= 6 \left(\frac{1}{3}\hat{i} - \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k} \right)$$

$$= 2\hat{i} - 4\hat{j} + 4\hat{k}$$

16. Given: Vectors $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$

We know that the cross-product of two vectors, $\vec{a} \times \vec{b}$ is a vector perpendicular to both \vec{a} and \vec{b}

Hence, vector \vec{d} which is also perpendicular to both \vec{a} and \vec{b} is $\vec{d} = \lambda (\vec{a} \times \vec{b})$ where $\lambda = 1$ or some other scalar.

$$\text{Therefore, } \vec{d} = \lambda \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 4 & 2 \\ 3 & -2 & 7 \end{vmatrix}$$

$$= \lambda [\hat{i}(28+4) - \hat{j}(7-6) + \hat{k}(-2-12)]$$

$$\Rightarrow \vec{d} = 32\lambda\hat{i} - \lambda\hat{j} - 14\lambda\hat{k} \dots (i)$$

Now given $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$ and $\vec{c} \cdot \vec{d} = 15$

$$\vec{c} \cdot \vec{d} = 15$$

$$= 2(32\lambda) + (-1)(-\lambda) + 4(-14\lambda) = 15$$

$$\Rightarrow 64\lambda + \lambda - 56\lambda = 15$$

$$\Rightarrow 9\lambda = 15$$

$$\Rightarrow \lambda = \frac{15}{9}$$

$$\Rightarrow \lambda = \frac{5}{3}$$

Putting $\lambda = \frac{5}{3}$ in eq. (i), we get

$$\vec{d} = \frac{5}{3} [32\hat{i} - \hat{j} - 14\hat{k}]$$


$$\Rightarrow \vec{d} = \frac{1}{3} [160\hat{i} - 5\hat{j} - 70\hat{k}]$$

17. Let the initial point of departure is origin (0, 0) and the girl walks a distance OA = 4 km towards west.

Through the point A, draw a line AQ parallel to a line OP, which is 30° East of North, i.e., in East-North quadrant making an angle of 30° with North.

Again, let the girl walks a distance AB = 3 km along this direction \vec{OQ}

$$\therefore \vec{OA} = 4(-\hat{i}) = -4\hat{i} \dots (i) \quad [\because \text{Vector } \vec{OA} \text{ is along } OX']$$



Now, draw BM perpendicular to x - axis.

In $\triangle AMB$ by Triangle Law of Addition of vectors,

$$\vec{AB} = \vec{AM} + \vec{MB} = (AM)\hat{i} + (MB)\hat{i}$$

Dividing and multiplying by AB in R.H.S.,

$$\overrightarrow{AB} = AB \frac{AM}{AB} \hat{i} + AB \frac{MB}{AB} \hat{j} = 3 \cos 60^\circ \hat{i} + 3 \sin 60^\circ \hat{j}$$

$$\Rightarrow AB = 3 \frac{1}{2} \hat{i} + 3 \frac{\sqrt{3}}{2} \hat{j} = \frac{3}{2} \hat{i} + \frac{3\sqrt{3}}{2} \hat{j} \dots (ii)$$

\therefore Girl's displacement from her initial point O of departure to final point B,

$$\overrightarrow{OB} = \overrightarrow{OA} + \overrightarrow{AB} = -4\hat{i} + \left(\frac{3}{2} \hat{i} + \frac{3\sqrt{3}}{2} \hat{j} \right) = \left(-4 + \frac{3}{2} \right) \hat{i} + \frac{3\sqrt{3}}{2} \hat{j}$$

$$\Rightarrow \overrightarrow{OB} = -\frac{5}{2} \hat{i} + \frac{3\sqrt{3}}{2} \hat{j}$$

18. $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$

Let $\vec{d} = x\hat{i} + y\hat{j} + z\hat{k}$

ATQ, $\vec{d} \cdot \vec{a} = 0$, $\vec{d} \cdot \vec{b} = 0$ and $\vec{c} \cdot \vec{d} = 15$, then,

$$x + 4y + 2z = 0 \dots (1)$$

$$3x - 2y + 7z = 0 \dots (2)$$

$$2x - y + 4z = 15 \dots (3)$$

On solving equation (1) and (2)

$$\frac{x}{\begin{array}{cc} 4 & 2 \\ -2 & 7 \end{array}} = \frac{y}{\begin{array}{cc} 2 & 1 \\ 7 & 3 \end{array}} = \frac{z}{\begin{array}{cc} 1 & 4 \\ 3 & -2 \end{array}} = K$$

$$\frac{x}{28-4} = \frac{y}{6-7} = \frac{z}{-2-12} = k$$

$$x = 32k, y = -k, z = -14k$$

Put x, y, z in equation (3)

$$2(32k) - (-k) + 4(-14k) = 15$$

$$64k + k - 56k = 15$$

$$9k = 15$$

$$k = \frac{15}{9}$$

$$k = \frac{5}{3}$$

$$x = 32 \times \frac{5}{3} = \frac{160}{3}$$

$$y = -\frac{5}{3}$$

$$z = -14 \times \frac{5}{3} = -\frac{70}{3}$$

$$\vec{d} = \frac{160}{3} \hat{i} - \frac{5}{3} \hat{j} - \frac{70}{3} \hat{k}$$