

# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 10


A quantity that has both magnitude and direction is called a vector.  
The distance between the initial and terminal points of a vector is called its magnitude. Magnitude of vector  $\overrightarrow{AB}$  is  $|AB|$ .


Position vector of a point  $P(x, y, z)$  is  $x\hat{i} + y\hat{j} + z\hat{k}$  and its magnitude is  $OP(r) = \sqrt{x^2 + y^2 + z^2}$ . For eg: Position vector of  $P(2, 3, 5)$  is  $2\hat{i} + 3\hat{j} + 5\hat{k}$  and its magnitude is  $\sqrt{2^2 + 3^2 + 5^2} = \sqrt{38}$ .

The scalar components of a vector are its direction ratios, and represent its projections along the respective axes.  
The magnitude ( $r$ ) direction ratios ( $a, b, c$ ) and direction cosines ( $l, m, n$ ) of vector  $a\hat{i} + b\hat{j} + c\hat{k}$  are related as:  
 $l = \frac{a}{r}, m = \frac{b}{r}, n = \frac{c}{r}$   
For eg: If  $\overrightarrow{AB} = \hat{i} + 2\hat{j} + 3\hat{k}$ , then  $r = \sqrt{1 + 4 + 9} = \sqrt{14}$   
Direction ratios are  $(1, 2, 3)$   $\left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$   
and direction cosines are  $\left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

- (i) Zero vector (initial and terminal points coincide)
- (ii) Unit vector (magnitude is unity)
- (iii) Coinitial vectors (same initial points)
- (iv) Collinear vectors (parallel to the same line)
- (v) Equal vectors (same magnitude and direction)
- (vi) Negative of a vector (same magnitude, opp. direction)

The vector sum of the three sides of a triangle taken in order is 0 i.e  
if  $ABC$  is given triangle, then  
 $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0$ .



The Vector sum of two coinitial vectors is given by the diagonal of the parallelogram whose adjacent sides are given vectors.  
  
if  $\overrightarrow{AB}, \overrightarrow{AC}$  are the given vectors, then  $\overrightarrow{AB} + \overrightarrow{AC} = \overrightarrow{AD}$

If we have two vectors  
 $a = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ ,  $b = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  and  $\lambda$  is any scalar, then-  
 $a + b = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}$   
 $\lambda a = (\lambda a_1)\hat{i} + (\lambda a_2)\hat{j} + (\lambda a_3)\hat{k}$   
 $a \cdot b = a_1b_1 + a_2b_2 + a_3b_3$  and  
 $a \times b = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$

$a \times b = |a||b|\sin\theta\hat{n}$ ,  $\hat{n}$  is a unit vector perpendicular to line joining  $a, b$ .

If  $a, b$  are the vectors and ' $\theta$ ', angle between them, then their scalar product  $a \cdot b = |a||b|\cos\theta$   
 $\Rightarrow \cos\theta = \frac{a \cdot b}{|a||b|}$

The Position vector of a point  $R$  dividing a line segment joining  $P, Q$  whose position vectors are  $a, b$  resp., in the ratio  $m : n$   
(i) internally is  $\frac{na + mb}{m + n}$ , (ii) externally is  $\frac{mb - na}{m - n}$

For a given vector  $a$ , the vector  $\hat{a} = \frac{a}{|a|}$  gives the unit vector in the direction of  $a$ . for eg, if  $a = 5\hat{i}$ , then  $\hat{a} = \frac{5\hat{i}}{|5|} = \hat{i}$ , which is a unit vector.

If  $\lambda$  Multiplied to vector  $AB$ , then the magnitude is multiplied by  $|\lambda|$  and direction remain same (or opp.) according as  $\lambda$  is the +ve or, -ve.

### Vector Algebra

Multiplication of vector by a scalar  
Unit vector  
Position of vectors

Vector

Position Vector  
Direction ratios and direction cosines

Types of Vectors

Properties of Vector

Scalar product of two vectors  
Cross product of two vectors