

**Topics : Magnetic Effect of Current and Magnetic Force on Charge/current, Electromagnet Induction, Rotation, Center of Mass, Geometrical Optics, Current Electricity**

**Type of Questions**

**Single choice Objective ('-1' negative marking) Q.1 to Q.4**

**(3 marks, 3 min.)**

**M.M., Min.**

**[12, 12]**

**Subjective Questions ('-1' negative marking) Q.5 to Q.6**

**(4 marks, 5 min.)**

**[8, 10]**

**Comprehension ('-1' negative marking) Q.7 to Q.9**



**(3 marks, 3 min.)**

**[9, 9]**

1. The direction of the field B at P is :

The V shaped wire is in x-y plane.


- (A) along + x-axis
- (B) along + z-axis
- (C) along (-x)-axis
- (D) along + y-axis



2. If the magnetic field at 'P' can be written as  $K \tan\left(\frac{\alpha}{2}\right)$  then K is :  
[Refer to the figure of above question ]

- (A)  $\frac{\mu_0 I}{4\pi d}$
- (B)  $\frac{\mu_0 I}{2\pi d}$
- (C)  $\frac{\mu_0 I}{\pi d}$
- (D)  $\frac{2\mu_0 I}{\pi d}$

3. A circular loop of radius r is moved with a velocity v as shown in the diagram. The work needed to maintain its velocity constant is :



- (A)  $\frac{\mu_0 i v r}{2\pi a}$
- (B)  $\frac{\mu_0 i v r}{2\pi(a+r)}$
- (C)  $\frac{\mu_0 i v r}{2\pi} \ln\left(\frac{2r+a}{a}\right)$
- (D) zero

4. The magnifying power of a simple microscope can be increased if an eyepiece of :


- (A) shorter focal length is used
- (B) longer focal length is used
- (C) shorter diameter is used
- (D) longer diameter is used

5. A rod of negligible mass and length  $\ell$  is pivoted at its centre. A particle of mass m is fixed to its left end & another particle of mass 2m is fixed to the right end. If the system is released from rest,




- (a) what is the speed v of the two masses when the rod is vertical.
- (b) what is the angular speed  $\omega$  of the system at that instant.

6. A ball is given velocity  $\sqrt{3g\ell}$  as shown. If the ratio of centripetal acceleration to tangential acceleration is  $1 : y\sqrt{2}$  at the point where the ball leaves circular path then write the value of  $y$ . [Neglect the size of ball]



### COMPREHENSION

A fixed cylindrical tank having large cross-section area is filled with two liquids of densities  $\rho$  and  $2\rho$  and in equal volumes as shown in the figure. A small hole of area of cross-section  $a = \sqrt{6}\text{ cm}^2$  is made at height  $h/2$  from the bottom.




7. Velocity of efflux will be :  
 (A)  $\sqrt{2gh}$       (B)  $\sqrt{3gh}$       (C)  $\sqrt{gh}$       (D)  $2\sqrt{gh}$
8. Distance (R) of the point at which the liquid will strike from container is :  
 (A)  $2h$       (B)  $h$       (C)  $\frac{h}{2}$       (D)  $\sqrt{2}h$
9. Area of cross section of stream of liquid just before it hits the ground.  
 (A)  $2\text{ cm}^2$       (B)  $\sqrt{3}\text{ cm}^2$       (C)  $1\text{ cm}^2$       (D)  $\sqrt{5}\text{ cm}^2$

## Answers Key

1. (B)      2. (B)      3. (D)      4. (A)  
 5. (a)  $V = \sqrt{g\ell/3}$ ,  $\omega = \sqrt{4g/3\ell}$  ]  
 6.  $y = 2$       7. (A)      8. (D)      9. (A)

# Hints & Solutions

1. By right hand thumb rule, the field by both the segments are out of the plane i.e. along **+ve z-axis**.
2. Let us compute the magnetic field due to any one segment :



$$B = \frac{\mu_0 i}{4\pi(d \sin \alpha)} (\cos 0^\circ + \cos(180 - \alpha))$$

$$= \frac{\mu_0 i}{4\pi(d \sin \alpha)} (1 - \cos \alpha) = \frac{\mu_0 i}{4\pi d} \tan \frac{\alpha}{2}$$

Resultant field will be :

$$B_{\text{net}} = 2B = \frac{\mu_0 i}{2\pi d} \tan \frac{\alpha}{2} \Rightarrow k = \frac{\mu_0 i}{2\pi d}$$

3. Due to the motion of the loop, there will be an induced current flowing in the circuit, resulting in a force acting on each element of the loop equally & radially. Therefore the net force on the loop is zero.

Hence (D).

5. Decrease in PE =

Increase in rotation K.E


$$\Rightarrow 2mg \cdot \frac{\ell}{2} - mg \cdot \frac{\ell}{2} = \frac{1}{2} I \omega^2$$

$$= \frac{1}{2} \left( 2m \frac{\ell^2}{4} + m \cdot \frac{\ell^2}{4} \right) \omega^2$$

$$\frac{mg\ell}{2} = \frac{1}{2} \cdot \frac{3m\ell^2}{4} \cdot \omega = \frac{3m\ell^2}{8} \omega^2$$

$$\omega = \sqrt{\frac{4g}{3\ell}} \quad \text{and } v = r\omega = \frac{\ell}{2} \sqrt{\frac{4g}{3\ell}} = \sqrt{\frac{g\ell}{3}}$$

[ Ans.: (a)  $V = \sqrt{g\ell/3}$ ,  $\omega = \sqrt{4g/3\ell}$  ]



$$mg \cos \theta + T_1 = \frac{mv_1^2}{\ell}$$

for leaving circle  $T_1 = 0$

$$mv_1^2 = mg \ell \cos \theta \quad \dots(i)$$

and by energy conservation

$$0 + \frac{1}{2} m (\sqrt{3g\ell})^2 = \frac{1}{2} mv_1^2 + mg (\ell + \ell \cos \theta)$$

$$\frac{1}{2} m (3g\ell) = \frac{1}{2} mv_1^2 + mg\ell (1 + \cos \theta)$$

$$\frac{3mg\ell}{2} = \frac{mg\ell \cos \theta}{2} + mg\ell + mg\ell \cos \theta$$

(by equation (i))

$$\frac{mg\ell}{2} = \frac{3}{2} mg\ell \cos \theta$$

$$\cos \theta = \frac{1}{3}$$

$$\sin \theta = \sqrt{1 - \frac{1}{9}} = \frac{\sqrt{8}}{3}$$

$$a_c = \frac{v_1^2}{\ell} = \frac{g\ell \cos \theta}{\ell} = g \cos \theta$$

$$a_t = g \sin \theta$$

then  $\frac{a_c}{a_t}$

$$= \frac{g \cos \theta}{g \sin \theta} = \frac{1/3}{\sqrt{8}/3} = \frac{1}{\sqrt{8}} = \frac{1}{2\sqrt{2}} = \frac{1}{y\sqrt{2}}$$

so  $y = 2$

**Ans.  $y = 2$**

**Sol. (7 to 9)**

Applying Bernoulli's equation

$$P_0 + \frac{1}{2} \times 2\rho \times V^2 = P_0 + 2\rho g \times \frac{h}{2} + \rho gh$$

$$v = \sqrt{2gh}$$

$$\frac{1}{2} \times g \times t^2 = \frac{h}{2}$$

$$\Rightarrow t = \sqrt{\frac{h}{g}}$$

$$R = v \times t$$

$$\Rightarrow \sqrt{2h}$$

Applying continuity equation

$$\sqrt{6} \times \sqrt{2gh} = \sqrt{3gh} \times A$$

$$A = 2\text{cm}^2$$