

# Three Dimensional Geometry

Question 1.

The angle between the line  $\frac{x-5}{7} = \frac{y+2}{-5} = \frac{z}{1}$  and

$\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$  is

- (a) 0      (b)  $\frac{\pi}{2}$       (c)  $\frac{\pi}{3}$       (d)  $\frac{\pi}{4}$

Answer:

- (b)  $\frac{\pi}{2}$

Question 2.

The angle between the lines passing through the points (4, 7, 8), (2, 3, 4) and (-1, -2, 1), (1, 2, 5) is

- (a) 0  
(b)  $\frac{\pi}{2}$   
(c)  $\frac{\pi}{4}$   
(d)  $\frac{\pi}{6}$

Answer:

- (a) 0

Question 3.

**Find the equation of line passing through the origin**

**which intersect the line  $\frac{x-3}{2} = \frac{y-3}{1} = \frac{z-0}{1}$  at angle**

$$\frac{\pi}{3}.$$

(a)  $\frac{x}{1} = \frac{y}{2} = \frac{z}{-1}$

(b)  $\frac{x}{-1} = \frac{y}{1} = \frac{z}{-2}$

(c)  $\frac{x}{1} = \frac{y}{3} = \frac{z}{2}$

(d) Both (a) and (b)

Answer:

(d) Both (a) and (b)

Question 4.

**Equation of a line passing through (1, 2, -3) and parallel**

**to the line  $\frac{x-2}{1} = \frac{y+1}{3} = \frac{z-1}{4}$  is**

(a)  $\frac{x-1}{1} = \frac{y-2}{3} = \frac{z+3}{4}$  (b)  $\frac{x-2}{1} = \frac{y+1}{2} = \frac{z-1}{-3}$

(c)  $\frac{x-1}{1} = \frac{y-3}{2} = \frac{z-4}{-3}$  (d) None of these

Answer:

(a)  $\frac{x-1}{1} = \frac{y-2}{3} = \frac{z+3}{4}$

Question 5.

**If lines  $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$  and  $\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5}$**

**are mutually perpendicular, then  $k$  is equal to**

(a)  $-\frac{10}{7}$  (b)  $-\frac{7}{10}$  (c) -10 (d) -7

Answer:

(a)  $-\frac{10}{7}$

Question 6.

Equation of the plane passing through three points A, B, C with position vectors

$$-6\hat{i} + 3\hat{j} + 2\hat{k}, 3\hat{i} - 2\hat{j} + 4\hat{k}, 5\hat{i} + 7\hat{j} + 3\hat{k}$$

(a)  $\vec{r} \cdot (\hat{i} - \hat{j} - 2\hat{k}) + 23 = 0$

(b)  $\vec{r} \cdot (\hat{i} + \hat{j} + 7\hat{k}) = 23$

(c)  $\vec{r} \cdot (\hat{i} + \hat{j} - 7\hat{k}) + 23 = 0$

(d)  $\vec{r} \cdot (\hat{i} - \hat{j} - 7\hat{k}) = 23$

Answer:

(a)  $\pi(\hat{i} - \hat{j} - 2\hat{k}) + 23 = 0$

Question 7.

Four points (0, -1, -1) (-4, 4, 4) (4, 5, 1) and (3, 9, 4) are coplanar. Find the equation of the plane containing them.

(a)  $5x + 7y + 11z - 4 = 0$

(b)  $5x - 7y + 11z + 4 = 0$

(c)  $5x - 7y - 11z - 4 = 0$

(d)  $5x + 7y - 11z + 4 = 0$

Answer:

(b)  $5x - 7y + 11z + 4 = 0$

Question 8.

Find the equation of plane passing through the points P(1, 1, 1), Q(3, -1, 2), R(-3, 5, -4).

(a)  $x + 2y = 0$

(b)  $x - y = 2$

(c)  $-x + 2y = 2$

(d)  $x + y = 2$

Answer:

(d)  $x + y = 2$

Question 9.

The vector equation of the plane passing through the origin and the line of intersection of the plane  $\vec{r} \cdot \vec{a} = \lambda$  and  $\vec{r} \cdot \vec{b} = \mu$  is

(a)  $\vec{r} \cdot (\lambda \vec{a} - \mu \vec{b}) = 0$

(b)  $\vec{r} \cdot (\lambda \vec{b} - \mu \vec{a}) = 0$

(c)  $\vec{r} \cdot (\lambda \vec{a} + \mu \vec{b}) = 0$

(d)  $\vec{r} \cdot (\lambda \vec{b} + \mu \vec{a}) = 0$

Answer:

(b)  $\vec{r} \cdot (\lambda \vec{b} - \mu \vec{a}) = 0$

Question 10.

The vector equation of a plane passing through the intersection of the planes  $r \cdot (\hat{i} + \hat{j} + \hat{k}) = 6$  and  $r \cdot (2\hat{i} + 3\hat{j} + 4\hat{k}) = -5$  and the point  $(1, 1, 1)$  is

- (a)  $\vec{r} \cdot (3\hat{i} + 4\hat{j} + 5\hat{k}) = 1$  (b)  $\vec{r} \cdot (8\hat{i} + 5\hat{j} + 2\hat{k}) = 99$   
(c)  $\vec{r} \cdot (20\hat{i} + 23\hat{j} + 26\hat{k}) = 69$   
(d)  $\vec{r} \cdot (20\hat{i} - 23\hat{j} - 26\hat{k}) = 69$

Answer:

(c)  $r \cdot (20\hat{i} + 23\hat{j} + 26\hat{k}) = 69$

Question 11.

The lines  $\frac{x-1}{2} = \frac{y+1}{-3} = \frac{z+10}{8}$  and

$\frac{x-4}{4} = \frac{y+3}{k} = \frac{z+1}{7}$  are coplanar if  $k =$

- (a) 4 (b) -4  
(c) 2 (d) -2

Answer:

(b) -4

Question 12.

The lines  $\frac{x+3}{-3} = \frac{y-1}{1} = \frac{z-5}{5}$  and  $\frac{x+1}{-1} = \frac{y-2}{2} = \frac{z-5}{5}$

are

- (a) coplanar  
(b) non-coplanar  
(c) perpendicular  
(d) None of the above

Answer:

(a) coplanar

Question 13.

The angle between the planes  $3x + 2y + z - 5 = 0$  and  $x + y - 2z - 3 = 0$  is

(a)  $\cos^{-1}\left(\frac{3}{2\sqrt{7}}\right)$       (b)  $\cos^{-1}\left(\frac{3}{2\sqrt{14}}\right)$

(c)  $\cos^{-1}\left(\frac{3}{2\sqrt{21}}\right)$       (d) None of these

Answer:

(c)  $\cos^{-1}\left(\frac{3}{2\sqrt{21}}\right)$

Question 14.

The equation of the plane through the point (0, -4, -6) and (-2, 9, 3) and perpendicular to the plane  $x - 4y - 2z = 8$  is

(a)  $3x + 3y - 2z = 0$

(b)  $x - 2y + z = 2$

(c)  $2x + y - z = 2$

(d)  $5x - 3y + 2z = 0$

Answer:

(c)  $2x + y - z = 2$

Question 15.

The angle between the planes  $r \cdot (\hat{i} + 2\hat{j} + \hat{k}) = 4$  and  $r(-\hat{i} + \hat{j} + 2\hat{k}) = 9$  is

(a)  $30^\circ$

(b)  $60^\circ$

(c)  $45^\circ$

(d) None of these

Answer:

(b)  $60^\circ$

Question 16.

**The value of p, so that the line**

$$\frac{1-x}{3} = \frac{7y-14}{2p} = \frac{z-3}{2} \text{ and } \frac{7-7x}{3p} = \frac{y-5}{1} = \frac{6-z}{5}$$

**intersect at right angle, is**

(a)  $\frac{10}{11}$       (b)  $\frac{70}{11}$       (c)  $\frac{10}{7}$       (d)  $\frac{70}{9}$

Answer:

(b)  $\frac{70}{11}$

Question 17.

The shortest distance between the lines

$$\vec{r} = -(\hat{i} + \hat{j} + \hat{k})\lambda(2\hat{i} + 3\hat{j} + 4\hat{k}) \text{ and}$$

$$\vec{r} = -\hat{i} + \mu(3\hat{i} + 4\hat{j} + 5\hat{k}) \text{ is}$$

- (a) 1      (b)  $\frac{1}{\sqrt{2}}$       (c)  $\frac{1}{\sqrt{3}}$       (d)  $\frac{1}{\sqrt{6}}$

Answer:

(d)  $\frac{1}{\sqrt{6}}$

Question 18.

The shortest distance between the lines  $\frac{x-3}{3} = \frac{y-8}{-1} = \frac{z-3}{1}$  and  $\frac{x+3}{-3} = \frac{y+7}{2} = \frac{z-6}{4}$  is equal

- (a)  $3\sqrt{30}$   
(b)  $\sqrt{30}$   
(c)  $2\sqrt{30}$   
(d) None of these

Answer:

(a)  $3\sqrt{30}$

Question 19.

The shortest distance between the lines  $x = y = z$  and  $x + 1 - y = \frac{z}{0}$  is

- (a)  $\frac{1}{2}$   
(b)  $\frac{1}{\sqrt{2}}$   
(c)  $\frac{1}{\sqrt{3}}$   
(d)  $\frac{1}{\sqrt{6}}$

Answer:

(d)  $\frac{1}{\sqrt{6}}$

Question 20.

The shortest distance between the lines  $x = y + 2 = 6z - 6$  and  $x + 1 = 2y = -12z$  is

- (a)  $\frac{1}{2}$   
(b) 2  
(c) 1  
(d)  $\frac{3}{2}$

Answer:

(b) 2

Question 21.

The angle  $\theta$  between the line  $r = a + \lambda b$  is given by

- (a)  $\sin^{-1}\left(\frac{\vec{a} \cdot \vec{b} \hat{n}}{|b|}\right)$       (b)  $\cos^{-1}\left(\frac{\vec{a} \cdot \vec{b} \hat{n}}{|b|}\right)$   
(c)  $\sin^{-1}\left(\frac{\vec{a} \cdot \hat{n}}{|a|}\right)$       (d)  $\cos^{-1}\left(\frac{\vec{a} \cdot \hat{n}}{|a|}\right)$

Answer:

(a)  $\sin^{-1}\left(\frac{\vec{a} \cdot \hat{n}}{|b|}\right)$

Question 22.

**Find the angle between the line**  
 $\vec{r} = \hat{i} + 2\hat{j} - \hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$  **and the plane**  
 $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 4$ .

- (a)  $\sin^{-1}\left(\frac{2\sqrt{2}}{3}\right)$       (b)  $\sin^{-1}\left(\frac{2}{3}\right)$   
(c)  $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$       (d)  $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$

Answer:

(a)  $\sin^{-1}\left(\frac{2\sqrt{2}}{3}\right)$

Question 23.

The angle between the straight line  $\frac{x-1}{2} = \frac{y+3}{-1} = \frac{z-5}{2}$  and the plane  $4x - 2y + 4z = 9$  is

- (a)  $60^\circ$   
(b)  $90^\circ$   
(c)  $45^\circ$   
(d)  $30^\circ$

Answer:

(b)  $90^\circ$

Question 24.

Distance of the point  $(\alpha, \beta, \gamma)$  from y-axis is

- (a)  $\beta$

- (b)  $|\beta|$
- (c)  $|\beta| + |\gamma|$
- (d)  $\sqrt{\alpha^2 + \gamma^2}$

Answer:

- (d)  $\sqrt{\alpha^2 + \gamma^2}$

Question 25.

The distance of the plane  $r \cdot \left( \frac{2}{7} \hat{i} + \frac{3}{7} \hat{j} - \frac{6}{7} \hat{k} \right) = 1$  from the origin is

- (a) 1
- (b) 7
- (c)  $\frac{1}{7}$
- (d) None of these

Answer:

- (a) 1

Question 26.

**The sine of the angle between the straight line**

$\frac{x-2}{3} = \frac{y-3}{4} = \frac{z-4}{5}$  **and the plane**  $2x - 2y + z = 5$  **is**

- (a)  $\frac{10}{6\sqrt{5}}$
- (b)  $\frac{4}{5\sqrt{2}}$
- (c)  $\frac{2\sqrt{3}}{5}$
- (d)  $\frac{\sqrt{2}}{10}$

Answer:

- (d)  $\frac{\sqrt{2}}{10}$

Question 27.

The reflection of the point  $(\alpha, \beta, \gamma)$  in the xy-plane is

- (a)  $(\alpha, \beta, 0)$
- (b)  $(0, 0, \gamma)$
- (c)  $(-\alpha, -\beta, -\gamma)$
- (d)  $(\alpha, \beta, -\gamma)$

Answer:

- (d)  $(\alpha, \beta, -\gamma)$

Question 28.

The area of the quadrilateral ABCD, where A(0, 4, 1), B(2, 3, -1), C(4, 5, 0) and D(2, 6, 2), is equal to

- (a) 9 sq. units
- (b) 18 sq. units
- (c) 27 sq. units
- (d) 81 sq. units

Answer:

- (a) 9 sq. units

Question 29.

The locus represented by  $xy + yz = 0$  is

- (a) A pair of perpendicular lines
- (b) A pair of parallel lines
- (c) A pair of parallel planes
- (d) A pair of perpendicular planes

Answer:

- (d) A pair of perpendicular planes

Question 30.

Direction cosines of the line that makes equal angles with the three axes in space are

- (a)  $\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{3}, \pm \frac{1}{3}$
- (b)  $\pm \frac{6}{7}, \pm \frac{2}{7}, \pm \frac{3}{7}$
- (c)  $\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}$
- (d)  $\sqrt{\frac{1}{7}}, \pm \sqrt{\frac{3}{14}}, \frac{1}{\sqrt{14}}$

Answer:

- (c)  $\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}$

Question 31.

If the direction ratios of a line are 1, -3, 2, then its direction cosines are

- (a)  $\frac{1}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{2}{\sqrt{14}}$
- (b)  $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$
- (c)  $\frac{-1}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{2}{\sqrt{14}}$
- (d)  $\frac{-1}{\sqrt{14}}, \frac{-2}{\sqrt{14}}, \frac{-3}{\sqrt{14}}$

Answer:

- (a)  $\frac{1}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{2}{\sqrt{14}}$

Question 32.

The cosines of the angle between any two diagonals of a cube is

- (a)  $\frac{1}{3}$

- (b)  $\frac{1}{2}$
- (c)  $\frac{2}{3}$
- (d)  $\frac{1}{\sqrt{3}}$

Answer:

- (a)  $\frac{1}{3}$

Question 33.

Which of the following is false?

- (a)  $30^\circ, 45^\circ, 60^\circ$  can be the direction angles of a line in space.
- (b)  $90^\circ, 135^\circ, 45^\circ$  can be the direction angles of a line in space.
- (c)  $120^\circ, 60^\circ, 45^\circ$  can be the direction angles of a line in space.
- (d)  $60^\circ, 45^\circ, 60^\circ$  can be the direction angles of a line in space.

Answer:

- (a)  $30^\circ, 45^\circ, 60^\circ$  can be the direction angles of a line in space.

Question 34.

A line makes angles  $\alpha, \beta$  and  $\gamma$  with the co-ordinate axes. If  $\alpha + \beta = 90^\circ$ , then  $\gamma$  is equal to

- (a)  $0^\circ$
- (b)  $90^\circ$
- (c)  $180^\circ$
- (d) None of these

Answer:

- (b)  $90^\circ$

Question 35.

If a line makes an angle  $\theta_1, \theta_2, \theta_3$  with the axis respectively, then  $\cos 2\theta_1 + \cos 2\theta_2 + \cos 2\theta_3 =$

- (a) -4
- (b) -2
- (c) -3
- (d) -1

Answer:

- (d) -1

Question 36.

The coordinates of a point P are (3, 12, 4) w.r.t. origin O, then the direction cosines of OP are

- (a) 3, 12, 4
- (b)  $\frac{1}{4}, \frac{1}{3}, \frac{1}{2}$
- (c)  $\frac{3}{\sqrt{13}}, \frac{12}{\sqrt{13}}, \frac{4}{\sqrt{13}}$
- (d)  $\frac{3}{13}, \frac{12}{13}, \frac{4}{13}$

Answer:

(d)  $\frac{3}{13}, \frac{12}{13}, \frac{4}{13}$

Question 37.

Find the direction cosines of the line joining A(0, 7, 10) and B(-1, 6, 6).

(a)  $\frac{-1}{3\sqrt{2}}, \frac{-1}{3\sqrt{2}}, \frac{2}{3\sqrt{2}}$       (b)  $\frac{1}{3\sqrt{2}}, \frac{1}{3\sqrt{2}}, \frac{4}{3\sqrt{2}}$

(c)  $\frac{1}{3}, \frac{-1}{3}, \frac{4}{3}$       (d) None of these

Answer:

(b)  $\frac{1}{3\sqrt{2}}, \frac{1}{3\sqrt{2}}, \frac{4}{3\sqrt{2}}$

Question 38.

The direction cosines of a line passing through two points P( $x_1, y_1, z_1$ ) and Q( $x_2, y_2, z_2$ ) are

(a)  $(x_2 - x_1), (y_2 - y_1), (z_2 - z_1)$

(b)  $(x_2 + x_1), (y_2 + y_1), (z_2 + z_1)$

(c)  $\frac{x_2 - x_1}{PQ}, \frac{y_2 - y_1}{PQ}, \frac{z_2 - z_1}{PQ}$

(d)  $\frac{x_2 + x_1}{PQ}, \frac{y_2 + y_1}{PQ}, \frac{z_2 + z_1}{PQ}$

Answer:

(c)  $\frac{x_2 - x_1}{PQ}, \frac{y_2 - y_1}{PQ}, \frac{z_2 - z_1}{PQ}$

Question 39.

The equation of a line which passes through the point (1, 2, 3) and is parallel to the vector

$3\hat{i} + 2\hat{j} - 2\hat{k}$ , is

(a)  $\vec{r} = (3\hat{i} + 2\hat{j} - 2\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$

(b)  $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k})$

(c)  $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} - 5\hat{k})$

(d)  $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(4\hat{i} + 4\hat{j} + \hat{k})$

Answer:

(b)  $r = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k})$

Question 40.

The equation of line passing through the point  $(-3, 2, -4)$  and equally inclined to the axes are

- (a)  $x - 3 = y + 2 = z - 4$
- (b)  $x + 3 = y - 2 = z + 4$
- (c)  $\frac{x+3}{1} = \frac{y-2}{2} = \frac{z+4}{3}$
- (d) None of these

Answer:

- (b)  $x + 3 = y - 2 = z + 4$

Question 41.

If  $l, m$  and  $n$  are the direction cosines of line  $l$ , then the equation of the line  $(l)$  passing through  $(x_1, y_1, z_1)$  is


- (a)  $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$
- (b)  $\left(\frac{x-x_1}{l}\right)\left(\frac{y-y_1}{m}\right) = \left(\frac{z-z_1}{n}\right)$
- (c)  $\frac{x+x_1}{l} = \frac{y+y_1}{m} = \frac{z+z_1}{n}$
- (d) None of these

Answer:

- (a)  $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$

Question 42.

In the figure,  $\vec{a}$  be the position vector of the point  $A$  with respect to the origin  $O$ .  $l$  is a line parallel to  $\vec{a}$  and  $\vec{b}$ . The vector equation of line  $l$  is



- (a)  $\vec{r} = \lambda(\vec{a} \times \vec{b})$
- (b)  $\vec{r} = \lambda\vec{a} - \vec{b}$
- (c)  $\vec{r} = \vec{a} + \lambda\vec{b}$
- (d)  $\vec{r} = \lambda(\vec{a} \cdot \vec{b})$

Answer:

(c)  $r = a + \lambda b$

Question 43.

The cartesian equation of the line  $l$  when it passes through the point  $(x_1, y_1, z_1)$  and parallel to the vector

$$b = a\hat{i} + b\hat{j} + c\hat{k}, \text{ is}$$

(a)  $x - x_1 = y - y_1 = z - z_1$

(b)  $x + x_1 = y + y_1 = z + z_1$

(c)  $\frac{x+x_1}{a} = \frac{y+y_1}{b} = \frac{z+z_1}{c}$

(d)  $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$

Answer:

(d)  $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$

Question 44.

The equation of the straight line passing through the point  $(a, b, c)$  and parallel to Z-axis is

(a)  $\frac{x-a}{1} = \frac{y-b}{1} = \frac{z-c}{0}$

(b)  $\frac{x-a}{0} = \frac{y-b}{1} = \frac{z-c}{1}$

(c)  $\frac{x-a}{1} = \frac{y-b}{0} = \frac{z-c}{0}$

(d)  $\frac{x-a}{0} = \frac{y-b}{0} = \frac{z-c}{1}$

Answer:

(d)  $\frac{x-a}{0} = \frac{y-b}{0} = \frac{z-c}{1}$

Question 45.

The coordinates of a point on the line  $\frac{x+2}{3} = \frac{y+1}{2} = \frac{z-3}{2}$  at a distance of  $\frac{6}{\sqrt{12}}$  from the point  $(1, 2, 3)$  is

(a)  $(56, 43, 111)$

(b)  $\left(\frac{56}{17}, \frac{43}{17}, \frac{111}{17}\right)$

(c)  $(2, 1, 3)$

(d)  $(-2, -1, -3)$

Answer:

(b)  $\left(\frac{56}{17}, \frac{43}{17}, \frac{111}{17}\right)$

Question 46.

Find the coordinates of the point where the line through the points (5, 1, 6) and (3, 4, 1) crosses the yz-plane.

(a)  $\left(0, -\frac{17}{2}, \frac{13}{2}\right)$

(b)  $\left(0, \frac{17}{2}, -\frac{13}{2}\right)$

(c)  $\left(10, \frac{19}{2}, \frac{13}{2}\right)$

(d) (0, 17, 13)

Answer:

(b)  $\left(0, \frac{17}{2}, -\frac{13}{2}\right)$

Question 47.

The point A(1, 2, 3), B(-1, -2, -1) and C(2, 3, 2) are three vertices of a parallelogram ABCD. Find the equation of CD.

(a)  $\frac{x}{1} = \frac{y}{2} = \frac{z}{2}$

(b)  $\frac{x+2}{1} = \frac{y+3}{2} = \frac{z-2}{2}$

(c)  $\frac{x}{2} = \frac{y}{3} = \frac{z}{2}$

(d)  $\frac{x-2}{1} = \frac{y-3}{2} = \frac{z-2}{2}$

Answer:

(d)  $\frac{x-2}{1} = \frac{y-3}{2} = \frac{z-2}{2}$

Question 48.

The equation of the line joining the points (-3, 4, 11) and (1, -2, 7) is

(a)  $\frac{x+3}{2} = \frac{y-4}{3} = \frac{z-11}{4}$

(b)  $\frac{x+3}{-2} = \frac{y-4}{3} = \frac{z-11}{2}$

(c)  $\frac{x+3}{-2} = \frac{y+4}{3} = \frac{z+11}{4}$

(d)  $\frac{x+3}{2} = \frac{y+4}{-3} = \frac{z+11}{2}$

Answer:

(b)  $\frac{x+3}{-2} = \frac{y-4}{3} = \frac{z-11}{2}$

Question 49.

The vector equation of the line through the points A(3, 4, -7) and B(1, -1, 6) is

- (a)  $\vec{r} = (3\hat{i} - 4\hat{j} - 7\hat{k}) + \lambda(\hat{i} - \hat{j} + 6\hat{k})$   
 (b)  $\vec{r} = (\hat{i} - \hat{j} + 6\hat{k}) + \lambda(3\hat{i} - 4\hat{j} - 7\hat{k})$   
 (c)  $\vec{r} = (3\hat{i} + 4\hat{j} - 7\hat{k}) + \lambda(-2\hat{i} - 5\hat{j} + 13\hat{k})$   
 (d)  $\vec{r} = (\hat{i} - \hat{j} + 6\hat{k}) + \lambda(4\hat{i} + 3\hat{j} - \hat{k})$

Answer:

(c)  $r = (3\hat{i} + 4\hat{j} - 7\hat{k}) + \lambda(-2\hat{i} - 5\hat{j} + 13\hat{k})$

Question 50.

The angle between the lines  $\vec{r} = (4\hat{i} - \hat{j}) + s(2\hat{i} + \hat{j} - 3\hat{k})$

and  $\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + t(\hat{i} - 3\hat{j} + 2\hat{k})$  is

- (a)  $\frac{3\pi}{2}$     (b)  $\frac{\pi}{3}$     (c)  $\frac{2\pi}{3}$     (d)  $\frac{\pi}{6}$

Answer:

(d)  $\frac{\pi}{6}$

Question 51.

The angle between the line  $2x = 3y = -z$  and  $6x = -y = -4z$  is

- (a)  $30^\circ$   
 (b)  $45^\circ$   
 (c)  $90^\circ$   
 (d)  $0^\circ$

Answer:

(c)  $90^\circ$

Question 52.

The angle between the lines  $3x = 6y = 2z$  and  $\frac{x-2}{-5} = \frac{y-1}{7} = \frac{z-3}{1}$  is

- (a)  $\frac{\pi}{6}$   
 (b)  $\frac{\pi}{4}$   
 (c)  $\frac{\pi}{3}$   
 (d)  $\frac{\pi}{2}$

Answer:

(d)  $\frac{\pi}{2}$

Question 53.

Find the angle between the pair of lines given by

$$\vec{r} = 3\hat{i} + 2\hat{j} - 4\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k}) \text{ and}$$

$$\vec{r} = 5\hat{i} - 2\hat{j} + \mu(3\hat{i} + 2\hat{j} + 6\hat{k}).$$

(a)  $\cos^{-1}\left(\frac{19}{21}\right)$

(b)  $\cos^{-1}\left(\frac{23}{19}\right)$

(c)  $\cos^{-1}\left(\frac{17}{13}\right)$

(d)  $\cos^{-1}\left(\frac{13}{9}\right)$

Answer:

(a)  $\cos^{-1}\left(\frac{19}{21}\right)$

Question 54.

The angle between the lines  $x = 1, y = 2$  and  $y = -1, z = 0$  is

(a)  $90^\circ$

(b)  $30^\circ$

(c)  $60^\circ$

(d)  $0^\circ$

Answer:

(a)  $90^\circ$

Question 55.

**Shortest distance between the two lines**

$$\frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7} \text{ and } \frac{x-2}{1} = \frac{y-4}{3} = \frac{z-6}{5} \text{ is}$$

(a) 0      (b)  $\frac{1}{2\sqrt{3}}$       (c) 3      (d) None of these

Answer:

(a) 0

Question 56.

The distance between line  $\vec{r} = \vec{a}_1 + t\vec{b}$  and  $\vec{r} = \vec{a}_2 + s\vec{b}$  is

- (a)  $|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|$       (b)  $\frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{b}|}$   
 (c)  $\frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{a}_2 - \vec{a}_1|}$       (d)  $\frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{a}_2 - \vec{a}_1| |\vec{b}|}$

Answer:

(b)  $\frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{b}|}$

Question 57.

The distance between the lines given by

$\vec{r} = \hat{i} + \hat{j} + \lambda(\hat{i} - 2\hat{j} + 3\hat{k})$  and

$\vec{r} = (2\hat{i} - 3\hat{k}) + \mu(\hat{i} - 2\hat{j} + 3\hat{k})$  is

- (a)  $\sqrt{\frac{59}{14}}$     (b)  $\sqrt{\frac{59}{7}}$     (c)  $\sqrt{\frac{118}{7}}$     (d)  $\frac{\sqrt{59}}{7}$

Answer:

(b)  $\sqrt{\frac{59}{7}}$

Question 58.

The distance between the lines  $\frac{x-4}{5} = \frac{y+1}{2} = \frac{z}{1}$  and

$\frac{x-1}{5} = \frac{y-2}{2} = \frac{z-3}{1}$  is

- (a)  $5\sqrt{129}$     (b)  $\frac{\sqrt{129}}{5}$     (c)  $\sqrt{\frac{129}{10}}$     (d)  $\sqrt{\frac{129}{5}}$

Answer:

(d)  $\sqrt{\frac{129}{5}}$

Question 59.

The direction cosines of the unit vector perpendicular to the plane  $r \cdot (6\hat{i} - 3\hat{j} - 2\hat{k}) + 1 = 0$  passing through the origin are

- (a)  $\frac{6}{7}, \frac{3}{7}, \frac{2}{7}$
- (b) 6, 3, 2
- (c)  $-\frac{6}{7}, \frac{3}{7}, \frac{2}{7}$
- (d) -6, 3, 2

Answer:

- (c)  $-\frac{6}{7}, \frac{3}{7}, \frac{2}{7}$

Question 60.

The coordinate of the foot of perpendicular drawn from origin to the plane  $2x - 3y + 4z - 6 = 0$  is

- (a)  $\left(\frac{2}{\sqrt{29}}, \frac{-3}{\sqrt{29}}, \frac{4}{\sqrt{29}}\right)$
- (b)  $\left(\frac{12}{29}, \frac{-18}{29}, \frac{24}{29}\right)$
- (c) (12, -18, 24)
- (d)  $\left(\frac{12}{\sqrt{29}}, \frac{-18}{\sqrt{29}}, \frac{24}{\sqrt{29}}\right)$

Answer:

- (d)  $\left(\frac{12}{\sqrt{29}}, \frac{-18}{\sqrt{29}}, \frac{24}{\sqrt{29}}\right)$

Question 61.

The vector equation of a plane which is at a distance of 7 units from the origin and normal to the vector

$3\hat{i} + 5\hat{j} - 6\hat{k}$  is

- (a)  $\vec{r} \cdot (3\hat{i} + 5\hat{j} - 6\hat{k}) = 7$
- (b)  $\vec{r} \cdot (3\hat{i} + 5\hat{j} - 6\hat{k}) = \frac{7}{\sqrt{70}}$
- (c)  $\vec{r} \cdot \left(\frac{3}{70}\hat{i} + \frac{5}{70}\hat{j} - \frac{6}{70}\hat{k}\right) = 7\sqrt{70}$
- (d)  $\vec{r} \cdot \left(\frac{3\hat{i}}{70} + \frac{5\hat{j}}{70} - \frac{6\hat{k}}{70}\right) = 7$

Answer:

- (d)  $r \cdot \left(\frac{3\hat{i}}{70} + \frac{5\hat{j}}{70} - \frac{6\hat{k}}{70}\right) = 7$

Question 62.

Find the vector equation of the plane which is at a distance of 8 units from the origin and which is

normal to the vector  $2\hat{i} + \hat{j} + 2\hat{k}$ .

(a)  $\vec{r} \cdot (2\hat{i} + \hat{j} + 2\hat{k}) = 0$  (b)  $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 24$

(c)  $\vec{r} \cdot (2\hat{i} + \hat{j} + 2\hat{k}) = 24$  (d) None of these

Answer:

(c)  $r \cdot (2\hat{i} + \hat{j} + 2\hat{k}) = 24$

Question 63.

Find the length of perpendicular from the origin to the plane  $r(3\hat{i} - 4\hat{j} + 12\hat{k})$ .

(a)  $\frac{5}{13}$

(b)  $\frac{5}{\sqrt{13}}$

(c)  $\frac{5}{23}$

(d)  $\frac{\sqrt{5}}{13}$

Answer:

(a)  $\frac{5}{13}$

Question 64.

The equation of the plane passing through three non-collinear points with position vectors a, b, c is

(a)  $r \cdot (b \times c + c \times a + a \times b) = 0$

(b)  $r \cdot (b \times c + c \times a + a \times b) = [abc]$

(c)  $r \cdot (a \times (b + c)) = [abc]$

(d)  $r \cdot (a + b + c) = 0$

Answer:

(b)  $r \cdot (b \times c + c \times a + a \times b) = [abc]$