

9. Quadrilaterals and Parallelograms

Exercise 9A

1. Question

Three angles of a quadrilateral measure 56° , 115° and 84° . Find the measure of the fourth angle.

Answer

Let the measure of the fourth angle be x° .

Since the sum of the angles of a quadrilateral is 360° , we have:

$$\therefore 56^\circ + 115^\circ + 84^\circ + x^\circ = 360^\circ$$

$$\therefore 255^\circ + x^\circ = 360^\circ$$

$$\therefore x^\circ = 105^\circ$$

Hence, the measure of the fourth angle is 105° .

2. Question

The angles of a quadrilateral are in the ratio 2:4:5:7. Find the angles.

Answer

Our given ratio of angles is 2:4:5:7. Let common multiplying factor be x° .

Hence, $\angle A = 2x^\circ$, $\angle B = 4x^\circ$, $\angle C = 5x^\circ$ and $\angle D = 7x^\circ$

Since the sum of the angles of a quadrilateral is 360° , we have:

$$\therefore 2x + 4x + 5x + 7x = 360^\circ$$

$$\therefore 18x = 360^\circ$$


$$\therefore x = 20^\circ$$

$$\therefore \angle A = 40^\circ; \angle B = 80^\circ; \angle C = 100^\circ; \angle D = 140^\circ$$

Hence, the measure of the angles are 40° , 80° , 100° and 140°

3. Question

In the adjoining figure, $ABCD$ is a trapezium in which $AB \parallel DC$. If $\angle A = 55^\circ$ and $\angle B = 70^\circ$, find $\angle C$ and $\angle D$.



Answer

Here given that ABCD is trapezium where $AB \parallel DC$.

We observe that $\angle A$ and $\angle D$ are the interior angles on the same side of transversal line AD, whereas $\angle B$ and $\angle C$ are the interior angles on the same side of transversal line BC.

As $\angle A$ and $\angle D$ are interior angles, we have,

$$\angle A + \angle D = 180^\circ$$

$$\therefore \angle D = 180^\circ - \angle A$$

$$\therefore \angle D = 180^\circ - 55^\circ = 125^\circ$$

Similarly for $\angle B$ and $\angle C$,

$$\angle B + \angle C = 180^\circ$$

$$\therefore \angle C = 180^\circ - \angle B$$


$$\therefore \angle C = 180^\circ - 70^\circ = 110^\circ$$

Hence, measure of $\angle D$ and $\angle C$ are 125° and 110° respectively.

4. Question

In the adjoining figure, ABCD is a square and $\triangle EDC$ is an equilateral triangle. Prove that

(i) $AE = BE$ (ii) $\angle DAE = 15^\circ$



Answer

(i) Here it is given that in ABCD is a square and $\triangle EDC$ is an equilateral triangle.

Hence, we say that $AB = BC = CD = DA$ and $ED = EC = DC$

Now in $\triangle ADE$ and $\triangle BCE$, we have,

$AD = BC$... given

$DE = EC \dots$ given

$\angle ADE = \angle BCE \dots$

as both angles are sum of 60° and 90°

$\therefore \triangle ADE \cong \triangle BCE$

Now by *cpct*,

$AE = BE \dots(1)$

(ii) Here $\angle ADE = 90^\circ + 60^\circ = 150^\circ$

$DA = DC \dots$ given

$DC = DE \dots$ given

$\therefore DA = DE$


This means that sides of square and triangles are equal.

$\therefore \triangle ADE$ and $\triangle BCE$ are isosceles triangles.

Hence, $\angle DAE = \angle DEA = \frac{1}{2}(180^\circ - 150^\circ) = 30^\circ/2 = 15^\circ$

5. Question

In the adjoining figure, $BM \perp AC$ and $DN \perp AC$. If $BM = DN$, prove that AC bisects BD .



Answer

Given: In $ABCD$, in which $BM \perp AC$ and $DN \perp AC$ and $BM = DN$.

To prove: AC bisects BD ie. $DO = BO$

Proof:

Now, in $\triangle OND$ and $\triangle OMB$, we have,

$\angle OND = \angle OMB \dots 90^\circ$ each

$\angle DON = \angle BOM \dots$ Vertically opposite angles

Also, $DN = BM \dots$ Given Hence, by AAS congruence rule,

$\triangle OND \cong \triangle OMB$

$\therefore OD = OB \dots$ CPCT

Hence, AC bisects BD .


6. Question

In the given figure, $ABCD$ is a quadrilateral in which $AB = AD$ and $BC = DC$. Prove that

(i) AC bisects $\angle A$ and $\angle C$,

(ii) $BE = DE$,

(iii) $\angle ABC = \angle ADC$.



Answer

Given: In $ABCD$, $AB = AD$ and $BC = DC$.

To prove: (i) AC bisects $\angle A$ and $\angle C$,

(ii) $BE = DE$,

(iii) $\angle ABC = \angle ADC$.

Proof:

(i) In $\triangle ABC$ and $\triangle ADC$, we have,

$AB = AD$...given

$BC = DC$...given

$AC = AC$... common side

Hence, by SSS congruence rule,

$\triangle ABC \cong \triangle ADC$

$\therefore \angle BAC = \angle DAC$ and $\angle BCA = \angle DCA$...By cpct

Thus, AC bisects $\angle A$ and $\angle C$.

(ii) Now, in $\triangle ABE$ and $\triangle ADE$, we have,

$AB = AD$...given

$\angle BAE = \angle DAE$...from i

$AE = AE$...common side

Hence, by SAS congruence rule,

$\triangle ABE \cong \triangle ADE$

$\therefore BE = DE$...by cpct

(iii) $\triangle ABC \cong \triangle ADC$ from ii


$\therefore \angle ABC = \angle ADC$...by cpct

7. Question

In the given figure, ABCD is a square and $\angle PQR = 90^\circ$. If $PB = QC = DR$, prove that

(i) $QB = RC$, (ii) $PQ = QR$,

(iii) $\angle QPR = 45^\circ$.



Answer

Given: ABCD is where $\angle PQR = 90^\circ$ and $PB = QC = DR$,

To prove: (i) $QB = RC$, (ii) $PQ = QR$,

(iii) $\angle QPR = 45^\circ$.

Proof:

(i) Here,

$BC = CD$...Sides of square

$CQ = DR$...Given

$BC = BQ + CQ$

$\therefore CQ = BC - BQ$

$\therefore DR = BC - BQ$...(1)

Also,

$CD = RC + DR$

$\therefore DR = CD - RC = BC - RC$...(2)

From (1) and (2), we have,

$BC - BQ = BC - RC$

$\therefore BQ = RC$

(ii) Now in $\triangle RCQ$ and $\triangle QBP$, we have,

$PB = QC$...Given

$BQ = RC$...from (i)

$\angle RCQ = \angle QBP$... 90° each

Hence by SAS congruence rule,

$\triangle RCQ \cong \triangle QBP$

$\therefore QR = PQ$...by cpct

(iii) $\triangle RCQ \cong \triangle QBP$ and $QR = PQ$... from (ii)

$\therefore$ In $\triangle RPQ$,

$$\angle QPR = \angle QRP = \frac{1}{2} (180^\circ - 90^\circ) = \frac{90^\circ}{2} = 45^\circ$$

$\therefore \angle QPR = 45^\circ$

8. Question


If O is a point within a quadrilateral $ABCD$, show that $OA + OB + OC + OD > AC + BD$.

Answer

Given: In $ABCD$, O is any point within the quadrilateral.

To prove: $OA + OB + OC + OD > AC + BD$.

Proof:



We know that the sum of any two sides of a triangle is greater than the third side. So, in $\triangle AOC$,

$$OA + OC > AC \dots (1)$$

Also, in $\triangle BOD$,

$$OB + OD > BD \dots (2)$$

Adding 1 and 2, we get,

$$(OA + OC) + (OB + OD) > (AC + BD)$$

$$\therefore OA + OB + OC + OD > AC + BD$$

Hence proved.


9. Question

In the adjoining figure, $ABCD$ is a quadrilateral and AC is one of its diagonals. Prove that:

(i) $AB + BC + CD + DA > 2AC$

(ii) $AB + BC + CD > DA$

(iii) $AB + BC + CD + DA > AC + BD$



Answer

Given: In $ABCD$, AC is one of diagonals.

To prove:

(i) $AB + BC + CD + DA > 2AC$

(ii) $AB + BC + CD > DA$

(iii) $AB + BC + CD + DA > AC + BD$

Proof:

(i) We know that the sum of any two sides of a triangle is greater than the third side. In $\triangle ABC$,

$$AB + BC > AC \dots(1)$$

In $\triangle ACD$,

$$CD + DA > AC \dots(2)$$

Adding (1) and (2), we get,

$$AB + BC + CD + DA > 2AC$$

(ii) In $\triangle ABC$, we have,

$$AB + BC > AC \dots(1)$$

We also know that the length of each side of a triangle is greater than the positive difference of the length of the other two sides.

In $\triangle ACD$, we have:

$$AC > DA - CD \dots(2)$$

From (1) and (2), we have,

$$AB + BC > DA - CD$$

$$\therefore AB + BC + CD > DA$$

(ii) In $\triangle ABC$,

$$AB + BC > AC \dots(1)$$

In $\triangle ACD$,

$$CD + DA > AC \dots(2)$$

In $\triangle BCD$,

$$BC + CD > BD \dots(3)$$

In $\triangle ABD$,

$$DA + AB > BD \dots(4)$$

Adding 1, 2, 3 and 4, we get,


$$2(AB + BC + CD + DA) > 2(AC + BD)$$

$$\therefore AB + BC + CD + DA > AC + BD$$

10. Question

Prove that the sum of all the angles of a quadrilateral is 360° .

Answer



Given: Consider a PQRS where QS is diagonal.

To prove: $\angle P + \angle Q + \angle R + \angle S = 360^\circ$

Proof:

For $\triangle PQS$, we have,

$$\angle P + \angle PQS + \angle PSQ = 180^\circ \dots (1) \dots \text{Using Angle sum property of Triangle}$$

Similarly, in $\triangle QRS$, we have,

$$\therefore \angle SQR + \angle R + \angle QSR = 180^\circ \dots (2) \dots \text{Using Angle sum property of Triangle}$$

On adding (1) and (2), we get

$$\angle P + \angle PQS + \angle PSQ + \angle SQR + \angle R + \angle QSR = 180^\circ + 180^\circ$$

$$\therefore \angle P + \angle PQS + \angle SQR + \angle R + \angle QSR + \angle PSQ = 360^\circ$$


$$\therefore \angle P + \angle Q + \angle R + \angle S = 360^\circ$$

$\therefore$ The sum of all the angles of a quadrilateral is 360° .

Exercise 9B

1. Question

In the adjoining figure, $ABCD$ is a parallelogram in which $\angle A = 72^\circ$. Calculate $\angle B$, $\angle C$ and $\angle D$.



Answer

In $ABCD$, $\angle A = 72^\circ$

We know that opposite angles of a parallelogram are equal.

Hence, $\angle A = \angle C$ and $\angle B = \angle D$

$$\therefore \angle C = 72^\circ$$

$\angle A$ and $\angle B$ are adjacent angles.

$$\therefore \angle A + \angle B = 180^\circ$$

$$\angle B = 180^\circ - \angle A$$


$$\angle B = 180^\circ - 72^\circ = 108^\circ$$

$$\therefore \angle B = \angle D = 108^\circ$$

Hence, $\angle B = \angle D = 108^\circ$ and $\angle C = 72^\circ$

2. Question

In the adjoining figure, $ABCD$ is a parallelogram in which $\angle DAB = 80^\circ$ and $\angle DBC = 60^\circ$. Calculate $\angle CDB$ and $\angle ADB$.



Answer

It is given that $ABCD$ is parallelogram and $\angle DAB = 80^\circ$ and $\angle DBC = 60^\circ$

We need to find measure of $\angle CDB$ and $\angle ADB$

In $ABCD$, $AD \parallel BC$, BD as transversal,

$$\angle DBC = \angle ADB = 60^\circ \text{ ...Alternate interior angles ... (i)}$$

As $\angle DAB$ and $\angle ADC$ are adjacent angles,

$$\angle DAB + \angle ADC = 180^\circ$$

$$\therefore \angle ADC = 180^\circ - \angle DAB$$

$$\angle ADC = 180^\circ - 80^\circ = 100^\circ$$

Also,

$$\angle ADC = \angle ADB + \angle CDB$$

$$\therefore \angle ADC = 100^\circ$$

$$\angle ADB + \angle CDB = 100^\circ \text{ ... (ii)}$$

From (i) and (ii), we get:

$$60^\circ + \angle CDB = 100^\circ$$


$$\Rightarrow \angle CDB = 100^\circ - 60^\circ = 40^\circ$$

Hence, $\angle CDB = 40^\circ$ and $\angle ADB = 60^\circ$

3. Question

In the adjoining figure, $ABCD$ is a parallelogram in which $\angle A = 60^\circ$. If the bisectors of $\angle A$ and $\angle B$ meet DC at P , prove that

(i) $\angle APB = 90^\circ$, (ii) $AD = DP$ and $PB = PC = BC$, (iii) $DC = 2AD$.



Answer

Given: $ABCD$ is a parallelogram. The bisectors of $\angle A$ and $\angle B$ meet DC at P .

To prove: (i) $\angle APB = 90^\circ$, (ii) $AD = DP$ and $PB = PC = BC$, (iii) $DC = 2AD$.

Proof:

$\therefore \angle A = \angle C$ and $\angle B = \angle D$... Opposite angles

And $\angle A + \angle B = 180^\circ$... Adjacent angles

$$\therefore \angle B = 180^\circ - \angle A$$

$$180^\circ - 60^\circ = 120^\circ \text{ ... as } \angle A = 60^\circ$$

$$\therefore \angle A = \angle C = 60^\circ \text{ and } \angle B = \angle D = 120^\circ$$

(i) In $\triangle APB$,

$$\angle PAB = \frac{60^\circ}{2} = 30^\circ \text{ and } \angle PBA = \frac{120^\circ}{2} = 60^\circ$$

$$\therefore \angle APB = 180^\circ - (30^\circ + 60^\circ) = 90^\circ$$

(ii) In $\triangle ADP$, $\angle PAD = 30^\circ$ and $\angle ADP = 120^\circ$

$$\therefore \angle APB = 180^\circ - (30^\circ + 120^\circ) = 30^\circ$$

Hence,

$$\angle PAD = \angle APB = 30^\circ$$

Hence, $\triangle ADP$ is an isosceles triangle and $AD = DP$. In $\triangle PBC$,

$$\angle PBC = 60^\circ$$

$$\angle BPC = 180^\circ - (90^\circ + 30^\circ) = 60^\circ \text{ and } \angle BCP = 60^\circ \dots \text{Opposite angle of } \angle A$$

$$\therefore \angle PBC = \angle BPC = \angle BCP$$

Hence, $\triangle PBC$ is an equilateral triangle and, therefore, $PB = PC = BC$.

$$(iii) DC = DP + PC$$

From (ii), we have


$$DC = AD + BC \dots AD = BC \quad DC = AD + AD$$

$$DC = 2 AD$$

4. Question

In the adjoining figure, $ABCD$ is a parallelogram in which $\angle BAO = 35^\circ$, $\angle DAO = 40^\circ$ and $\angle COD = 105^\circ$.

Calculate (i) $\angle ABO$, (ii) $\angle ODC$, (iii) $\angle ODC$, (iv) $\angle CBD$.



Answer

In $ABCD$, $\angle BAO = 35^\circ$, $\angle DAO = 40^\circ$ and $\angle COD = 105^\circ$.

(i) In $\triangle AOB$,

$$\angle BAO = 35^\circ$$

$$\angle AOB = \angle COD = 105^\circ \dots \text{Vertically opposite angles}$$

$$\therefore \angle ABO = 180^\circ - (35^\circ + 105^\circ) = 40^\circ \dots \text{Using Angle sum property of Triangle}$$

(ii) $\angle ODC$ and $\angle ABO$ are alternate angles for transversal BD

$$\therefore \angle ODC = \angle ABO = 40^\circ$$

(iii) $\angle ACB = \angle CAD = 40^\circ$...Alternate angles for transversal AC

$$(iv) \angle CBD = \angle ABC - \angle ABD \dots (1)$$

$\angle ABC = 180^\circ - \angle BAD$...Adjacent angles are supplementary

$$\angle ABC = 180^\circ - 75^\circ = 105^\circ$$

$$\angle CBD = 105^\circ - \angle ABD \dots \text{as } \angle ABD = \angle ABO$$

$$\angle CBD = 105^\circ - 40^\circ = 65^\circ$$

5. Question

In a ||gm ABCD, if $\angle A = (2x + 25)^\circ$ and $\angle B = (3x - 5)^\circ$, find the value of x and the measure of each angle of the parallelogram.

Answer

It is given that in ABCD, $\angle A = (2x + 25)^\circ$ and $\angle B = (3x - 5)^\circ$,

We know that opposite angles of parallelogram are equal.

$$\therefore \angle A = \angle C \text{ and } \angle B = \angle D$$

Also,

$\angle A + \angle B = 180^\circ$...Adjacent angles of parallelogram are supplementary

$$\therefore (2x + 25)^\circ + (3x - 5)^\circ = 180^\circ$$

$$5x^\circ + 20^\circ = 180^\circ$$

$$5x^\circ = 160^\circ$$

$$x^\circ = 32^\circ$$

$$\therefore \angle A = 2 \times 32 + 25 = 89^\circ$$

$$\therefore \angle B = 3 \times 32 - 5 = 91^\circ$$

Hence, $x = 32^\circ$, $\angle A = \angle C = 89^\circ$ and $\angle B = \angle D = 91^\circ$

6. Question

If an angle of a parallelogram is four-fifths of its adjacent angle, find the angles of the parallelogram.

Answer

Let ABCD be the parallelogram.

We know that opposite angles of parallelogram are equal.

$$\therefore \angle A = \angle C \text{ and } \angle B = \angle D \text{ By given conditions,}$$

Let $\angle A = x^\circ$ and $\angle B = \frac{4x^\circ}{5}$

Also, adjacent angles of parallelogram are supplementary,

$$\therefore x^\circ + \frac{4x^\circ}{5} = 180^\circ$$

$$\frac{9x^\circ}{5} = 180^\circ$$

$$\therefore x = 100^\circ$$

Hence, $\angle A = 100^\circ$ and $\angle B = \frac{4 \times 100^\circ}{5} = 80^\circ$

Hence, $\angle A = \angle C = 100^\circ$; $\angle B = \angle D = 80^\circ$

7. Question

Find the measure of each angle of a parallelogram, if one of its angles is 30° less than twice the smallest angle.

Answer

Let ABCD be the parallelogram.

We know that opposite angles of parallelogram are equal.

$$\therefore \angle A = \angle C \text{ and } \angle B = \angle D$$

Let $\angle A$ be the smallest angle whose measure is x° .

$$\therefore \angle B = (2x - 30)^\circ$$

We know that adjacent angles of parallelogram are supplementary,

$$\angle A + \angle B = 180^\circ$$

$$x + 2x - 30^\circ = 180^\circ$$

$$3x = 210^\circ$$

$$x = 70^\circ$$

$$\therefore \angle B = 2 \times 70^\circ - 30^\circ = 110^\circ$$

Hence, $\angle A = \angle C = 70^\circ$ and $\angle B = \angle D = 110^\circ$

8. Question

ABCD is a parallelogram in which $AB = 9.5$ cm and its perimeter is 30 cm. Find the length of each side of the parallelogram.

Answer

Here ABCD is parallelogram.

We know that the opposite sides of a parallelogram are parallel and equal.

Hence, $AB = DC = 9.5 \text{ cm}$

Also let $BC = AD = x \text{ cm}$

Now,

Perimeter of ABCD = 30 cm ...(given)

$$\therefore AB + BC + CD + DA = 30 \text{ cm}$$

$$\therefore 9.5 + x + 9.5 + x = 30$$

$$\therefore 19 + 2x = 30$$


$$\therefore 2x = 11$$

$$\therefore x = 5.5 \text{ cm}$$

Hence, length of each side is $AB = DC = 9.5 \text{ cm}$ and $BC = DA = 5.5 \text{ cm}$

9. Question

In each of the figures given below, ABCD is a rhombus. Find the value of x and y in each case.



Answer

(i) ABCD is a rhombus.

We know that rhombus is type of parallelogram whose all sides are equal.

$$\text{In } \triangle ABC, \angle BAC = \angle BCA = \frac{1}{2} (180^\circ - 110^\circ) = 35^\circ$$

$$\text{Hence } x = 35^\circ$$

But $AB \parallel DC$...opposite sides of rhombus are parallel

$$\angle BAC = \angle DCA \text{ ...for transversal AC}$$

$$\therefore \angle BAC = \angle DCA = 35^\circ$$

$$\text{Hence, } x = y = 35^\circ$$

(ii) ABCD is a rhombus.

We know that the diagonals of a rhombus are perpendicular bisectors of each other.

$\therefore$ in $\triangle AOB$,

$$\angle OAB = 40^\circ, \angle AOB = 90^\circ$$

$$\therefore \angle ABO = 180^\circ - (40^\circ + 90^\circ) = 50^\circ$$

$$\text{Hence } x = 50^\circ$$

Now in $\triangle DAB$,

$AB = AD$... as rhombus has all sides equal.

ie. $\triangle AOB$ is *isosceles triangle*.

Also base angles of *isosceles triangle* are equal.

$$\text{Hence, } x = y = 50^\circ$$

(iii) ABCD is a rhombus.

We know that rhombus is type of parallelogram whose all sides are equal.

So in $\triangle DCB$,

$$DC = BC$$

$$\therefore \angle CDB = \angle CBD = y^\circ \text{ base angles of } \textit{isosceles triangle} \text{ are equal.}$$

Now, $x = \angle CAB$...alternate angles with transversal AC

$$\therefore x = \frac{1}{2} \angle BAD$$

$$\therefore x = \frac{1}{2} \times 62^\circ$$

$$\therefore x = 31^\circ$$

In $\triangle DOC$,

We know sum of angles of triangle is 180°

$$\angle CDO + \angle DOC + \angle OCD = 180^\circ$$

$$\therefore \angle CDO + 90^\circ + 31^\circ = 180^\circ$$

$$\therefore \angle CDO = 59^\circ$$

$$\therefore y = 59^\circ$$


$$\text{Hence, } x = 31^\circ \text{ and } y = 59^\circ$$

10. Question

The lengths of the diagonals of a rhombus are 24 cm and 18 cm respectively. Find the length of each side of the rhombus.

Answer

Let ABCD be rhombus.



Here, AC and BD are the diagonals of ABCD, where AC = 24 cm and BD = 18 cm.

Let the diagonals intersect each other at O.

We know that the diagonals of a rhombus are perpendicular bisectors of each other.

$\therefore \triangle AOB$ is a right angle triangle in which $OA = \frac{24}{2} = 12$ cm and $OB = \frac{18}{2} = 9$ cm.

Now, $AB^2 = OA^2 + OB^2$...Pythagoras theorem

$$\therefore AB^2 = (12)^2 + (9)^2$$

$$\therefore AB^2 = 144 + 81 = 225$$

$$\therefore AB = 15 \text{ cm}$$


Hence, the side of the rhombus is 15 cm

11. Question

Each side of a rhombus is 10 cm long and one of its diagonals measures 16 cm. Find the length of the other diagonal and hence find the area of the rhombus.

Answer

Let ABCD be rhombus.



We know that rhombus is type of parallelogram whose all sides are equal.

$$\therefore AB = BC = CD = DA = 10 \text{ cm}$$

Let the diagonals AC and BD intersect each other at O, where AC = 16 cm and let BD = x

We know that the diagonals of a rhombus are perpendicular bisectors of each other.

$\therefore \triangle AOB$ is a right angle triangle, in which $OB = BD \div 2 = x \div 2$ and $OA = AC \div 2 = 16 \div 2 = 8$ cm.

Now, $AB = OA^2 + OB^2$...by pythagoras theorem $\therefore 10^2 = \left(\frac{x}{2}\right)^2 + 8^2$

$$\text{ie. } 100 - 64 = \frac{x^2}{4}$$

$$36 \times 4 = x^2$$

$$\therefore x^2 = 144$$

$$\therefore x = 12 \text{ cm}$$

Hence, the length of the other diagonal is 12 cm

We know that area of rhombus is,

$$\text{Area of rhombus} = \frac{1}{2} \times (\text{Diagonal1}) \times (\text{Diagonal2})$$

Hence,

$$\text{Area of ABCD} = \frac{1}{2} \times AC \times BD$$


$$= \frac{1}{2} \times 16 \times 12$$

$$= 96 \text{ cm}^2$$

Hence, the area of rhombus is 96 cm^2

12. Question

In each of the figures given below, $ABCD$ is a rectangle. Find the values of x and y in each case.



Answer

(i) Here, ABCD is rectangle.

We know that the diagonals of a rectangle are congruent and bisect each other.

$\therefore$ In $\triangle AOB$, we have $OA = OB$

This means that $\triangle AOB$ is isosceles triangle.

We know that base angles of isosceles triangle are equal.

$$\therefore \angle OAB = \angle OBA = 35^\circ$$

$$\therefore \therefore x = 90^\circ - 35^\circ = 55^\circ$$

$$\text{Also, } \angle AOB = 180^\circ - (35^\circ + 35^\circ) = 110^\circ$$

$$\therefore y = \angle AOB = 110^\circ \dots \text{Vertically opposite angles}$$

$$\text{Hence, } x = 55^\circ \text{ and } y = 110^\circ$$

(ii) Here, ABCD is rectangle.

We know that the diagonals of a rectangle are congruent and bisect each other.

$$\therefore \text{In } \triangle AOB, \text{ we have } OA = OB$$

This means that $\triangle AOB$ is isosceles triangle.

We know that base angles of isosceles triangle are equal.

$$\therefore \angle OAB = \angle OBA = \frac{1}{2} \times (180^\circ - 110^\circ) = 35^\circ$$

$$\therefore y = \angle BAC = 35^\circ \dots \text{alternate angles with transversal AC}$$


$$\text{Also, } x = 90^\circ - y \dots \therefore \angle C = 90^\circ = x + y$$

$$\therefore x = 90^\circ - 35^\circ = 55^\circ$$

$$\text{Hence, } x = 55^\circ \text{ and } y = 35^\circ$$

13. Question

In the adjoining figures, ABCD is a square. A line segment CX cuts AB at X and the diagonal BD at O such that $\angle COD = 80^\circ$ and $\angle OXA = x^\circ$. Find the value of x .



Answer

Here, ABCD is square.

Here AC and BD are diagonals.

We know that the angles of a square are bisected by the diagonals.

$$\therefore \angle OBX = 45^\circ \because \angle ABC = 90^\circ \text{ and } BD \text{ bisects } \angle ABC$$

$$\text{And } \angle BOX = \angle COD = 80^\circ \dots \text{Vertically opposite angles}$$

$$\therefore \text{In } \triangle BOX, \text{ we have:}$$

$$\angle AXO = \angle OBX + \angle BOX \dots \text{Exterior angle theorem}$$


$$\Rightarrow \angle AXO = 45^\circ + 80^\circ = 125^\circ$$

$$\therefore x = 125^\circ$$

14. Question

In the adjoining figures, AL and CM are perpendiculars to the diagonal BD of a ||gm $ABCD$. Prove that

(i) $\triangle ALD \cong \triangle CMB$, (ii) $AL = CM$.



Answer

Here, $ABCD$ is parallelogram.

Hence, $AD \parallel BC$ and $AD = BC$

(i) In $\triangle ALD$ and $\triangle CMB$, we have, $AD = BC$


$$\angle ALD = \angle CMB \text{ (} 90^\circ \text{ each)}$$

$$\angle ADL = \angle CBM \text{ (Alternate interior angle)} \therefore \triangle ALD \cong \triangle CMB$$

(ii) As $\triangle ALD \cong \triangle CMB$...from 1. $\therefore AL = CM$...by cpct

15. Question

In the adjoining figures, $ABCD$ is a parallelogram in which the bisectors of $\angle A$ and $\angle B$ intersect at a point P . Prove that $\angle APB = 90^\circ$.



Answer

$ABCD$ is parallelogram.

We know that the sum of the adjacent angles in parallelogram is 180°

$$\therefore \angle A + \angle B = 180^\circ$$

$$\therefore \frac{\angle A}{2} + \frac{\angle B}{2} = \frac{180^\circ}{2} = 90^\circ$$

In $\triangle APB$, we have:

$$\angle PAB = \angle A / 2$$

$$\angle PBA = \angle B / 2$$

$\therefore \angle APB = 180 - (\angle PAB + \angle PBA)$...Angle sum property of triangle


$$\therefore \angle APB = 180 - \left(\frac{\angle A}{2} + \frac{\angle B}{2} \right)$$

$$\therefore \angle APB = 180 - 90 = 90^\circ$$

Hence, proved.

16. Question

In the adjoining figures, $ABCD$ is a parallelogram. If P and Q are points on AD and BC respectively such that $AP = \frac{1}{3}AD$ and $CQ = \frac{1}{3}BC$, prove that $AQCP$ is a parallelogram.



Answer

$ABCD$ is parallelogram

We know that opposite sides and angles of parallelogram *are equal*.

$$\therefore \angle B = \angle D \text{ and } AD = BC \text{ and } AB = DC$$

Also, $AD \parallel BC$ and $AB \parallel DC$

It is given that $AP = \frac{1}{3}AD$ and $CQ = \frac{1}{3}BC$,

Hence, $AP = CQ$... $\because AD = BC$

In $\triangle DPC$ and $\triangle BQA$, we have,

$$AB = CD$$

$$\angle B = \angle D$$

$$DP = QB \text{ ... as } AP = \frac{1}{3}AD \text{ and } CQ = \frac{1}{3}BC,$$

Hence, by SAS test for congruency,

$$\triangle DPC \cong \triangle BQA$$

$$\therefore PC = QA \text{ ... by cpct}$$


Hence, from above, in $AQCP$, we have,

$$AP = CQ \text{ and } PC = QA$$

$\therefore AQCP$ is a parallelogram.

17. Question

In the adjoining figures, $ABCD$ is a parallelogram whose diagonals intersect each other at O . A line segment EOF is drawn to meet AB at E and DC at F . Prove that $OE = OF$.



Answer

$ABCD$ is parallelogram.

$\therefore$ in $\triangle ODF$ and $\triangle OBE$, we have:

$OD = OB$... Diagonals bisect each other

$\angle DOF = \angle BOE$... Vertically opposite angles

$\angle FDO = \angle OBE$... Alternate interior angles

Hence, by SAA test for congruency,


$$\triangle ODF \cong \triangle OBE$$

$\therefore OF = OE$...by cpct

Hence, proved.

18. Question

In the adjoining figures, $ABCD$ is a parallelogram in which AB is produced to E so that $BE = AB$. Prove that ED bisects BC .



Answer

$ABCD$ is parallelogram.

In $\triangle ODC$ and $\triangle OEB$, we have,

$$DC = BE \text{ ...as } DC = AB$$

$\angle COD = \angle BOE$... Vertically opposite angles are equal

$\angle OCD = \angle OBE$... Alternate angles with transversal BC

Hence, by SAA test for congruency, we get,

$$\triangle ODC \cong \triangle OEB$$


$\therefore OC = OB$...by cpct

We know that $BC = OC + OB$.

$\therefore$ ED bisects BC.

19. Question

In the adjoining figures, $ABCD$ is a parallelogram and E is the midpoint of side BC . If DE and AB when produced meet at F , prove that $AF = 2AB$.



Answer

$ABCD$ is parallelogram.

Also given that $BE = CE$

In $ABCD$, $AB \parallel DC$

$\angle DCE = \angle EBF$... Alternate angles with transversal DF

In $\triangle DCE$ and $\triangle BFE$, we have,

$\angle DCE = \angle EBF$...from above

$\angle DEC = \angle BEF$... Vertically opposite angles

Also, $BE = CE$... givenHence, by ASA congruence rule,

$\triangle DCE \cong \triangle BFE$

$\therefore DC = BF$... by cpct

But $DC = AB$, as $ABCD$ is a parallelogram.

$\therefore DC = AB = BF$

Now, $AF = AB + BF$


From above, we get,

$AF = AB + AB = 2AB$

Hence, proved.

20. Question

A $\triangle ABC$ is given. If lines are drawn through A, B, C , parallel respectively to the sides BC, CA and AB , forming $\triangle PQR$, as shown in the adjoining figure, show that $BC = \frac{1}{2} QR$.



Answer

Here given that $BC \parallel QA$ and $CA \parallel QB$ which means that $BCQA$ is a parallelogram.

$$\therefore BC = QA \dots(1)$$

Similarly, $BC \parallel AR$ and $AB \parallel CR$, which means $BCRA$ is a parallelogram.

$$\therefore BC = AR \dots(2)$$

$$\text{But } QR = QA + AR$$

From (1) and (2), we get,


$$QR = BC + BC$$

$$\therefore QR = 2BC$$

$$\text{Hence, } BC = \frac{1}{2} QR$$

21. Question

In the adjoining figure, $\triangle ABC$ is a triangle and through A, B, C lines are drawn, parallel respectively to BC, CA and AB , intersecting at P, Q and R . Prove that the perimeter of $\triangle PQR$ is double the perimeter of $\triangle ABC$.



Answer

$$\text{Here, Perimeter of } \triangle ABC = AB + BC + CA$$

$$\text{And Perimeter of } \triangle PQR = PQ + QR + PR$$

Given that $BC \parallel QA$ and $CA \parallel QB$ which means $BCQA$ is a parallelogram.

$$\therefore BC = QA \dots(1)$$

Similarly, $BC \parallel AR$ and $AB \parallel CR$, which means $BCRA$ is a parallelogram.

$$\therefore BC = AR \dots(2)$$

$$\text{But, } QR = QA + AR$$

From 1 and 2,

$$QR = BC + BC$$

$$\therefore QR = 2BC$$

$$\therefore BC = \frac{1}{2} QR$$

$$\text{Similarly, } CA = \frac{1}{2} PQ \text{ and } AB = \frac{1}{2} PR$$

Now,

$$\text{Perimeter of } \triangle ABC = AB + BC + CA$$

$$= \frac{1}{2} QR + \frac{1}{2} PQ + \frac{1}{2} PR$$

$$= \frac{1}{2} (PR + QR + PQ)$$

This states that,


$$\text{Perimeter of } \triangle ABC = \frac{1}{2} (\text{Perimeter of } \triangle PQR)$$

$$\therefore \text{Perimeter of } \triangle PQR = 2 \times \text{Perimeter of } \triangle ABC$$


Exercise 9C

1. Question

In the adjoining figure, $ABCD$ is a trapezium in which $AB \parallel DC$ and E is the midpoint of AD . A line segment $EF \parallel AB$ meets BC at F . Show that F is the midpoint of BC .



Answer



Here, $ABCD$ is trapezium.

Join BD to cut EF at O .

It is given that, in $\triangle DAB$, E is the mid point of AD and $EO \parallel AB$.


$\therefore O$ is the midpoint of BD ...By converse of mid point theorem

Now in $\triangle BDC$, O is the mid point of BD and $OF \parallel DC$.

$\therefore F$ is the midpoint of BC ... By converse of mid point theorem

2. Question

In the adjoining figure, $ABCD$ is a $\parallel$ gm in which E and F are the midpoints of AB and CD respectively. If GH is a line segment that cuts AD , EF and BC at G , P and H respectively, prove that $GP = PH$.



Answer

Here, $ABCD$ is parallelogram.

By the properties of parallelogram,

$AD \parallel BC$ and $AB \parallel DC$

$AD = BC$ and $AB = DC$

Also,

$AB = AE + BE$ and $DC = DF + FC$

This means that,

$AE = BE = DF = FC$

Now, $DF = AE$ and $DF \parallel AE$, that is $AEFD$ is a parallelogram.

Hence, $AD \parallel EF$

Similarly, $BEFC$ is also a parallelogram.

Hence, $EF \parallel BC$

$\therefore AD \parallel EF \parallel BC$

Thus, AD , EF and BC are three parallel lines cut by the transversal line DC at D , F and C , respectively such that $DF = FC$.

Also, the lines AD , EF and BC are also cut by the transversal AB at A , E and B , respectively such that $AE = BE$.

Similarly, they are also cut by GH .


Hence by intercept theorem,

$\therefore GP = PH$

Hence proved.

3. Question

In the adjoining figure, $ABCD$ is a trapezium in which $AB \parallel DC$ and P, Q are the midpoints of AD and BC respectively. DQ and AB when produced meet at E . Also, AC and PQ intersect at R . Prove that (i) $DQ = QE$, (ii) $PR \parallel AB$, (iii) $AR = RC$.



Answer

Here, $ABCD$ is trapezium.

Hence, $AB \parallel DC$

Also given that $AP = PD$ and $BQ = CQ$

(i) In $\triangle QCD$ and $\triangle QBE$, we have,

$\angle DQC = \angle BQE$...Vertically opposite angles

$\angle DCQ = \angle EBQ$...Alternate angles with transversal BC

$BQ = CQ$... Q is the midpoint

Hence, by AAS test of congruency,

$\triangle QCD \cong \triangle QBE$

Hence, $DQ = QE$...by cpct

(ii) Also, in $\triangle ADE$, P and Q are the midpoints of AD and DE respectively

$\therefore PQ \parallel AE$

Hence, $PQ \parallel AB \parallel DC$

ie. $AB \parallel PR \parallel DC$

(iii) PQ , AB and DC are cut by transversal AD at P such that $AP = PD$.

Also they are cut by transversal BC at Q such that $BQ = QC$.


Similarly, lines PQ , AB and DC are also cut by AC at R .

Hence, by intercept theorem,

$\therefore AR = RC$

4. Question

In the adjoining figure, AD is a median of $\triangle ABC$ and $DE \parallel BA$. Show that BE is also a median of $\triangle ABC$.



Answer

In $\triangle ABC$, AD is median.

$$\therefore BD = DC$$

We know that the line drawn through the midpoint of one side of a triangle and parallel to another side bisects the third side.

So, in $\triangle ABC$, D is the mid point of BC and $DE \parallel BA$.

Hence, DE bisects AC .


$$\therefore AE = EC$$

This means that E is the midpoint of AC .

$\therefore BE$ is median of $\triangle ABC$.

5. Question

In the adjoining figure, AD and BE are the medians of $\triangle ABC$ and $DF \parallel BE$. Show that $CF = \frac{1}{4}AC$.



Answer

Here in $\triangle ABC$ AD and BE are medians.

Hence, in $\triangle ABC$, we have: $AC = AE + EC$

But $AE = EC$... as E is midpoint of AC

$$\therefore AC = 2EC \dots (1)$$

Now in $\triangle BEC$,

$DF \parallel BE$

Also, $EF = CF$... by midpoint theorem, as D is the midpoint of BC

But,

$$EC = EF + CF$$

$$\therefore EC = 2CF \dots (2)$$

From 1 and 2, we get,


$$AC = 4 CF$$

$$\therefore CF = \frac{1}{4} AC.$$

6. Question

In the adjoining figure, $ABCD$ is a parallelogram. E is the midpoint of DC and through D , a line segment is drawn parallel to EB to meet CB produced at G and it cuts AB at F . Prove that

$$(i) AD = \frac{1}{2} GC, (ii) DG = 2EB.$$



Answer

$ABCD$ is parallelogram.

(i) In $\triangle DCG$, we have:

$$DG \parallel EB$$

$$DE = EC \dots E \text{ is the midpoint of } DC$$

Also, $GB = BC \dots$ by midpoint theorem

$\therefore B$ is the midpoint of GC .

$$\text{Also, } GC = GB + BC$$

$$GC = 2BC$$

$$GC = 2 AD \dots \text{as } AD = BC$$

$$\therefore AD = \frac{1}{2} GC$$

(ii) Now, in $\triangle DCG$, $DG \parallel EB$ and E is the midpoint of DC and B is the midpoint of GC .

$$\therefore EB = \frac{1}{2} DG \dots \text{by midpoint theorem}$$


$$\therefore DG = 2 EB$$

7. Question

Prove that the line segments joining the middle points of the sides of a triangle divide it into four congruent triangles.

Answer

Let triangle be $\triangle ABC$. D , E and F are the midpoints of sides AB , BC and CA , respectively.



By midpoint theorem, for D and E as midpoints of sides AB and BC ,

$$DE \parallel AC$$

Similarly, $DF \parallel BC$ and $EF \parallel AB$.

$\therefore ADEF$, $BDFE$ and $DFCE$ are all parallelograms.

But, DE is the diagonal of the $BDFE$.

$$\therefore \triangle BDE \cong \triangle FED \dots (1)$$

Similarly, DF is the diagonal of the parallelogram $ADEF$.

$$\therefore \triangle DAF \cong \triangle FED \dots (2)$$


And, EF is the diagonal of the parallelogram $DFCE$.

$$\therefore \triangle EFC \cong \triangle FED \dots (3)$$

Hence, all the four triangles are congruent.

8. Question

In the adjoining figure, D , E , F are the midpoints of the sides BC , CA and AB respectively, of $\triangle ABC$. Show that $\angle EDF = \angle A$, $\angle DEF = \angle B$ and $\angle DFE = \angle C$.



Answer

Here, in $\triangle ABC$, D , E , F are the midpoints of the sides BC , CA and AB respectively.

By mid point theorem, as F and E are the mid points of sides AB and AC ,

$$FE \parallel BC$$

Similarly, $DE \parallel FB$ and $FD \parallel AC$.

Therefore, $AFDE$, $BDEF$ and $DCEF$ are all parallelograms.

We know that opposite angles in parallelogram are equal.

∴ In AFDE, we have,

$$\angle A = \angle EDF$$

In BDEF, we have,

$$\angle B = \angle DEF$$

In DCEF, we have,

$$\angle C = \angle DFE$$


Hence proved.

9. Question

Show that the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a rectangle is a rhombus.

Answer

Let ABCD be the rectangle and P, Q, R and S be the midpoints of AB, BC, CD and DA, respectively.



Join diagonals of the rectangle.

In $\triangle ABC$, we have, by midpoint theorem, ∴ $PQ \parallel AC$ and $PQ = \frac{1}{2} AC$

Similarly, $SR \parallel AC$ and $SR = \frac{1}{2} AC$.

As, $PQ \parallel AC$ and $SR \parallel AC$, then also $PQ \parallel SR$

Also, $PQ = SR$, each equal to $\frac{1}{2} AC \dots (1)$

So, PQRS is a parallelogram

Now, in $\triangle SAP$ and $\triangle QBP$, we have,

$$AS = BQ \quad \angle A = \angle B = 90^\circ \quad AP = BP$$

∴ By SAS test of congruency,

$$\triangle SAP \cong \triangle QBP$$

Hence, $PS = PQ \dots \text{by cpct} \dots (2)$

Similarly, $\triangle SDR \cong \triangle QCR$

$\therefore SR = RQ \dots$ by cpct ... (3)

Hence, from 1, 2 and 3 we have,


$PQ = PQ = SR = RQ$ Hence, PQRS is a rhombus.

Hence, the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a rectangle is a rhombus.

10. Question

Show that the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a rhombus is a rectangle.

Answer



In $\triangle ABC$, P and Q are mid points of AB and BC respectively.

$\therefore PQ \parallel AC$ and $PQ = \frac{1}{2}AC \dots (1) \dots$ Mid point theorem

Similarly in $\triangle ACD$, R and S are mid points of sides CD and AD respectively.

$\therefore SR \parallel AC$ and $SR = \frac{1}{2}AC \dots (2) \dots$ Mid point theorem

From (1) and (2), we get

$PQ \parallel SR$ and $PQ = SR$

Hence, PQRS is parallelogram (pair of opposite sides is parallel and equal)

Now, $RS \parallel AC$ and $QR \parallel BD$.

Also, $AC \perp BD \dots$ as diagonals of rhombus are perpendicular bisectors of each other.

$\therefore RS \perp QR$.


Thus, PQRS is a rectangle.

Hence, the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a rhombus is a rectangle.

11. Question

Show that the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a square is a square.

Answer



Let $ABCD$ be the square and P , Q , R and S be the midpoints of AB , BC , CD and DA , respectively.

Join diagonals of the square.

In $\triangle ABC$, we have, by midpoint theorem,

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC$$

$$\text{Similarly, } SR \parallel AC \text{ and } SR = \frac{1}{2} AC.$$

As, $PQ \parallel AC$ and $SR \parallel AC$, then also $PQ \parallel SR$

$$\text{Also, } PQ = SR, \text{ each equal to } \frac{1}{2} AC \dots (1)$$

So, $PQRS$ is a parallelogram

Now, in $\triangle SAP$ and $\triangle QBP$, we have,

$$AS = BQ$$

$$\angle A = \angle B = 90^\circ$$

$$AP = BP$$

$\therefore$ By SAS test of congruency,

$$\triangle SAP \cong \triangle QBP$$

$$\text{Hence, } PS = PQ \dots \text{by cpct} \dots (2)$$

$$\text{Similarly, } \triangle SDR \cong \triangle QCR$$

$$\therefore SR = RQ \dots \text{by cpct} \dots (3)$$

Hence, from 1, 2 and 3 we have,

$$PQ = PS = SR = RQ$$

We know that the diagonals of a square bisect each other at right angles.

$$\therefore \angle EOF = 90^\circ$$

Now, $RQ \parallel DB$

$$\Rightarrow RE \parallel FO$$

Also, $SR \parallel AC$

$$\Rightarrow FR \parallel OE$$

$\therefore OERF$ is a parallelogram.

So, $\angle FRE = \angle EOF = 90^\circ$ (Opposite angles are equal)

Thus, $PQRS$ is a parallelogram with $\angle R = 90^\circ$ and $PQ = PS = SR = RQ$.


This means that $PQRS$ is square.

Hence, the quadrilateral formed by joining the midpoints of the pairs of adjacent sides of a square is a square.

12. Question

Prove that the line segments joining the midpoints of opposite sides of a quadrilateral bisect each other.

Answer



In $\triangle ADC$, S and R are the midpoints of AD and DC respectively.

By midpoint theorem,

$$\text{Hence } SR \parallel AC \text{ and } SR = \frac{1}{2} AC \dots (1)$$

Similarly, in $\triangle ABC$, P and Q are midpoints of AB and BC respectively.

$$PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC \dots (2) \dots \text{By midpoint theorem}$$

From equations (1) and (2), we get

$$PQ \parallel SR \text{ and } PQ = SR \dots (3)$$

Here, one pair of opposite sides of quadrilateral PQRS is equal and parallel.

Hence PQRS is a parallelogram


Hence the diagonals of parallelogram PQRS bisect each other.

Thus PR and QS bisect each other.

Hence, the line segments joining the midpoints of opposite sides of a quadrilateral bisect each other.

13. Question

In the given figure, $ABCD$ is a quadrilateral whose diagonals intersect at right angles. Show that the quadrilateral formed by joining the midpoints of the pairs of adjacent sides is a rectangle.



Answer

Here, in $ABCD$, diagonals intersect at 90°

Also, in $ABCD$, P , Q , R and S be the midpoints of AB , BC , CD and DA , respectively.

In $\triangle ABC$, we have,

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC \text{ ...by midpoint theorem}$$

Similarly, in $\triangle DAC$,

$$SR \parallel AC \text{ and } SR = \frac{1}{2} AC \text{ ...by midpoint theorem}$$

Now, $PQ \parallel AC$ and $SR \parallel AC$

$$\therefore PQ \parallel SR$$

$$\text{Also, } PQ = SR = \frac{1}{2} AC$$

Hence, $PQRS$ is parallelogram.

We know that the diagonals of the given quadrilateral bisect each other at right angles.

$$\therefore \angle EOF = 90^\circ$$

Also, $RQ \parallel DB$

$$\therefore RE \parallel FO$$

Also, $SR \parallel AC$

$$\therefore FR \parallel OE$$

$\therefore OERF$ is a parallelogram.

So, $\angle FRE = \angle EOF = 90^\circ$...Opposite angles of parallelogram are equal

Thus, $PQRS$ is a parallelogram with $\angle R = 90^\circ$.

$\therefore PQRS$ is a rectangle.

CCE Questions

1. Question

Three angles of a quadrilateral are 80° , 95° and 112° . Its fourth angle is

- A. 78°
- B. 73°
- C. 85°
- D. 100°

Answer

Let the fourth angle be x

$$80^\circ + 95^\circ + 112^\circ + x^\circ = 360^\circ \text{ (Sum of angles of quadrilateral)}$$

$$287^\circ + x^\circ = 360^\circ$$

$$x = 360^\circ - 287^\circ$$

$$= 73^\circ$$

Hence, option (B) is correct

2. Question

Three angles of a quadrilateral are in the ratio 3 : 4 : 5 : 6. The smallest of these angles is

- A. 45°
- B. 60°
- C. 36°
- D. 48°

Answer

Let the angles be $3x$, $4x$, $5x$ and $6x$

$$3x + 4x + 5x + 6x = 360^\circ \text{ (Sum of angles of a quadrilateral)}$$

$$18x = 360^\circ$$

$$x = \frac{360}{18}$$

$$x = 20^\circ$$

$\therefore$ Angles of the quadrilateral are:

$$3x = 3 \times 20^\circ = 60^\circ$$

$$4x = 4 \times 20^\circ = 80^\circ$$

$$5x = 5 \times 20^\circ = 100^\circ$$


$$6x = 6 \times 20^\circ = 120^\circ$$

Hence, the smallest angle is 60°

$\therefore$ Option (B) is correct

3. Question

In the given figure, ABCD is a parallelogram in which $\angle BAD = 75^\circ$ and $\angle CBD = 60^\circ$. Then, $\angle BDC = ?$



A. 60°

B. 75°

C. 45°

D. 50°

Answer

It is given in the question that,

In parallelogram ABCD: $\angle BAD = 75^\circ$, $\angle CBD = 60^\circ$

Now, $\angle DAB = \angle DCB = 75^\circ$ (Opposite angles)

Also, in triangle DBC we know that sum of angles of a triangle is 180°

$$\angle DBC + \angle BDC + \angle DCB = 180^\circ$$

$$60^\circ + \angle BDC + 75^\circ = 180^\circ$$

$$135^\circ + \angle BDC = 180^\circ$$

$$\angle BDC = 180^\circ - 135^\circ$$

$$\angle BDC = 45^\circ$$

Hence, option (C) is correct

4. Question

In which of the following figures are the diagonals equal?

A. Parallelogram

- B. Rhombus
- C. Trapezium
- D. Rectangle

Answer

As we know that from all the quadrilaterals given below, diagonals of a rectangle are equal

Hence, option (D) is correct

5. Question

If the diagonals of a quadrilateral bisect each other at right angles, then the figure is a

- A. Trapezium
- B. Parallelogram
- C. Rectangle
- D. Rhombus

Answer

As we know that from all the quadrilaterals given below the diagonals of rhombus bisect each other at right angles

Hence, option (D) is correct

6. Question

The lengths of the diagonals of a rhombus are 16 cm and 12 cm. The length of each side of the rhombus is

- A. 10 cm
- B. 12 cm
- C. 9 cm
- D. 8 cm

Answer

Let us assume a rhombus ABCD where,

$$AB = BC = CD = DA$$

Now, in triangle OBC by using Pythagoras theorem we get:

$$BC^2 = OB^2 + OC^2$$

$$BC^2 = 6^2 + 8^2$$

$$BC^2 = 36 + 64$$

$$BC^2 = 100$$

$$BC = \sqrt{100}$$

$$BC = 10 \text{ cm}$$

$$\therefore AB = BC = CD = DA = 10 \text{ cm}$$

Hence, option (A) is correct

7. Question

The length of each side of a rhombus is 10cm and one of its diagonals is of length 16 cm. The length of the other diagonal is

A. 13 cm

B. 12 cm

C. $2\sqrt{39}$ cm

D. 6 cm

Answer

It is given in the question that,

ABCD is rhombus where, $AB = BC = CD = DA$

Now, by using Pythagoras theorem in triangle BOC we have:

$$BC^2 = OB^2 + OC^2$$

$$(10)^2 = OB^2 + (8)^2$$

$$100 = OB^2 + 64$$

$$OB^2 = 100 - 64$$

$$OB^2 = 36$$

$$OB = 6 \text{ cm}$$

$$\therefore \text{Length of diagonal, } BC = OB + OD$$

$$BC = 6 + 6$$

$$BC = 12 \text{ cm}$$

Hence, option (B) is correct

8. Question

If ABCD is a parallelogram with two adjacent angles $\angle A = \angle B$, then the parallelogram is a

A. rhombus

B. trapezium

C. rectangle

D. none of these

Answer

It is given in the question that,

ABCD is a parallelogram where two adjacent angles $\angle A = \angle B$

We know that, sum of adjacent angles is 180°

$$\therefore \angle A + \angle B = 180^\circ$$

$$2\angle A = 180^\circ$$

$$\angle A = 180/2$$

$$\angle A = 90^\circ$$

$$\text{As, } \angle A = \angle B = \angle C = \angle D = 90^\circ$$

$\therefore$ ABCD is a rectangle as all the angles are equal to 90°

Hence, option (C) is correct

9. Question

In a quadrilateral ABCD, if AO and BO are the bisectors of $\angle A$ and $\angle B$ respectively, $\angle C = 70^\circ$ and $\angle D = 30^\circ$. Then, $\angle AOB = ?$

A. 40°

B. 50°

C. 80°

D. 100°

Answer

It is given in the question that, ABCD is a quadrilateral where AO and BO are the bisectors of $\angle A$ and $\angle B$

We know that, sum of all angles of a quadrilateral is equal to 360°

$$\therefore \angle A + \angle B + \angle C + \angle D = 360^\circ$$

$$\angle A + \angle B + 70^\circ + 30^\circ = 360^\circ$$

$$\angle A + \angle B = 360^\circ - 100^\circ$$

$$\angle A + \angle B = 260^\circ$$

$$1/2 (\angle A + \angle B) = 1/2 \times 260^\circ$$

$$1/2 (\angle A + \angle B) = 130^\circ$$

Now, in triangle AOB

$$\frac{1}{2}(\angle A + \angle B) + \angle AOB = 180^\circ$$

$$130^\circ + \angle AOB = 180^\circ$$

$$\angle AOB = 180^\circ - 130^\circ$$

$$\angle AOB = 50^\circ$$

Hence, option (B) is correct

10. Question

The bisectors of any two adjacent angles of a parallelogram intersect at

A. 30°

B. 45°

C. 60°

D. 90°

Answer

We know that,

Sum of two adjacent angles = 180°

Also, sum of bisector of adjacent angles = $180/2 = 90^\circ$

As sum of angles of a triangle = 180°

$\therefore$ Sum of 2 adjacent angles + Intersection angle = 180°

$$90^\circ + \text{Intersection angle} = 180^\circ$$

$$\therefore \text{Intersection angle} = 180^\circ - 90^\circ$$

$$= 90^\circ$$

Hence, option (D) is correct

11. Question

The bisectors of the angles of a parallelogram enclose a

A. Rhombus

B. Square

C. Rectangle

D. Parallelogram

Answer

From all the given quadrilateral we know that the bisectors of the angles of a parallelogram enclose a rectangle

Hence, option (C) is correct

12. Question

The figure formed by joining the mid-points of the adjacent sides of a quadrilateral is a

- A. Rhombus
- B. Square
- C. Rectangle
- D. Parallelogram

Answer

We know that, the figure formed by joining the mid-points of the adjacent sides of a quadrilateral is a parallelogram

Hence, option (D) is correct

13. Question

The figure formed by joining the mid-points of the adjacent sides of a square is a

- A. Rhombus
- B. Square
- C. Rectangle
- D. Parallelogram

Answer

We know that, the figure formed by joining the mid-points of the adjacent sides of a square is a square

Hence, option (B) is correct

14. Question

The figure formed by joining the mid-points of the adjacent sides of a parallelogram is a

- A. rhombus
- B. square
- C. rectangle
- D. parallelogram

Answer

We know that, the figure formed by joining the mid-points of the adjacent sides of a parallelogram is parallelogram

Hence, option (D) is correct

15. Question

The figure formed by joining the mid-points of the adjacent sides of a rectangle is a

- A. rhombus
- B. square
- C. rectangle
- D. parallelogram

Answer

We know that, the figure formed by joining the mid-points of the adjacent sides of a rectangle is a rhombus

Hence, option (A) is correct

16. Question

The figure formed by joining the mid-points of the adjacent sides of a rhombus is a

- A. rhombus
- B. square
- C. rectangle
- D. parallelogram

Answer

We know that, the figure formed by joining the mid-points of the adjacent sides of a rhombus is a rectangle

Hence, option (C) is correct

17. Question

If an angle of a parallelogram is two-third of its adjacent angle, the smallest angle of the parallelogram is

- A. 108°
- B. 54°
- C. 72°
- D. 81°

Answer

We know that,

Sum of two adjacent angles is equal to 180°

$$\therefore \angle A + \angle B = 180^\circ$$

According to the condition given in the question, we have

$$\angle A = x^\circ \text{ then } \angle B = \frac{2}{3} x^\circ$$

$$\therefore x^\circ + \frac{2x}{3}^\circ = 180^\circ$$

$$\frac{5x}{3}^\circ = 180^\circ$$

$$\Rightarrow x = \frac{180 \times 3}{5}$$

$$\Rightarrow x = 540^\circ / 5$$

$$\Rightarrow x = 108^\circ$$

$$\therefore \angle A = 108^\circ \text{ and,}$$

$$\angle B = \frac{2}{3} \times 108^\circ$$

$$\angle B = 2 \times 36^\circ = 72^\circ$$

Thus, the smallest angle = $\angle B = 72^\circ$

Hence, option (C) is correct

18. Question

If one angle of a parallelogram is 24° less than twice the smallest angle, then the largest angle of the parallelogram is

A. 68°

B. 102°

C. 112°

D. 136°

Answer

As per the question,

Let the smallest angle be x° and the largest angle be $(2x - 24)^\circ$

Since, the sum of adjacent angles of a parallelogram is 180°

$$\therefore x + (2x - 24) = 180^\circ$$

$$3x - 24 = 180^\circ$$


$$x = 68^\circ$$

$$\text{Hence, the largest angle is: } 2x - 24 = 2(68) - 24 = 136 - 24 = 112$$

$\therefore$ Option A is correct

19. Question

In the given figure, ABCD is a parallelogram in which $\angle BDC = 45^\circ$ and $\angle BAD = 75^\circ$. Then, $\angle CBD = ?$



A. 45°

B. 55°

C. 60°

D. 75°

Answer

As per the question,

$\angle BAD = \angle BCD = 75^\circ$ (opposite angles of parallelogram)

Now, in $\triangle ABC$,

$$\angle BAC + \angle ABC + \angle BCD = 180^\circ$$

$$45 + \angle ABC + 75 = 180^\circ$$

$$\angle ABC = 60^\circ$$

$\therefore$ Option C is correct

20. Question

If area of a ||gm with sides a and b is A and that of a rectangle with sides a and b is B, then

A. $A > B$

B. $A = B$

C. $A < B$

D. $A \geq B$

Answer

Let the height of the parallelogram be 'h'

Now, $h < b$ (Since, perpendicular distance is the shortest)


$$\therefore a \times h < a \times b$$

$$A < B$$

$\therefore$ Option C is correct

21. Question

In the given figure, ABCD is a ||gm and E is the mid-point of BC. Also, DE and AB when produced meet at F. Then,



A. $AF = \frac{3}{2}AB$

B. $AF = 2AB$

C. $AF = 3AB$

D. $AF^2 = 2AB^2$

Answer

According to the condition given in the question, we have

In triangle DCE and FBE

$$BE = EC \text{ (E is the mid-point of BC)}$$

$$\angle CED = \angle BEF \text{ (Vertically opposite angles)}$$

$$\angle CDE = \angle EFB \text{ (Alternate interior angles)}$$

$$\therefore \triangle DCE \cong \triangle FBE \text{ (By AAS congruence rule)}$$

$$DC = BF \text{ (BY CPCT)}$$

$$\text{As AB is parallel to DC, then } AB = DC$$

$$\therefore AB = DC = BF$$

$$AF = AB + BF$$

$$AF = AB + AB$$

$$AF = 2AB$$

Hence, option (B) is correct

22. Question

The parallel sides of a trapezium are a and b respectively. The line joining the mid-points of its non-parallel sides will be

A. $\frac{1}{2}(a - b)$

B. $\frac{1}{2}(a + b)$

C. $\frac{2ab}{(a + b)}$

D. $\sqrt{ab}$

Answer

It is given in the question that,

ABCD is a trapezium

Draw EF parallel to AB and DC, and join BD intersecting EF at point M.

Now, E is the midpoint of AD and $EM \parallel AB$. Hence, using midpoint theorem,

$$EM = \frac{1}{2} AB$$

$$\Rightarrow EM = \frac{1}{2} b$$

$$\text{Similarly, } FM = \frac{1}{2} a$$

$$\Rightarrow DC = \frac{1}{2} a$$

$$EF = EM + FM$$


$$EF = \frac{1}{2} a + \frac{1}{2} b$$

$$EF = \frac{1}{2} (a + b)$$

$\therefore$ Option B is correct

23. Question

In a trapezium ABCD, if E and F be the mid-point of the diagonals AC and BD respectively. Then, EF = ?



A. $\frac{1}{2} AB$

B. $\frac{1}{2} CD$

C. $\frac{1}{2}(AB + CD)$

D. $\frac{1}{2}(AB - CD)$

Answer

Construction: Join CF and extend it to cut AB at point M

Firstly, in triangle MFB and triangle DFC

DF = FB (As F is the mid-point of DB)

$\angle DFC = \angle MFB$ (Vertically opposite angle)

$\angle DFC = \angle FBM$ (Alternate interior angle)

$\therefore$ By ASA congruence rule

$\triangle MFB \cong \triangle DFC$

Now, in triangle CAM

E and F are the mid-points of AC and CM respectively

$\therefore EF = \frac{1}{2}(AM)$


$EF = \frac{1}{2}(AB - MB)$

$EF = \frac{1}{2}(AB - CD)$

Hence, option D is correct

24. Question

In the given figure, ABCD is a parallelogram, M is the mid-point of BD and BD bisects $\angle B$ as well as $\angle D$. Then, $\angle AMB = ?$



A. 45°

B. 60°

C. 90°

D. 30°

Answer

Since, ABCD is a parallelogram,

$$\therefore \angle B = \angle D \text{ (opposite angle)}$$

$$\frac{1}{2} \angle B = \frac{1}{2} \angle D$$

$$\angle ADB = \angle ABD$$

$\therefore$ ADB is an isosceles triangle.

Since, M is the midpoint of BD


$\therefore$ AM is a median of $\triangle ADB$.

Now, $\angle AMB = 90^\circ$ (AM is perpendicular to BD)

$\therefore$ Option C is correct

25. Question

In the given figure, ABCD is a rhombus. Then,



A. $AC^2 + BD^2 = AB^2$

B. $AC^2 + BD^2 = 2AB^2$

C. $AC^2 + BD^2 = 4AB^2$

D. $2(AC^2 + BD^2) = 3AB^2$

Answer

Since, we know that the diagonals of a rhombus bisect each other at 90° .

Hence, $OA = \frac{1}{2} AC$, $OB = \frac{1}{2} BD$ and $\angle AOB = 90^\circ$

$$AB^2 = OA^2 + OB^2$$

$$AB^2 = \left(\frac{1}{2} AC\right)^2 + \left(\frac{1}{2} BD\right)^2$$

$$= \frac{1}{4}(AC)^2 + \frac{1}{4}(BD)^2$$


$$AB^2 = \frac{1}{4}(AC^2 + BD^2)$$

$$4AB^2 = (AC^2 + BD^2)$$

∴ Option C is correct

26. Question

In a trapezium ABCD, if $AB \parallel CD$, then $(AC^2 + BD^2) = ?$



- A. $BC^2 + AD^2 + 2BC \cdot AD$
- B. $AB^2 + CD^2 + 2AB \cdot CD$
- C. $AB^2 + CD^2 + 2AD \cdot BC$
- D. $BC^2 + AD^2 + 2AB \cdot CD$

Answer

Draw perpendicular from D on AB meeting it on E and from C on AB meeting AB at F

∴ DEFC will be a parallelogram and thus, $EF = CD$

Now, In $\triangle ABC$

Since, $\angle B$ is acute

$$\therefore AC^2 = BC^2 + AB^2 - 2AB \times AE \text{ (i)}$$

Similarly, In $\triangle ABD$,

Since $\angle A$ is acute

$$\therefore BD^2 = AD^2 + AB^2 - 2AB \times AF \text{ (ii)}$$

Adding (i) and (ii),

$$AC^2 + BD^2 = (BC^2 + AD^2) + (AB^2 + AB^2) - 2AB (AE + BF)$$

$$= (BC^2 + AD^2) + 2AB (AB - AE - BF) \text{ [Since, } AB = AE + EF + FB \text{ and } AB - AE = BE]$$

$$= (BC^2 + AD^2) + 2AB (BE - BF)$$

$$= (BC^2 + AD^2) + 2AB \cdot EF$$

Now, we know that $CD = EF$

$$\text{Thus, } AC^2 + BD^2 = (BC^2 + AD^2) + 2AB \cdot CD$$

∴ Option D is correct

27. Question

Two parallelograms stand on equal bases and between the same parallels. The ratio of their areas is

- A. 1:2
- B. 2:1
- C. 1:3
- D. 1:1

Answer

We know that,

Area of a parallelogram = base $\times$ height


Now, if both parallelograms are on the same base and between the same parallels, then their heights will be equal.

Hence, their areas will also be equal

$\therefore$ Option D is correct

28. Question


In the given figure, AD is a median of $\triangle ABC$ and E is the mid-point of AD. If BE is joined and produced to meet AC in F, then AF = ?



- A. $\frac{1}{2}AC$
- B. $\frac{1}{3}AC$
- C. $\frac{2}{3}AC$
- D. $\frac{3}{4}AC$

Answer

Let G be the mid-point of FC and join DG



In $\triangle BCF$,

G is the mid-point of FC and D is the mid-point of BC

Thus, $DG \parallel BF$

$DG \parallel EF$

Now, In $\triangle ADG$,

E is the mid-point of AD and EF is parallel to DG.

Thus, F is the mid-point of AG.

$AF = FG = GC$ [G is the mid-point of FC]

Hence, $AF = \frac{1}{3} AC$

$\therefore$ Option B is correct

29. Question

If $\angle A$, $\angle B$, $\angle C$ and $\angle D$ of a quadrilateral ABCD taken in order, are in the ratio 3 : 7 : 6 : 4, then ABCD is a

A. Rhombus

B. Kite

C. Trapezium

D. Parallelogram

Answer

Let the required angles be $3x$, $7x$, $6x$ and $4x$

$3x + 7x + 6x + 4x = 360^\circ$ (Sum of angles of quadrilateral)

$20x = 360^\circ$

$x = 18^\circ$

Hence, angles are:

$3x = 3 \times 18^\circ = 54^\circ$

$7x = 7 \times 18^\circ = 126^\circ$

$6x = 6 \times 18^\circ = 108^\circ$

$$4x = 4 \times 18^\circ = 72^\circ$$

Now we can observe that, $54^\circ + 126^\circ = 180^\circ$ and $72^\circ + 108^\circ = 180^\circ$

Thus, ABCD is a trapezium.

Hence option C is correct.

30. Question

Which of the following is not true for a parallelogram?

- A. Opposite sides are equal.
- B. Opposite angles are equal.
- C. Opposite angles are bisected by the diagonals.
- D. Diagonals bisect each other.

Answer

We know that,

In any parallelogram, opposite angles are bisected by the diagonals

$\therefore$ Option C is correct

31. Question

If APB and CQD are two parallel lines, then the bisectors of $\angle APQ$, $\angle BPQ$, $\angle CQP$ and $\angle PQD$ enclose a

- A. square
- B. rhombus
- C. rectangle
- D. kite

Answer

It is given in the question that,


APB and CQD are two parallel lines,

Thus, the bisectors of $\angle CQP$, $\angle APQ$, $\angle BPQ$ and $\angle PQD$ enclose a rectangle.

Hence, option C is correct.

32. Question

The diagonals AC and BD of a parallelogram ABCD intersect each other at the point O such that $\angle DAC = 30^\circ$ and $\angle AOB = 70^\circ$. Then, $\angle DBC = ?$



A. 40°

B. 35°

C. 45°

D. 50°

Answer

In the given figure,

$\angle OAD = \angle OCB$ (Alternate interior angle)

$\angle OCB = 30^\circ$

$\angle AOB + \angle BOC = 180^\circ$ (Linear pair)

$70^\circ + \angle BOC = 180^\circ$

$\angle BOC = 110^\circ$

Now, In $\triangle BOC$,

$\angle OBC + \angle BOC + \angle OCB = 180^\circ$

$\angle OBC + 110^\circ + 30^\circ = 180^\circ$

$\angle OBC = 40^\circ$

$\therefore \angle DBC = 40^\circ$

Hence, Option A is correct.

33. Question

Three statements are given below:

I. In a $\parallel$ gm, the angle bisectors of two adjacent angles enclose a right angle.

II. The angle bisectors of a $\parallel$ gm form a rectangle.

III. The triangle formed by joining the mid-points of the sides of an isosceles triangle is not necessarily an isosceles triangle.

Which is true?

A. I only

B. II only

- C. I and II
- D. II and III

Answer

We can clearly observe that statement I and statement II are correct. Whereas Statement III is not correct because the triangle formed by joining the midpoints of the sides of an isosceles triangle is always an isosceles triangle

Therefore, Option C is correct

34. Question

Three statements are given below:

- I. In a rectangle ABCD, the diagonal AC bisects $\angle A$ as well as $\angle C$.
- II. In a square ABCD, the diagonal AC bisects $\angle A$ as well as $\angle C$.
- III. In a rhombus ABCD, the diagonal AC bisects $\angle A$ as well as $\angle C$.

Which is true?

- A. I only
- B. II and III
- C. I and III
- D. I and II

Answer

We can clearly observe that statement II and statement III are correct and Statement I is wrong because the diagonals of a rectangle does not bisect $\angle A$ and $\angle C$. And this is so because the adjacent sides are unequal in a rectangle.

$\therefore$ Option B is correct

35. Question

In each of the questions one question is followed by two statements I and II. Choose the correct option.

Is quadrilateral ABCD a ||gm?

- I. Diagonals AC and BD bisect each other.
 - II. Diagonals AC and BD are equal.
- A. if the question can be answered by one of the given statements alone and not by the other;
 - B. if the question can be answered by either statement alone;
 - C. if the question can be answered by both the statements together but not by any one of the two;
 - D. if the question cannot be answered by using both the statements together.

Answer

Here, as we know that if the diagonals of a quadrilateral bisect each other, then it is a parallelogram.

But as per II, if the diagonals of a quadrilateral are equal, then it is not necessarily a parallelogram which is not true. Thus, II does not give the answer.

Therefore Option A is correct.

36. Question

In each of the questions one question is followed by two statements I and II. Choose the correct option.

Is quadrilateral ABCD a rhombus?

I. Quad. ABCD is a ||gm.

II. Diagonals AC and BD are perpendicular to each other.

A. if the question can be answered by one of the given statements alone and not by the other;

B. if the question can be answered by either statement alone;

C. if the question can be answered by both the statements together but not by any one of the two;

D. if the question cannot be answered by using both the statements together.

Answer

Here, we can observe that neither I nor II can alone justify the answer to the given question. But if we consider both I and II together then they completely satisfy the answer.

∴ Option C is correct.

37. Question

In each of the questions one question is followed by two statements I and II. Choose the correct option.

Is ||gm ABCD a square?

I. Diagonals of ||gm ABCD are equal.

II. Diagonals of ||gm ABCD intersect at right angles.

A. if the question can be answered by one of the given statements alone and not by the other;

B. if the question can be answered by either statement alone;

C. if the question can be answered by both the statements together but not by any one of the two;

D. if the question cannot be answered by using both the statements together.

Answer

We know that when the diagonals of a parallelogram are equal, it might be a square or a rectangle. But if the diagonals of that parallelogram intersect at a right angle, then it is definitely a square. Thus, it can be concluded that both I and II together will give the answer.

Therefore, Option C is correct.

38. Question

In each of the questions one question is followed by two statements I and II. Choose the correct option.

Is quad. ABCD a parallelogram?

I. Its opposite sides are equal.

II. Its opposite angles are equal.

A. if the question can be answered by one of the given statements alone and not by the other;

B. if the question can be answered by either statement alone;

C. if the question can be answered by both the statements together but not by any one of the two;

D. if the question cannot be answered by using both the statements together.

Answer

We know that a quadrilateral is a parallelogram when either I or II holds true.

Hence, the correct answer is (b)

39. Question

Each question consists of two statements, namely, Assertion (A) and Reason (R). Choose the correct option.

Assertion (A)	Reason (R)
If three angles of a quadrilateral are 130° , 70° , and 60° , then the fourth angle is 100° .	The sum of all the angle of a quadrilateral is 360° .

A. Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).

B. Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).

C. Assertion (A) is true and Reason (R) is false.

D. Assertion (A) is false and Reason (R) is true.

Answer

Let the fourth angle be x ,

$$130^\circ + 70^\circ + 60^\circ + x^\circ = 360^\circ \text{ (angle sum of quadrilateral)}$$

$$x^\circ = 360^\circ - (130^\circ + 70^\circ + 60^\circ)$$

$$x^\circ = 100^\circ$$

Thus, it can be observed that reason and assertion both are true and the reason explains the assertion.

Therefore Option A is correct.

40. Question

Each question consists of two statements, namely, Assertion (A) and Reason (R). Choose the correct option.

Assertion (A)	Reason (R)
ABCD is a quadrilateral in which P, Q, R and S are the mid-points of AB, BC, CD and DA respectively. Then, PQRS is a parallelogram.	The line segment joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

- A. Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).
- B. Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).
- C. Assertion (A) is true and Reason (R) is false.
- D. Assertion (A) is false and Reason (R) is true.

Answer

It is given that, ABCD is a quadrilateral in which P, Q, R and S are the mid-points of AB, BC, CD and DA respectively. Then, PQRS is a parallelogram

Also, the line segment joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

Hence, both assertion and reason are true and reason is correct explanation of the assertion

$\therefore$ Option (a) is correct

41. Question

Each question consists of two statements, namely, Assertion (A) and Reason (R). Choose the correct option.

Assertion (A)	Reason (R)
In a rhombus ABCD, the diagonal AC bisects $\angle A$ as well as $\angle C$.	The diagonals of a rhombus bisect each other at right angles.

- A. Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).
- B. Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).
- C. Assertion (A) is true and Reason (R) is false.
- D. Assertion (A) is false and Reason (R) is true.

Answer

It is given that,

In a rhombus ABCD, the diagonal AC bisects $\angle A$ as well as $\angle C$ which is true

And we know that, the diagonals of a rhombus bisect each other at right angles.

Hence, both assertion and reason are true but reason is not the correct explanation of assertion

$\therefore$ Option (b) is correct

42. Question

Each question consists of two statements, namely, Assertion (A) and Reason (R). Choose the correct option.

Assertion (A)	Reason (R)
Every parallelogram is a rectangle.	The angle bisectors of a parallelogram form a rectangle.

- A. Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).
- B. Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).
- C. Assertion (A) is true and Reason (R) is false.
- D. Assertion (A) is false and Reason (R) is true.

Answer

The statement given in assertion is not true as every parallelogram is not a rectangle whereas, statement given in the reason is true as the angle bisectors of a parallelogram form a rectangle

Hence, assertion is false whereas reason is true

∴ Option (d) is correct

43. Question

Each question consists of two statements, namely, Assertion (A) and Reason (R). Choose the correct option.

Assertion (A)	Reason (R)
The diagonals of a gm bisect each other.	If the diagonals of a gm are equal and intersect at right angles, then the parallelogram is a square.

- A. Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).
- B. Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).
- C. Assertion (A) is true and Reason (R) is false.
- D. Assertion (A) is false and Reason (R) is true.

Answer

We know that,

The diagonals of a ||gm bisect each other

Also we know that, if the diagonals of a ||gm are equal and intersect at right angles, then the parallelogram is a square

Hence, both assertion and reason are true but reason is not the correct explanation of the assertion

Hence, option (b) is correct

44. Question

Match the following columns:

Column I	Column II
(a) Angle bisectors of a parallelogram form a	(p) parallelogram
(b) The quadrilateral formed by joining the mid-points of the pairs of adjacent sides of a square is a	(q) rectangle
(c) The quadrilateral formed by joining the mid-points of the pairs of adjacent sides of a rectangle is a	(r) square
(d) The figure formed by joining the mid-points of the pairs of adjacent sides of a quadrilateral is a	(s) rhombus

The correct answer is:

(a) -....., (b) -.....,

(c) -....., (d) -.....,

Answer



The correct match for the above given table is as follows:

Column I	Column II
(a) Angle bisectors of a parallelogram form a	(q) Rectangle
(b) The quadrilateral formed by joining the mid-points of the pairs of adjacent sides of a square is a	(r) Square
(c) The quadrilateral formed by joining the mid-points	(s) Rhombus
(d) The figure formed by joining the mid-points of the pairs of adjacent sides of a quadrilateral is a	(p) Parallelogram

45. Question

Match the following columns:

Column I	Column II
(a) In the given figure, ABCD is a trapezium in which $AB = 10$ cm and $CD = 7$ cm. If P and Q are the mid-points of AD and BC respectively. then $PO =$	(p) equal

	
(b) In the given figure, PQRS is a gm whose diagonals intersect at O. If PR = 13 cm, then OR = 	(q) at right angles
(c) The diagonals of a square are	(r) 8.5 cm
(d) The diagonals of a rhombus bisect each other	(s) 6.5 cm

The correct answer is:

(a) -....., (b) -.....,

(c) -....., (d) -.....,

Answer

$$a) PQ = \frac{1}{2} (AB + CD)$$

$$PQ = \frac{1}{2} (17)$$

$$PQ = 8.5 \text{ cm}$$

$$(b) OR = \frac{1}{2} (PR)$$

$$OR = \frac{1}{2} (13)$$

$$OR = 6.5 \text{ cm}$$

(c) We know that,

The diagonals of a square are equal

(d) We also know that,

The diagonals of a rhombus bisect each other at right angles

$\therefore$ The correct match is as follows:

(a) - (r)

(b) - (s)

(c) - (p)

(d) - (q)

Formative Assessment (Unit Test)

1. Question

Which is false?

- A. In a $\parallel\text{gm}$, the diagonals are equal.
- B. In a $\parallel\text{gm}$, the diagonals bisect each other.
- C. If a pair of opposite sides of a quadrilateral is equal, then it is a $\parallel\text{gm}$.
- D. If the diagonals of a $\parallel\text{gm}$ are perpendicular to each other, then it is a rhombus.


Answer

from the above given four statements option A is false as we know that in any parallelogram the diagonals are not equal

Hence, option A is correct

2. Question

If P is a point on the median AD of a $\triangle ABC$, then $\text{ar}(\triangle ABP) = \text{ar}(\triangle ACP)$.



A. True

B. False

Answer

In $\triangle ABC$,

Since, AD is the median

Thus, $BD = DC$

Let the height of $\triangle ABC$ be h

$$\text{ar}(\triangle ABD) = \text{ar}(\triangle ABD)$$

$$\frac{1}{2} \times h \times BD = \frac{1}{2} \times h \times BD$$

$$\frac{1}{2} \times h \times BD = \frac{1}{2} \times h \times CD$$

$$\therefore \text{ar}(\triangle ABD) = \text{ar}(\triangle ADC)$$

Let H be the height of $\triangle BPD$ and $\triangle PDC$

$$\therefore \text{ar}(\triangle BPD) = \text{ar}(\triangle PDC)$$

$$\text{Now, ar}(\triangle ABD) = \text{ar}(\triangle ABP) + \text{ar}(\triangle BPD)$$

$$\text{And, ar}(\triangle ACD) = \text{ar}(\triangle ACP) + \text{ar}(\triangle PDC)$$

$$\text{Thus, ar}(\triangle ABP) = \text{ar}(\triangle ACP)$$

$\therefore$ Option A is correct

3. Question

The angles of a quadrilateral are in the ratio 1:3:5:6. Find its greatest angle.

Answer

Let the angles be x, 3x, 5x and 6x.

$$x + 3x + 5x + 6x = 360^\circ \text{ (sum of angles of quadrilateral)}$$

$$15x^\circ = 360^\circ$$

$$x^\circ = 24^\circ$$

Therefore, angles are as follows:

$$x^\circ = 24^\circ$$

$$3x^\circ = 24^\circ \times 3 = 72^\circ$$


$$5x^\circ = 24^\circ \times 5 = 120^\circ$$

$$6x^\circ = 24^\circ \times 6 = 144^\circ$$

Hence, 144° is the greatest angle.

4. Question

In a $\triangle ABC$, D and E are the mid-points of AB and AC respectively and $DE = 5.6$ cm. Find the length of BC.



Answer

We know that in $\triangle ABC$, D and E are the midpoints of AB and AC, respectively.

Now using mid-point theorem,

$$DE = \frac{1}{2} (BC)$$

$$BC = 2 \times DE$$


$$BC = 2 \times 5.6$$

$$= 11.2 \text{ cm}$$

Thus, $BC = 11.2$ cm

5. Question

In the given figure, AD is the median and $DE \parallel AB$. Prove that BE is the median.



Answer

In $\triangle ABC$, using mid point theorem

We know that D is the mid-point of BC and $DE \parallel AB$.


$$\text{Thus, } AE = EC \text{ and } DE = \frac{1}{2} (AB)$$

Now, E is the mid point of AC

Thus, BE is the median

6. Question

In the given figure, lines l, m and n are parallel lines and the lines p and q are transversals. If $AB = 5$ cm, $BC = 15$ cm, then $DE : EF = ?$



Answer

Here, we have:

$$l \parallel m \parallel n$$

And p and q are the transversal lines

$$\text{Thus, } AB : BC = 5 : 15$$

$$AB : BC = 1 : 3$$

$\therefore$ Using intercept theorem,


$$DE : EF = 1 : 3$$

7. Question

ABCD is a rectangle in which diagonal BD bisects $\angle B$. Show that ABCD is a square.

Answer

Let there be a rectangle ABCD with $AB = CD$ and $BC = AD$ and $\angle A = \angle B = \angle C = \angle D = 90^\circ$



Since, BD bisects $\angle B$

$$\angle ABD = \angle DBC \text{ (i)}$$

And, $\angle ADB = \angle DBC$ [Alternate interior angles]

$$\angle ABD = \angle ADB. \text{ [From (i)]}$$

$AB = DA$. (Sides opposite to equal angles)


$\therefore AB = CD = DA = BC$

Since, all the sides are equal and all the angles are equal to 90° , thus the quadrilateral is a square.

Hence, ABCD is a square.

8. Question

The diagonals of a rectangle ABCD intersect at the point O. If $\angle BOC = 50^\circ$, then $\angle OAD = ?$



A. 50°

B. 55°

C. 65°

D. 75°

Answer

$\angle BOC = \angle AOD$ (Vertically opposite angles)

Angle AOD = 50°

In $\triangle AOD$, Since, the diagonals are equal, thus the bisectors will also be equal)

Thus, $OA = OD$

$\therefore \angle OAD = \angle ODA$

$$= \frac{1}{2} (180^\circ - 50^\circ)$$

$$= \frac{1}{2} (130^\circ)$$

$$= 65^\circ$$

$\therefore$ Option C is correct

9. Question

Match the following column:

Column I	Column II
(a) Sum of all the angles of a quadrilateral is	(p) Right angles
(b) In a gm, the angle bisectors of two adjacent angles intersect at	(q) Rectangle
(c) Angle bisectors of a gm form a	(r) 90°
(d) The diagonals of a square are equal and bisect each other at an angle of	(s) 4 right angles

The correct answer is:

(a) -....., (b) -.....,

(c) -....., (d) -.....,


Answer

The correct match for the above given table is as follows:

Column I	Column II
(a) Sum of all the angles of a quadrilateral is	(s) 4 right angles
(b) In a gm, the angle bisectors of two adjacent angles intersect at	(p) Right angles
(c) Angle bisectors of a gm form a	(q) Rectangle
(d) The diagonals of a square are equal and bisect each other at an angle of	(r) 90°

10. Question

The diagonals of a rhombus, ABCD intersect at the point O. If $\angle BDC = 50^\circ$, then $\angle OAB = ?$



- A. 50°
- B. 40°
- C. 25°

D. 20°

Answer

$\angle BDC = \angle ABD$ (Alternate interior angles)

$\angle ABD = 50^\circ$

Now, In $\triangle AOB$,

$\angle DBA = 50^\circ$ and $\angle AOB = 90^\circ$

Thus, $\angle OAB = 180^\circ - (90^\circ + 50^\circ)$


$\angle OAB = 180^\circ - 140^\circ$

$\angle OAB = 40^\circ$

$\therefore$ Option B is correct.

11. Question

ABCD is a trapezium in which $AB \parallel CD$ and $AD = BC$, then $\angle A = \angle B$ is



A. true

B. false

Answer

Construction: Draw perpendicular line from D and C to AB such that it cuts AB at F and E, respectively.

Now, In $\triangle ADF$ and $\triangle BCE$,

$AD = BC$ (Given)

$\angle AFD = \angle BEC$ (90° each)

$DF = CE$ (Perpendicular distance between the same parallels)

$\therefore$ By SSA axiom

$\triangle ADF \cong \triangle BCE$

$\angle A = \angle B$ (by c.p.c.t.)

Therefore Option A is correct.

12. Question

Look at the statements given below:

I. If AD, BE and CF be the altitudes of a $\triangle ABC$ such that $AD = BE = CF$, then $\triangle ABC$ is an equilateral triangle.

II. If D is the mid-point of hypotenuse AC of a right $\triangle ABC$, then $BD = AC$.

III. In an isosceles $\triangle ABC$ in which $AB = AC$, the altitude AD bisects BC.

Which is true?

A. I only B. II only


C. I and III D. II and III

Answer

We can clearly observe that statement I and statement III are correct.

We can prove the statement as follows:

In $\triangle ABC$, altitudes AD, BE and CF are equal



Now, In $\triangle ABE$ and $\triangle ACF$,

$BE = CF$ (Given)

$\angle A = \angle A$ (common)

$\angle AEB = \angle AFC$ (Each 90°)

Therefore, by AAS axiom,

$\triangle ABE \cong \triangle ACF$

$AB = AC$ (by cpct)

In the same way, $\triangle BCF \cong \triangle BAD$


thus, $BC = AB$ (by cpct)

Therefore $AB = AC = BC$

Thus, $\triangle ABC$ is an equilateral triangle.

We can prove the IIIrd statement as follows:

Let $\triangle ABC$ be an isosceles triangle with AD as an altitude



Now, In $\triangle ABD$ and $\triangle ADC$,

$AB = AC$ (Given)

$\angle B = \angle C$ (Angles opposite to equal sides)

$\angle BDA = \angle CDA$ (each 90°)

Therefore by AAS axiom,

$\triangle ABD \cong \triangle ADC$

$BD = DC$ (by congruent parts of congruent triangles)


$\therefore D$ is the mid-point of BC and hence AD bisects BC .

13. Question

In the given figure, D and E are two points on side BC of $\triangle ABC$ such that $BD = DE = EC$.

Prove that

$\text{ar}(\triangle ABD) = \text{ar}(\triangle ADE) = \text{ar}(\triangle AEC)$.



Answer

Area of a triangle = $\frac{1}{2}$ (Base $\times$ Height)

Now, draw AL perpendicular to BC and h be the height of $\triangle ABC$ i.e. AL

Thus, Height of $\triangle ABD$ = Height of $\triangle ADE$ = Height of $\triangle AEC$


It is given that the bases BD , DE and EC of $\triangle ABD$, $\triangle ADE$ and $\triangle AEC$ respectively are equal.

Now, since base and height both are equal of all the triangles therefore,

$\text{ar}(\triangle ABD) = \text{ar}(\triangle ADE) = \text{ar}(\triangle AEC)$

14. Question

In the given figure $ABCD$, $DCFE$ and $ABFE$ are parallelograms. Show that $\text{ar}(\triangle ADE) = \text{ar}(\triangle BCF)$.



Answer

Now, here in $\triangle ADE$ and $\triangle BCF$,

$AD = BC$ (Opposite sides of parallelogram ABCD)

$DE = CF$ (Opposite sides of parallelogram DCEF)

$AE = BF$ (Opposite sides of parallelogram ABFE)

$\therefore$ By SSS axiom,


$\triangle ADE \cong \triangle BCF$

And,

$\text{ar}(\triangle ADE) = \text{ar}(\triangle BCF)$ (By cpct)

15. Question

In the given figure, ABCD is a trapezium in which $AB \parallel DC$ and diagonals AC and BD intersect at O. Prove that $\text{ar}(\triangle AOD) = \text{ar}(\triangle BOC)$.



Answer

Here, in trapezium ABCD,

$AB \parallel DC$ and AC and BD are the diagonals intersecting at O.

Now, since $\triangle ACD$ and $\triangle BCD$ lie on the same base and between the same parallels.

Thus, $\text{ar}(\triangle ACD) = \text{ar}(\triangle BCD)$

Subtracting $\text{ar}(\triangle COD)$ from both the sides, we get:

$\text{ar}(\triangle ACD) - \text{ar}(\triangle COD) = \text{ar}(\triangle BCD) - \text{ar}(\triangle COD)$


$\therefore \text{ar}(\triangle AOD) = \text{ar}(\triangle BOC)$

16. Question

Show that a diagonal divides a parallelogram into two triangles of equal area.

Answer

Let there be a parallelogram ABCD and with one of its diagonal as AC.



Now, In $\triangle CDA$ and $\triangle ABC$,

$DA = BC$ (Opposite sides of parallelogram ABCD)

$AC = AC$ (Common)

$CD = AB$ (Opposite sides of parallelogram ABCD)

$\therefore$ By SSS axiom

$\triangle CDA \cong \triangle ABC$


$\text{ar}(\triangle CDA) = \text{ar}(\triangle ABC)$ (by cpct)

Thus, we can say that the diagonal of a parallelogram divides it into two triangles of equal area.

17. Question

In the given figure, AC is a diagonal of quad. ABCD in which $BL \perp AC$ and $DM \perp AC$. Prove that or

$$(\text{quad. ABCD}) = \frac{1}{2} \times AC \times (BL + DM).$$



Answer

Here we have ABCD as a quadrilateral with one of its diagonal as AC and BL and DM are perpendicular to AC

Thus, $\text{ar}(\text{ABCD}) = \text{ar}(\triangle ADC) + \text{ar}(\triangle ABC)$


Since, $(BL \perp AC)$ and $(DM \perp AC)$

$$\therefore \text{Area of ABCD} = \left(\frac{1}{2} \times AC \times BL\right) + \left(\frac{1}{2} \times AC \times DM\right)$$

$$= \frac{1}{2} \times AC \times (BL + DM)$$

18. Question

||gm ABCD and rectangle ABEF have the same base AB and are equal in areas. Show that the perimeter of the ||gm is greater than that of the rectangle.



Answer

Here we know that parallelogram ABCD and rectangle ABEF are on the same base AB and between the same parallels such that:

$$AB = CD \text{ and } AB = EF$$

$$\text{So, } CD = FE$$

Now, adding AB on both sides

$$AB + CD = AB + FE \text{ (i)}$$

Since we know that hypotenuse is the longest side of a triangle

$$\therefore AD > AF \text{ (ii)}$$

$$\text{And, } BC > BE \text{ (iii)}$$

Adding (ii) and (iii),

$$AD + BC > AF + BE \text{ (iv)}$$

$$\text{Now, Perimeter of ABCD} = AB + BC + CD + AD$$

$$\text{And, Perimeter of ABEF} = AB + BE + FE + AF$$

Adding (i) and (iv),

$$AB + CD + AD + BC > AB + FE + AF + BE$$

Thus, we can say that the perimeter of parallelogram ABCD is greater than that of rectangle ABEF.

19. Question

In the adjoining figure, ABCD is a ||gm and E is the mid-point of side BC. If DE and AB when produced meet at F, prove that $AF = 2AB$.

Answer

Here we have parallelogram ABCD with $AB \parallel DC$

Thus, $DC \parallel BF$

Now, in $\triangle DEC$ and $\triangle FEB$,

$$\angle DCF = \angle EBF \text{ (Alternate interior angle)}$$

$$CE = BE \text{ (E is the mid-point of BC)}$$

$\angle CED = \angle BEF$ (Vertically opposite angle)

Therefore, by ASA axiom,

$$\triangle DEC \cong \triangle FEB$$

$$CD = BF \text{ (by cpct)}$$


And $CD = AB$ (Opposite sides of a parallelogram ABCD)

$$\text{So, } AF = AB + BF = AB + AB = 2AB$$

20. Question

In the adjoining figure, ABCD and PQRC are rectangles, where Q is the mid-point of AC.

Prove that (i) $DP = PC$ (ii) $PR = \frac{1}{2}AC$.




Answer

(i) Here, we have

$$\angle CRQ = \angle CBA = 90^\circ$$

Thus, $RQ \parallel AB$

Now, In $\triangle ABC$,



Q is the mid-point of AC and $QR \parallel AB$.

Thus, R is the mid-point of BC.

In the same way, P is the midpoint of DC.

Hence, $DP = PC$

(ii) Here, let us join B to D.

Now, In $\triangle CDB$,

P and R are the mid points of DC and BC respectively.

Since, $AC = BD$

Thus, $PR \parallel DB$ and $PR = \frac{1}{2} DB = \frac{1}{2} AC$