
Triangles

• A triangle is a simple closed curve made up of three line segments.

It has three vertices, three sides and three angles.

- Triangles can be classified on the basis of their sides as:
 - 1. Scalene No side of the triangle is equal
 - 2. Isosceles Exactly two sides of the triangle are equal
 - 3. Equilateral All the sides of the triangle are equal \hat{I}
- On the basis of angles, triangles can be classified as:
 - 1. Acute-angled All the angles of the triangle are less than 90°
 - 2. Obtuse-angled Any one of the angles of the triangle is greater than 90°
 - 3. Right-angled Any one of the angles of the triangle is 90°
- Median of a triangle

A median is a line segment joining the vertex of a triangle to the mid-point of the opposite side.




In the given $\triangle ABC$, if AD = DC, then BD is the median of $\triangle ABC$ with respect to the side AC.


A triangle has three medians, one for each side.

• Altitude of a triangle

An altitude is the perpendicular drawn from the vertex of a triangle to its opposite side.




In the given figure, AD is the altitude of \triangle ABC with respect to side BC. A triangle has three altitudes, one from each vertex. The altitude of a triangle may or may not lie inside the triangle. For example, for \triangle PQR, its altitude lies outside it.



• Angle sum property of triangles:

The sum of all the three interior angles of a triangle is 180°.



Example:

If the measures of the angles of a triangle are in the ratio 2: 4: 6, then find all the angles of the triangle.

Solution:

Ratio of the measures of angles = 2: 4: 6

Therefore, let the angles of the triangle measure 2*x*, 4*x*, and 6*x*. Now, $2x + 4x + 6x = 180^{\circ}$ {By angle sum property of triangles} $\Rightarrow 12x = 180^{\circ}$ $\Rightarrow x = 15^{\circ}$ Thus, the angles of the triangle are $2x = 2 \times 15^{\circ} = 30^{\circ}$,

$$6x = 6 \times 15^\circ = 90^\circ.$$


 $4x = 3 \times 15^{\circ} = 60^{\circ}$

The measure of one of angle is 90°.


• Facts deduced from angle sum property of triangles:

There can be no triangle with two right angles or two obtuse angles. There can be no triangle with all angles less than or greater than 60°.

• Relation between the vertex angle and the angles made by the bisectors of the remaining angles:




• The angle formed by a side of a triangle with an extended adjacent side is called an **exterior angle of the triangle**.



It can be seen that in \triangle ABC, side CB is extended up to point D. This extended side forms an angle with side AB, i.e., \angle ABD. This angle lies exterior to the triangle. Hence, \angle ABD is an exterior angle of \triangle ABC.


• If a side of a triangle is produced, then the exterior angle so formed is equal to the sum of the two interior opposite angles.



This property is known as exterior angle property of a triangle.

Example:

Find the value of x in the following figure.



Solution:

 \angle QRS is an exterior angle of \triangle PQR. It is thus equal to the sum of its interior opposite angles.

 $\therefore \angle QRS = \angle QPR + \angle PQR$

 $\Rightarrow x = 65^{\circ} + 70^{\circ} = 135^{\circ}$


Thus, the value of x is 135°.

- Two exterior angles can be drawn at each vertex of triangle. The two angles thus drawn have an equal measure and are equal to the sum of the two opposite interior angles.
- Isosceles triangle

In an isosceles triangle, (1) two sides are of equal length and (2) the base angles opposite to the equal sides are equal.

Example:


In the following figure, AC = BC. Find m $\angle ACD$.



• Angles opposite to equal sides of a triangle are equal.

Example:

Find the missing angles in the following triangles.



Solution:

We know that angles opposite to equal sides of a triangle are equal.

 $\therefore \angle ABC = \angle ACB = x$ (say)

By angle sum property of triangles, we obtain

 $\angle ABC + \angle BCA + \angle CAB = 180^{\circ}$ $\Rightarrow x + x + 70^{\circ} = 180^{\circ}$ $\Rightarrow 2x = 110^{\circ}$ $\Rightarrow x = 55^{\circ}$ Thus, $\angle ABC = \angle BCA = 55^{\circ}$

• Sides opposite to equal angles of a triangle are equal in length.

Thus, we can say that if two angles of a triangle are equal then the sides opposite to them are also equal, therefore the triangle is isosceles.

- In a triangle, the sum of the lengths of any two sides is greater than the length of the third side.
- The difference between the lengths of any two sides of a triangle is less than the length of the third side.

Example:


Is it possible to construct a triangle with sides having lengths as 6 cm, 5 cm and 12 cm? Solution: In the given case, 12 cm - 5 cm = 7 cm > 6 cm

Here, the difference between two sides is greater than the third side.

Hence, a triangle with the given lengths cannot be constructed.

• If two sides of a triangle are unequal then the longer side has the greater angle opposite it. Thus, we can say that angle opposite to the shorter side of a triangle is smaller.

For example, in the given triangle, AC > AB, therefore $\angle ABC > \angle ACB$.



• If two angles of a triangle are unequal then the greater angle has the longer side opposite it. Thus, we can say that the smaller angle has the shorter side opposite it.

For example, in the given figure, $\angle BAC > \angle ACB$, therefore BC > AB.

