

# Redox Reactions

## Quick Revision

The reactions that involve oxidation and reduction as its two half reactions are called **redox reactions**.

### 1. Classical Idea of Redox Reactions

According to classical concepts, oxidation and reduction are defined as the process that involve:

#### Oxidation

- (i) Addition of oxygen
- (ii) Addition of electronegative element
- (iii) Removal of hydrogen
- (iv) Removal of electropositive element

#### Reduction

- (i) Removal of oxygen
- (ii) Removal of electronegative element
- (iii) Addition of hydrogen
- (iv) Addition of electropositive element

### 2. Oxidising and Reducing Agents

In a redox reaction, the substance which oxidises the other species or itself undergo reduction is called the oxidising agent. The substance that reduces the other species and itself undergo oxidation is called the reducing agent.

### 3. Redox Reactions in Term of Electron Transfer

Loss of electrons or an increase in oxidation state is called oxidation. Gain of electrons or a decrease in oxidation state is called reduction. Because of simultaneous loss and gain of electrons in oxidation-reduction processes, the redox reactions, (or the oxidation-reduction reactions) are also called electron transfer reactions.

### 4. Oxidation Number

It is defined as “the charge that an atom of the element possesses in its ion or appear to have when present in the combined state with other atoms.”

### 5. Rules for Calculating Oxidation Number

These rules are given below:

**Rule 1** The oxidation number of an atom in its free or elementary state or in any of its allotropes is zero. e.g. The oxidation state of H in  $H_2$ , S in  $S_8$ , P in  $P_4$ .

**Rule 2** In case of ions having only one kind of atoms, the oxidation number of each atom is equal to charge present on the ion. e.g. In case of  $Na^+$ ,  $Mg^{2+}$ ,  $Fe^{3+}$ ,  $Cl^-$  and  $O^{2-}$ , the oxidation state is respectively +1, +2, +3, -1, -2.

**Rule 3** The oxidation state of alkali metals in all their compounds is always +1. Similarly, in case of alkaline earth metals, it is always +2. For aluminium, oxidation state is always +3.

**Rule 4** The oxidation state of oxygen in most of its compounds is -2, with an exception of peroxides and superoxides in which the oxidation state of oxygen is respectively -1 and -1/2.

**Rule 5** The oxidation state of hydrogen is generally +1 with an exception of metallic hydrides like  $NaH$ ,  $CaH_2$  etc. In these hydrides, oxidation state of hydrogen is -1.


**Rule 6** The oxidation state of fluorine in all of its compounds is always  $-1$ . Other halogens (i.e. chlorine, bromine and iodine) also exhibit  $-1$  oxidation state but it is not always true.

In case of oxoacids and oxoanions, halogens (except fluorine) exhibit positive oxidation state.

**Rule 7** The algebraic sum of the oxidation numbers of all the atoms present in a compound must be equal to zero.

**Rule 8** In case of polyatomic ions, the algebraic sum of oxidation number of all the atoms present in the ion must be equal to the charge on the ion. e.g. In case of carbonate ion ( $\text{CO}_3^{2-}$ ), it is equal to  $-2$ .

### 6. Paradox of Fractional Oxidation Number




two carbon atoms are present in  $+2$  oxidation state each whereas third one is present in zero oxidation state.


The average of oxidation states of 3 C-atoms in

$$\text{C}_3\text{O}_2 = \frac{2 + 2 + 0}{3} = \frac{4}{3}$$

- Average oxidation state of Br in  $\text{Br}_3\text{O}_8$  is  $16/3$  while the oxidation states of three Br atoms are  $+6$ ,  $+4$  and  $+6$  as shown below:



- The average oxidation state of four S-atoms in  $\text{S}_4\text{O}_6^{2-}$  is  $2.5$  while the actual oxidation state of the four S-atoms are  $+5$ ,  $0$ ,  $0$  and  $+5$  as shown below:



### 7. Balancing of Redox Reactions

(i) **Ion electron method** The method involves the following steps:

- Write redox reaction in ionic form.
- Split redox reaction into oxidation half and reduction half reactions. Balance atoms of each half reactions by using simple multiples.

For balancing H and O, add  $\text{H}^+$  ion and  $\text{H}_2\text{O}$  to the appropriate sides, similarly add  $\text{OH}^-$  and  $\text{H}_2\text{O}$  to the appropriate sides.

Balance the charge on both sides and multiply one or both half reactions by suitable number to equalise number of electrons in both equations. Add the two balance half reactions and cancel common terms.

(ii) **Oxidation number method** The method involves the following steps:

- Assign oxidation number to the atoms in the equation and write separate equations for atoms undergoing oxidation and reduction.
- Find the change in oxidation number in each equation and make the change equal in both the equations by multiplying with suitable integers.

After adding both the equations complete the balancing (by balancing H and O).

# Objective Questions

## Multiple Choose Questions

1. Which of the following reactions represent(s) redox process?

- (a) Electrochemical process for extraction of highly reactive metals and non-metals
- (b) Manufacturing of caustic soda
- (c) Corrosion of metals
- (d) All of the above


2. Which of the following processes take place in oxidation?

- (a) Addition of oxygen
- (b) Addition of hydrogen
- (c) Removal of oxygen
- (d) Removal of chlorine

3. Which of the following processes take place in reduction?

- (a) Removal of oxygen
- (b) Addition of hydrogen
- (c) Removal of hydrogen
- (d) Both (a) and (b)

4. In the given reaction,



ferric chloride undergoes +2HCl(aq)



- (a) reduction process
- (b) oxidation process
- (c) addition process
- (d) All of the above

5. In the reaction given below, identify the species undergoing oxidation and reduction, respectively



- (a)  $\text{H}_2\text{S}$  is oxidised and  $\text{Cl}_2$  is reduced
- (b)  $\text{H}_2\text{S}$  is reduced and  $\text{Cl}_2$  is oxidised
- (c) Both  $\text{H}_2\text{S}$  and  $\text{Cl}_2$  are oxidised
- (d) Both  $\text{H}_2\text{S}$  and  $\text{Cl}_2$  are reduced


6. In the given reaction,



Which of the following processes takes place?

- (a) Oxidation due to removal of potassium
- (b) Oxidation due to removal of iron
- (c) Reduction due to removal of potassium
- (d) Oxidation due to removal of electronegative element


7. In the following reaction,



the process taking place w.r.t. Mg is known as

- (a) oxidation
- (b) reduction
- (c) redox reaction
- (d) None of these

8. In the given reaction,




- (a) oxidation of ethylene
- (b) reduction of ethylene
- (c) Both (a) and (b)
- (d) None of the above

9. In oxidation process,

- (a) oxidation number decreases
- (b) number of electrons decreases
- (c) oxygen content decreases
- (d) number of ions decreases

10. Given the reaction for the discharge of a cobalt-cadmium battery




Which species is oxidised during the discharge of the battery?

- (a)  $\text{Co}^{3+}$
- (b)  $\text{Co}^{2+}$
- (c) Cd
- (d)  $\text{Cd}^{2+}$

11. Both oxidation and reduction takes place in

- (a)  $\text{NaBr} + \text{HCl} \longrightarrow \text{NaCl} + \text{HBr}$
- (b)  $\text{HBr} + \text{AgNO}_3 \longrightarrow \text{AgBr} + \text{HNO}_3$
- (c)  $\text{H}_2 + \text{Br}_2 \longrightarrow 2\text{HBr}$
- (d)  $\text{CaO} + \text{H}_2\text{SO}_4 \longrightarrow \text{CaSO}_4 + \text{H}_2\text{O}$

12. In the reaction,




the elements which have been oxidised and reduced respectively are

- (a) chlorine and oxygen  
 (b) oxygen and chlorine  
 (c) potassium and oxygen  
 (d) oxygen and potassium

13. The compound that can work both as an oxidising and reducing agent is

- (a)  $\text{KMnO}_4$  (b)  $\text{H}_2\text{O}_2$   
 (c)  $\text{Fe}_2(\text{SO}_4)_3$  (d)  $\text{K}_2\text{Cr}_2\text{O}_7$

14. Water molecule is formed by the reaction,



What does happen in this reaction?

- (a) Electrons are transferred from H to O-atom  
 (b) Electrons are transferred from O to H-atom  
 (c) Electrons are accepted by H from O-atom  
 (d) Electrons are donated by O to H-atom

15. Which of the following is not an example of redox reaction? (NCERT Exemplar)

- (a)  $\text{CuO} + \text{H}_2 \longrightarrow \text{Cu} + \text{H}_2\text{O}$   
 (b)  $\text{Fe}_2\text{O}_3 + 3\text{CO} \longrightarrow 2\text{Fe} + 3\text{CO}_2$   
 (c)  $2\text{K} + \text{F}_2 \longrightarrow 2\text{KF}$   
 (d)  $\text{BaCl}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{BaSO}_4 + 2\text{HCl}$

16. Match the Column I (Reaction with underlined species) with Column II (Type of change shown by underlined species) and choose the correct option from the codes given below.

Column I (Reactions)	Column II (Type of change)
A. $2\underline{\text{Mg}} + \text{O}_2 \longrightarrow 2\text{MgO}$	1. Removal of hydrogen.
B. $\underline{\text{Mg}} + \text{Cl}_2 \longrightarrow \text{MgCl}_2$	2. Removal of electropositive element.
C. $2\underline{\text{H}_2\text{S}} + \text{O}_2 \longrightarrow 2\text{S} + 2\text{H}_2\text{O}$	3. Addition of oxygen.
D. $2\underline{\text{KI}} + \text{H}_2\text{O} + \text{O}_3 \longrightarrow 2\text{KOH} + \text{I}_2 + \text{O}_2$	4. Addition of electronegative element.

Codes

A	B	C	D	A	B	C	D
(a) 2	3	4	1	(b) 3	4	1	2
(c) 3	4	2	1	(d) 3	2	1	4

17. Strongest reducing agent is

- (a) K (b) Mg (c) Al (d) Ba

18. In the reaction,




sodium acts as a/an

- (a) oxidising agent (b) reducing agent  
 (c) complexing agent (d) None of these

19. Which reaction indicates the action of  $\text{HNO}_3$  as oxidising agent?


- (a)  $\text{NaOH} + \text{HNO}_3 \longrightarrow \text{NaNO}_3 + \text{H}_2\text{O}$   
 (b)  $\text{Ca}(\text{OH})_2 + 2\text{HNO}_3 \longrightarrow \text{Ca}(\text{NO}_3)_2 + 2\text{H}_2\text{O}$   
 (c)  $\text{C}_6\text{H}_6 + \text{HNO}_3 \longrightarrow \text{C}_6\text{H}_5\text{NO}_2 + \text{H}_2\text{O}$   
 (d)  $\text{NaCl} + \text{HNO}_3 \longrightarrow \text{HCl} + \text{NaNO}_3$

20. In the following reaction reducing agent is



- (a)  $\text{H}^+$  (b)  $\text{Cr}_2\text{O}_7^{2-}$  (c)  $\text{H}_2\text{O}$  (d) Ni

21. Which is the best description behaviour of bromine in the given equation?



- (a) Proton acceptor  
 (b) Both oxidised and reduced  
 (c) Oxidised  
 (d) Reduced

22. In which of the following compounds, nitrogen exhibits highest oxidation state?

- (a)  $\text{N}_2\text{H}_4$  (b)  $\text{NH}_3$   
 (c)  $\text{N}_3\text{H}$  (d)  $\text{NH}_2\text{OH}$

23. Oxidation states of X, Y, Z are +2, +5 and -2 respectively. The formula of the compound formed by these will be

- (a)  $\text{X}_2\text{YZ}_6$  (b)  $\text{XY}_2\text{Z}_6$   
 (c)  $\text{XY}_5$  (d)  $\text{X}_3\text{YZ}_4$

24. In which of the following, Fe exhibits minimum oxidation state?

- (a)  $K_4Fe(CN)_6$   
 (b)  $Fe_3O_4$   
 (c)  $Fe(CO)_5$   
 (d)  $FeSO_4(NH_4)_2SO_4 \cdot 6H_2O$

25. Which of the following have been arranged in the decreasing order of oxidation number of sulphur?

- (a)  $Na_2S_4O_6 > H_2S_2O_7 > Na_2S_2O_3 > S_8$   
 (b)  $H_2SO_4 > SO_2 > H_2S > H_2S_2O_8$   
 (c)  $SO_2^{2-} > SO_4^{2-} > SO_3^{2-} > HSO_4^-$   
 (d)  $H_2SO_5 > H_2SO_3 > SCl_2 > H_2S$

26. In which of the following compounds, an element exhibits two different oxidation states?

- (a)  $NH_2OH$  (b)  $NH_4NO_3$   
 (c)  $N_2H_4$  (d)  $N_3H$

27. State the oxidation number of carbonyl carbon in methanal and methanoic acid respectively.

- (a) 0 and 0 (b) 0 and +2  
 (c) +1 and +2 (d) +1 and +3

28. +3 oxidation state is most common in ...

- (a) Ni(28) (b) Fe(26)  
 (c) Zn(30) (d) Cu(29)

29. In the reaction,  $4Na + O_2 \longrightarrow 2Na_2O$ , sodium acts as a/an

- (a) oxidising agent (b) reducing agent  
 (c) complexing agent (d) None of these

30. The value of oxidation numbers of Cl, in  $Cl_2$ ,  $NaOCl$  and  $ClO_3^-$  are respectively.

- (a) +2, 0, +5 (b) 0, +2, +5  
 (c) +2, +3, +5 (d) 0, +1, +5

31. What is the average oxidation number of carbon in carbon suboxide?

- (a)  $+\frac{4}{3}$  (b)  $+\frac{10}{4}$  (c) +2 (d)  $+\frac{2}{3}$

32. The oxidation states of sulphur in  $H_2SO_4$ ,  $HSO_3^-$  and  $SO_2Cl_2$  respectively are

- (a) +6, +4, +6 (b) +6, +6, +4  
 (c) +6, -6, +4 (d) -4, +6, +6

33. The largest oxidation number exhibited by an element depends on its outer electronic configuration. With which of the following outer electronic configurations the element will exhibit largest oxidation number?

(NCERT Exemplar)

- (a)  $3d^1 4s^2$  (b)  $3d^3 4s^2$   
 (c)  $3d^5 4s^1$  (d)  $3d^5 4s^2$

34. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is  $1s^2 2s^2 2p^3$ , the simplest formula for this compound is

- (a)  $Mg_2X$  (b)  $MgX_2$   
 (c)  $Mg_2X_3$  (d)  $Mg_3X_2$

35. The valency of Cr in  $[Cr(H_2O)_4Cl_2]^+$  ion is

- (a) 3 (b) 1  
 (c) 6 (d) 5

36. Match the Column I with Column II and select the correct option for oxidation number of N-atom from the codes given below.

	Column I (Compounds)	Column II (Oxidation number)	
A.	$NH_2OH$	1.	-1
B.	$Mg_3N_2$	2.	-1
C.	$N_2O$	3.	+5
D.	$N_2O_5$	4.	-3

Codes

- A B C D  
 (a) 1 3 4 2  
 (b) 2 4 3 1  
 (c) 2 4 1 3  
 (d) 4 2 1 3

**37.** In oxygen difluoride ( $\text{OF}_2$ ) and dioxygen difluoride ( $\text{O}_2\text{F}_2$ ), the oxygen is assigned an oxidation number of

- (a) +1 and +2 respectively  
 (b) +2 and +2 respectively  
 (c) +1 and +1 respectively  
 (d) +2 and +1 respectively


**38.** Which of the following arrangements represent increasing oxidation number of the central atom? (NCERT Exemplar)

- (a)  $\text{CrO}_2^-$ ,  $\text{ClO}_3^-$ ,  $\text{CrO}_4^{2-}$ ,  $\text{MnO}_4^-$   
 (b)  $\text{ClO}_3^-$ ,  $\text{CrO}_4^{2-}$ ,  $\text{MnO}_4^-$ ,  $\text{CrO}_2^-$   
 (c)  $\text{CrO}_2^-$ ,  $\text{ClO}_3^-$ ,  $\text{MnO}_4^-$ ,  $\text{CrO}_4^{2-}$   
 (d)  $\text{CrO}_4^{2-}$ ,  $\text{MnO}_4^-$ ,  $\text{CrO}_2^-$ ,  $\text{ClO}_3^-$

**39.** Which of the following reactions is represented in basic medium?

- (a)  $\text{MnO}_4^-(\text{aq}) + \text{SO}_2(\text{g}) \longrightarrow \text{Mn}^{2+}(\text{aq}) + \text{HSO}_4^-(\text{aq})$   
 (b)  $\text{H}_2\text{O}_2(\text{aq}) + \text{Fe}^{2+}(\text{aq}) \longrightarrow \text{Fe}^{3+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$   
 (c)  $\text{MnO}_4^-(\text{aq}) + \text{I}^-(\text{aq}) \longrightarrow \text{MnO}_2(\text{s}) + \text{I}_2(\text{s})$   
 (d)  $\text{Cr}_2\text{O}_7^{2-} + \text{SO}_2(\text{g}) \longrightarrow \text{Cr}^{3+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq})$

**40.** In the following redox reaction,



the values of coefficients  $x$ ,  $y$  and  $z$  respectively, are

- (a) 3, 8, 2                      (b) 3, 8, 7  
 (c) 3, 2, 4                      (d) 3, 1, 8

**41.** A mixture of potassium chlorate, oxalic acid and sulphuric acid is heated. During the reaction which element undergoes maximum change in oxidation number?

- (a) S                      (b) H                      (c) Cl                      (d) C

### Assertion-Reasoning MCQs

**Directions** In the following questions (Q.No. 42-55) a statement of Assertion followed by a statement of Reason is given. Choose the correct answer out of the following choices.

- (a) Both Assertion and Reason are correct statements and Reason is the correct explanation of the Assertion.  
 (b) Both Assertion and Reason are correct statements, but Reason is not the correct explanation of the Assertion.  
 (c) Assertion is correct, but Reason is incorrect statement.  
 (d) Assertion is incorrect but Reason is correct statement.

**42. Assertion**  $\text{KMnO}_4$  is a stronger oxidising agent than  $\text{K}_2\text{Cr}_2\text{O}_7$ .

**Reason** This is due to the increasing stability of the lower species to which they are reduced.

**43. Assertion** In the reaction between potassium permanganate and potassium iodide, permanganate ions act as oxidising agent.

**Reason** Oxidation state of manganese changes from +2 to +7 during the reaction.

**44. Assertion** Average oxidation number of I in  $\text{KI}_3$  is  $-1/3$ .

**Reason**  $\text{KI}_3$  is made up of KI and  $\text{I}_2$ . Each species have different oxidation number.

**45. Assertion**  $\text{PbCl}_4$  is a powerful oxidising agent.

**Reason**  $\text{PbCl}_4$  is more stable than  $\text{PbCl}_2$ .

**46. Assertion** In some cases oxygen shows positive oxidation number though it is an electronegative element.

**Reason** Fluorine is more electronegative than oxygen.

**47. Assertion**  $\text{H}_2\text{SO}_4$  cannot act as a reducing agent.

**Reason** Sulphur cannot increase its oxidation number beyond +6.

**48. Assertion** Among halogens fluorine is the best oxidant.

**Reason** Fluorine is the most electronegative atom. (NCERT Exemplar)

- 49. Assertion** The two Fe atoms in  $\text{Fe}_3\text{O}_4$  have different oxidation numbers.

**Reason**  $\text{Fe}^{2+}$  ions decolourise  $\text{KMnO}_4$  solution.

- 50. Assertion** In the species,  $\text{Br}_3\text{O}_8$  each of two extreme bromine exhibits oxidation state of +6 and the middle bromine of +4.

**Reason** The average of three oxidation numbers of bromine of the  $\text{Br}_3\text{O}_8$  is 16/3.

- 51. Assertion** Amongst the halogens, fluorine cannot oxidise the element to the highest oxidation state.

**Reason** Due to small size of fluoride ion, it is difficult to oxidise fluoride ion to fluorine. Hence, reverse reaction takes place more easily.

- 52. Assertion** The oxidation number of O in  $\text{O}_3$  is zero and the oxidation number of S in  $\text{SO}_3$  is +4.

**Reason**  $\text{O}_3$  can act as an oxidising agent as well as a reducing agent but  $\text{SO}_2$  can act only as an oxidising agent.

- 53. Assertion** The formal oxidation number of sulphur in  $\text{Na}_2\text{S}_4\text{O}_6$  is 2.5.

**Reason** Two S-atoms are not directly linked with O-atoms.

- 54. Assertion**  $\text{MnO}_4^-$  is reduced to  $\text{Mn}^{2+}$  in acidic medium.

**Reason** In acidic medium, following reaction takes place.



- 55. Assertion**  $\text{Cl}_2$  gas bleaches the articles permanently.

**Reason**  $\text{Cl}_2$  is a strong reducing agent.

## Case Based MCQs

- 56.** Read the passage given below and answer the following questions :

The oxidation state of an individual atom is 0. The total oxidation state of all atoms in a neutral species is 0 and in an ion is equal to the ion charge. Group 1 metals have an oxidation state of +1 and group 2 an oxidation state of +2.

The oxidation state of fluorine is -1 in compounds. Hydrogen generally has an oxidation state of +1 in compounds. Oxygen generally has an oxidation state of -2 in compounds.

In binary metal compounds, group 17 elements have an oxidation state of -1, group 16 elements of -2, and group 15 elements of -3. The sum of the oxidation states is equal to zero for neutral compounds and equal to the charge for polyatomic ion species. An atom is oxidised if its oxidation number increases and an atom is reduced if its oxidation number decreases.

The atom that is oxidised is the reducing agent and the atom that is reduced is the oxidising agent.

(Note the oxidising and reducing agents can be the same element or compound).

Redox reactions are comprised of two parts, a reduced half and an oxidised half, that always occur together.

The reduced half gains electrons and the oxidation number decreases, while the oxidised half loses electrons and the oxidation number increases.

The ion or molecule that accepts electrons is called the oxidising agent, by accepting electrons it causes the oxidation of another species. Conversely, the species that donates electrons is called the **reducing agent**; when the reaction occurs, it reduces the other species.

The following questions (i-iv) are multiple choice questions. Choose the most appropriate answer :

- (i) One mole of acidified  $K_2Cr_2O_7$  on reaction with excess KI will liberate  $n$  mole of  $I_2$ , then the value of  $n$  is  
 (a) 6 (b) 1  
 (c) 3 (d) 7
- (ii) When electrons are transferred from Zn to  $Cu^{2+}$  in copper sulphate solution, the energy (heat) is  
 (a) absorbed  
 (b) evolved  
 (c) consumed  
 (d) Both (a) and (b)
- (iii) Negative  $E^\ominus$  indicates that redox couple is  
 (a) weaker reducing agent than  $H^+ / H_2$  couple  
 (b) stronger reducing agent than  $H^+ / H_2$  couple  
 (c) stronger oxidising agent than  $H^+ / H_2$  couple  
 (d) weaker oxidising agent than  $H^+ / H_2$  couple
- (iv) Which of the following statements is/are incorrect?  
 (a) The reactants, which undergo oxidation and reduction are called reductant and oxidant respectively  
 (b) In redox reaction, the oxidation number of oxidant increases, while that of reductant decreases  
 (c)  $HNO_2$  acts as an oxidising as well as reducing agent  
 (d) Oxidation is the process, in which electrons are lost

Or

In alkaline medium,  $ClO_2$  oxidises  $H_2O_2$  to  $O_2$  and itself gets reduced to  $Cl^-$ . How many moles of  $H_2O_2$  are oxidised by 1 mole of  $ClO_2$ ?

- (a) 1 (b) 1.5  
 (c) 2.5 (d) 5

57. Read the passage given below and answer the following questions :

The concept of electron transfer is found unable to explain the redox changes or electron shift in case of covalent compounds.

To explain these changes a new concept, called oxidation number is introduced. Oxidation number is defined as the charge that an atom of the element has in its ion or appear to have when present in the combined state with other atoms.

In other words, it is also defined as the charge that an atom appear to have in a compound when all other atoms are removed as ions from the compound.

The following steps are involved while calculating the oxidation number of an atom in a given compound/ion.

**Step I** Write down the formula of given compound/ion leaving some space between the atoms.

**Step II** Write the oxidation state of each element above its atoms. Write down  $x$  above the atom, oxidation state of which we have to find out.

**Step III** Multiply the oxidation numbers of each element with the number of atoms of that element present in the compound.

Enclose the product in a bracket.

**Step IV** Equate the algebraic sum of the oxidation numbers of all the atoms present in a compound to zero or to the charge in case of ionic species charge on the ion.

**Step V** Solve the equation obtained for the value of  $x$ .

The following questions (i-iv) are multiple choice questions. Choose the most appropriate answer:

- (i) Highest oxidation state of Mn is present in  
 (a)  $KMnO_4$  (b)  $K_2MnO_4$   
 (c)  $Mn_2O_3$  (d)  $MnO_2$
- (ii) Identify the element which never has positive oxidation number in any of its compound?  
 (a) Oxygen  
 (b) Chlorine  
 (c) Fluorine  
 (d) Bromine

(iii) When a manganous salt is fused with a mixture of  $\text{KNO}_3$  and solid  $\text{NaOH}$ , the oxidation number of  $\text{Mn}$  changes, from + 2 to

- (a) + 4                      (b) + 3  
(c) + 6                      (d) + 7

(iv) The brown ring complex compound is formulated as  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]\text{SO}_4$ . What will be the oxidation state of iron in the given complex?

- (a) + 2                      (b) + 3  
(c) + 4                      (d) + 1


*Or*

In which of the following reactions, there is no change in valency?

- (a)  $\text{SO}_2 + 2\text{H}_2\text{S} \longrightarrow 2\text{H}_2\text{O} + 3\text{S}$   
(b)  $2\text{Na} + \text{O}_2 \longrightarrow 2\text{Na}_2\text{O}_2$   
(c)  $\text{Na}_2\text{O}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O}_2$   
(d)  $4\text{KClO}_3 \longrightarrow 3\text{KClO}_4 + \text{KCl}$

**58.** Read the passage given below and answer the following questions :

In a redox reaction, the substance which oxidises the other or which itself undergoes reduction is called the oxidising agent and the substance that reduces the other and itself undergoes oxidation is called the reducing agent. e.g.




Thus, in the above example,  $\text{HgCl}_2$  is an oxidising agent (as it reduces) and  $\text{SnCl}_2$  is a reducing agent (as it oxidises).

In these questions (i-iv) a statement of Assertion followed by a statement of Reason is given. Choose the correct answer out of the following choices :

- (a) Assertion and Reason both are correct statements and Reason is correct explanation for Assertion.  
(b) Assertion and Reason both are correct statements but Reason is not correct explanation for Assertion.  
(c) Assertion is correct statement but Reason is incorrect statement.  
(d) Assertion is incorrect statement but Reason is correct statement.

(i) **Assertion** This reaction is redox reaction,



**Reason** Both oxidation and reduction take place in the given reaction.

(ii) **Assertion** The decomposition of hydrogen peroxide to form water and oxygen is an example of disproportionation reaction.

**Reason** The oxygen of peroxide is in  $-1$  oxidation state and it is converted to zero oxidation state in  $\text{O}_2$  and  $-2$  oxidation state in  $\text{H}_2\text{O}$ .

(iii) **Assertion** In the species,  $\text{Br}_3\text{O}_8$  each of two extreme bromine exhibits oxidation state of + 6 and the middle bromine of + 4.

**Reason** The average of three oxidation numbers of bromine of the  $\text{Br}_3\text{O}_8$  is  $16/3$ .

(iv) **Assertion** In the reaction between potassium permanganate and potassium iodide, permanganate ions act as oxidising agent.

**Reason** Oxidation state of manganese changes from +2 to +7 during the reaction.

*Or*

**Assertion** A negative value of  $E^\circ$  means that the redox couple is a weaker reducing agent than the  $\text{H}^+/\text{H}_2$  couple.

**Reason** A negative  $E^\circ$  means that the redox couple is stronger reducing agent than the  $\text{H}^+/\text{H}_2$ .

**59.** Read the passage given below and answer the following questions :

The real or imaginary charge which an atom appears to have in its combined state is called oxidation state or oxidation number of that atom. Fraction oxidation states are often used to represent the average oxidation states of several atom in

a structure. These oxidation states are very helpful in finding the oxidation and reduction process in redox reactions. Redox reactions are of two main, i.e. intermolecular redox reactions and intramolecular redox reactions.

The elements that show an increase in oxidation number (hydrogen and chlorine in the above reaction) are **oxidised**, while the elements that are reduced (oxygen and chlorine in the above reaction) show a decrease in their oxidation numbers from their initial values.

In these questions (i-iv) a statement of Assertion followed by a statement of Reason is given. Choose the correct answer out of the following choices :

- (i) **Assertion** Oxidation state of nitrogen in  $N_3H$  is  $-\frac{1}{3}$ .

**Reason** Nitrogen is less electronegative than hydrogen.

- (ii) **Assertion** The decomposition of hydrogen peroxide to form water and oxygen is an example of disproportionation reaction.

**Reason** The oxygen of peroxide is in  $-1$  oxidation state and it is converted to zero oxidation state in  $O_2$  and  $-2$  oxidation state in  $H_2O$ .

- (iii) **Assertion** The electrons are transferred from zinc to copper through the wire which connects the two rods.

**Reason** Electricity flows through the salt-bridge by migration of ions from one beaker to other.

- (iv) **Assertion** Redox couple is the combination of oxidised and reduced, form of a substance involved in an oxidation or reduction half-cell.

**Reason** In the representation  $E_{Fe^{3+}/Fe^{2+}}^{\ominus}$  and  $E_{Cu^{2+}/Cu}^{\ominus}$ ,  $Fe^{3+}/Fe^{2+}$  and  $Cu^{2+}/Cu$  are redox couples.

Or

**Assertion** Oxidation number of hydrogen  $-1$  in  $CaH_2$ .

**Reasons**  $CaH_2$  is a metal hydrides and for hydrides, hydrogen is assigned the oxidation number of  $-1$ .

## ANSWERS

### Multiple Choice Questions

1. (d) 2. (a) 3. (d) 4. (a) 5. (a) 6. (a) 7. (a) 8. (b) 9. (b) 10. (c)  
 11. (c) 12. (b) 13. (b) 14. (a) 15. (d) 16. (b) 17. (a) 18. (b) 19. (c) 20. (d)  
 21. (b) 22. (c) 23. (b) 24. (c) 25. (d) 26. (b) 27. (b) 28. (b) 29. (b) 30. (d)  
 31. (a) 32. (a) 33. (d) 34. (d) 35. (a) 36. (c) 37. (d) 38. (a) 39. (c) 40. (a)  
 41. (c)

### Assertion-Reasoning MCQs

42. (a) 43. (c) 44. (a) 45. (c) 46. (a) 47. (a) 48. (b) 49. (b) 50. (b) 51. (b)  
 52. (c) 53. (c) 54. (d) 55. (c)

### Case Based MCQs

56. (i)-(c), (ii)-(b), (iii)-(b), (iv)-(b) or-(c) 57. (i)-(a), (ii)-(c), (iii)-(c), (iv)-(b) or-(c)  
 58. (i)-(a), (ii)-(a), (iii)-(b), (iv)-(c) or-(d) 59. (i)-(c), (ii)-(a), (iii)-(b), (iv)-(b) or-(b)


# EXPLANATIONS

1. Electrochemical processes for the extraction of highly reactive metals and non-metals, manufacturing of chemical compounds like caustic soda, operation of dry and wet batteries and corrosion of metals fall within the range of redox processes.

2. Addition of oxygen/electronegative element and removal of hydrogen/electropositive element takes place in oxidation.

3. Reduction is a process which involves addition of hydrogen or electropositive elements to a substance or removal of oxygen or electronegative element from a substance.

4. In the given reaction, removal of electronegative element, i.e. chlorine from ferric chloride takes place. Hence, it is an example of reduction process.



Removal of chlorine  
(electronegative element)




Addition of hydrogen  
(reduction)

Removal of hydrogen(oxidation)

Thus,  $\text{H}_2\text{S}$  is oxidised and  $\text{Cl}_2$  is reduced.


6. In the given reaction,




Removal of potassium (electropositive element)

So, here oxidation takes place due to removal of one potassium atom.

7. Oxidation is a process, which involves addition of oxygen/electronegative element to a substance or removal of hydrogen/electropositive element from a substance.




Addition of oxygen = Oxidation



(Addition of hydrogen)


Reduction of ethylene occurs due to the addition of hydrogen.

9. In oxidation process, oxidation number increases and number of electrons decreases.




Here, oxidation number of Cd increases from 0 to +2, hence Cd is oxidised.

11. Both oxidation and reduction are taking place in the following reaction,



$\text{H}_2$ -reducing agent;  $\text{Br}_2$ -oxidising agent.


In other options, neither oxidation nor reduction takes place because oxidation number of elements involved in reaction remain same.



13. Oxygen in  $\text{H}_2\text{O}_2$  has oxidation number  $-1$  which can increase or decrease.

Hence, it can act as both oxidising agent or reducing agent.




14. Hydrogen is oxidised by loss of 1 electron



Oxygen is reduced by gain of 2 electrons

In this reaction, hydrogen (H) has transferred electrons to oxygen (O).


15. Following are the examples of redox reaction :




Option (d) is not an example of redox reaction.

16. The correct match is  
A → 3, B → 4, C → 1, D → 2


Oxidation (addition of oxygen)



Oxidation (addition of electronegative element)



Reduction (removal of hydrogen)




Oxidation (removal of electropositive element)

17. Since, K contains only one electron in its outermost shell, it has higher tendency to donate it, i.e. has higher tendency to get oxidised. Therefore, it is the strongest reducing agent among the given elements.



Loss of  $e^-$  (oxidation)

In the above reaction, Na converts into  $(\text{Na}^+)$  ion, i.e. Na donates its electron to oxygen atom. So, it behaves as reducing agent.




Reduction (gain of  $e^-$ )

In this reaction,  $\text{HNO}_3$  behaves as an oxidising agent while in rest of the reactions such as,


- (i)  $\text{NaOH} + \text{HNO}_3 \longrightarrow \text{NaNO}_3 + \text{H}_2\text{O}$   
(ii)  $\text{Ca}(\text{OH})_2 + 2\text{HNO}_3 \longrightarrow \text{Ca}(\text{NO}_3)_2 + 2\text{H}_2\text{O}$   
(iii)  $\text{NaCl} + \text{HNO}_3 \longrightarrow \text{HCl} + \text{NaNO}_3$

$\text{HNO}_3$  neither behaves as oxidising agent nor as reducing agent.

20.



As in this reaction Ni is oxidised to  $\text{Ni}^{2+}$  and reduced to  $\text{Cr}_2\text{O}_7^{2-}$ , thus it acts as reducing agent.



Here, oxidation number of bromine increases as well as decreases, i.e. bromine is oxidised as well as reduced.

22. Let, the oxidation state of nitrogen in the given compounds be  $x$ .

(a)  $\overset{x+1}{\text{N}_2}\text{H}_4, 2x + (+1)4 = 0$   
 $2x = -4$   
 $x = -2$

(b)  $\overset{x+1}{\text{N}}\text{H}_3, x + (+1)3 = 0$   
 $x = -3$

(c)  $\overset{x+1}{\text{N}_3}\text{H}, 3x + (+1) = 0$   
 $3x = -1$   
 $x = -1/3$

(d)  $\overset{x+1-2+1}{\text{N}}\text{H}_2\text{OH},$   
 $x + (+1)2 + (-2) + (+1) = 0$   
 $x + 2 - 2 + 1 = 0$   
 $x + 1 = 0 \Rightarrow x = -1$

The oxidation state of nitrogen is highest in  $\text{N}_3\text{H}$ .

23. We know that, the algebraic sum of the oxidation states is always zero in neutral compound.

Oxidation states of  $X = +2$   
 $Y = +5$   
 $Z = -2$

So, the algebraic sum of total  $X, Y$  and  $Z$  should be equal to zero which is found in  $\text{XY}_2\text{Z}_6$ .

$$\text{XY}_2\text{Z}_6 = +2 + (5 \times 2) + (-2 \times 6)$$

$$= +2 + 10 - 12 = 0$$

24. (a)  $\text{K}_4\text{Fe}(\text{CN})_6 \Rightarrow 1 \times 4 + x + (-1)6 \Rightarrow 0; \therefore x = 2$   
(b)  $\text{Fe}_3\text{O}_4 \Rightarrow \text{FeO} \cdot \text{Fe}_2\text{O}_3 \Rightarrow +2 + 3$   
(c)  $\text{Fe}(\text{CO})_5 \Rightarrow x + 0(5) = 0; \therefore x = 0$   
(d)  $\text{FeSO}_4(\text{NH}_4)_2 \cdot \text{SO}_4 \cdot 6\text{H}_2\text{O} \Rightarrow x + (-2) + 0 + 0 = 0$   
 $\therefore x = +2$

Thus, in  $\text{Fe}(\text{CO})_5$ , Fe shows minimum oxidation number.

25. (a)  $\overset{2.5}{\text{Na}}_2\overset{+6}{\text{S}}_4\overset{+2}{\text{O}}_6; \overset{+6}{\text{H}}_2\overset{+4}{\text{S}}_2\overset{-2}{\text{O}}_7; \overset{+2}{\text{Na}}_2\overset{+6}{\text{S}}_2\overset{+2}{\text{O}}_3; \overset{0}{\text{S}}_8$   
(b)  $\overset{+6}{\text{H}}_2\overset{+4}{\text{S}}\overset{-2}{\text{O}}_4; \overset{-2}{\text{S}}\overset{+6}{\text{O}}_2; \overset{+6}{\text{H}}_2\overset{+4}{\text{S}}\overset{-2}{\text{O}}_8$   
(c)  $\overset{+2}{\text{S}}\overset{+6}{\text{O}}_2; \overset{+4}{\text{S}}\overset{+6}{\text{O}}_4; \overset{-2}{\text{S}}\overset{+6}{\text{O}}_3^-; \overset{+6}{\text{H}}\overset{+4}{\text{S}}\overset{-2}{\text{O}}_4^-$   
(d)  $\overset{+6}{\text{H}}_2\overset{+6}{\text{S}}\overset{+4}{\text{O}}_5; \overset{+4}{\text{H}}_2\overset{+6}{\text{S}}\overset{-2}{\text{O}}_3; \overset{+2}{\text{S}}\overset{-2}{\text{Cl}}_2; \overset{-2}{\text{H}}_2\overset{+6}{\text{S}}$

Thus, in option (d), compounds are arranged in the decreasing order of oxidation number of sulphur.

26.  $\text{NH}_4\text{NO}_3$  is actually  $\text{NH}_4^+$  and  $\text{NO}_3^-$ . It is an ionic compound.

The oxidation number of nitrogen in the two species is different as shown below:

In  $\text{NH}_4^+$ ,

$$\Rightarrow x + (4 \times 1) = +1 \text{ or } x + 4 = +1$$

$$\text{or } x = -3$$

Let, oxidation number of N in  $\text{NO}_3^-$  is  $x$

$$\Rightarrow x + (3 \times -2) = -1 \text{ or } x - 6 = -1 \text{ or } x = +1$$

27. Let the oxidation number of carbonyl carbon in methanal ( $\text{HCHO}$ ) and methanoic acid ( $\text{HCOOH}$ ) is  $x$  and  $y$  respectively.

In  $\text{HCHO}$ ,  $2(+1) + x + (-2) = 0$

$$2 + x - 2 = 0 \Rightarrow x = 0$$

In  $\text{HCOOH}$ ,  $2(+1) + y + 2(-2) = 0$

$$2 + y - 4 = 0 \Rightarrow y = +2$$

28. (a)  $\text{Ni(III)} = 3d^7 4s^0 = \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\phantom{\uparrow\downarrow}}$   $4s$

(b)  $\text{Fe(III)} = 3d^5 4s^0 = \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}}$   $4s$

+ 3 oxidation state of Fe provides the extra stability due to half-filled  $d$ -orbitals.

(c)  $\text{Zn(III)} = 3d^9 4s^0 = \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}}$   $4s$

(d)  $\text{Cu(III)} = 3d^8 4s^0 = \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow\downarrow} \boxed{\uparrow} \boxed{\uparrow} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}} \boxed{\phantom{\uparrow\downarrow}}$   $4s$

29.  $4\overset{(0)}{\text{Na}} + \text{O}_2 \longrightarrow 2\overset{+1}{\text{Na}}_2\text{O}$

Loss of  $e^-$  (oxidation)

In this reaction,

Na converts into ( $\text{Na}^+$ ) ion, i.e. Na donates its electron to oxygen atom. So, it behaves as an reducing agent.

30. Oxidation number of Cl in molecular state (i.e. in  $\text{Cl}_2$ ) is zero.

Let, oxidation number of Cl in  $\text{NaOCl}$  is  $x$ .

$$\therefore 1 + (-2) + x = 0$$

$$x = +1$$

Let, oxidation number of Cl in  $\text{ClO}_3^-$ .

$$\therefore x + 3(-2) = -1$$

$$x = +5$$

31.  $\overset{+2}{\text{O}} = \overset{0}{\text{C}} = \overset{+2}{\text{C}} = \overset{+2}{\text{O}}$

Carbon suboxide

In  $\text{C}_3\text{O}_2$ , two C-atoms linked with oxygen atoms are present in +2 oxidation state and central carbon has zero oxidation state.

So, the average oxidation state of carbon is  $+\frac{4}{3}$ .

32. (i) Let oxidation number of S in  $\text{H}_2\text{SO}_4$  is  $x$ .

$$\therefore +1 \times 2 + x + (-2) \times 4 = 0$$

$$x = 8 - 2 \text{ or } x = +6$$

Therefore, oxidation number of S in  $\text{H}_2\text{SO}_4$  is +6.

(ii) Let oxidation number of S in  $\text{HSO}_3^-$  is  $x$ .

$$(+1) + x + (-2) \times 3 = -1$$

$$x = -1 + 5 \text{ or } x = +4$$

Therefore, oxidation number of S in  $\text{HSO}_3^-$  is +4.

(iii) Let the oxidation number of S in  $\text{SO}_2\text{Cl}_2$  be  $x$ .

$$x + 2(-2) + 2(-1) = 0$$

$$x = +6$$

Therefore, in  $\text{SO}_2\text{Cl}_2$  the oxidation number of sulphur is +6

33. Highest oxidation number of any transition element =  $(n - 1) d$  electrons +  $ns$  electrons. Therefore, larger the number of electrons in the  $3d$ -orbitals, higher is the maximum oxidation number.

(a)  $3d^1 4s^2 = 3$

(b)  $3d^3 4s^2 = 3 + 2 = 5$

(c)  $3d^5 4s^1 = 5 + 1 = 6$  and

(d)  $3d^5 4s^2 = 5 + 2 = 7$

Thus, option (d) is correct.

34. Given, electronic configuration of  $X = 1s^2 2s^2 2p^3$

$\therefore$  The valency of  $X$  will be 3.

The valency of Mg is +2.

$\therefore$  Magnesium reacts with element  $X$  to form an ionic compound with formula  $\text{Mg}_3\text{X}_2$ .

35. The valency of Cr in  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]^+$  ion is  $x$ .

$$x + 4 \times (0) - 2(-1) = +1$$

$$x = 1 + 2$$


$$x = 3$$

So, the valency of Cr in  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]^+$  ion is 3.

36. The correct match is

A → 2, B → 4, C → 1, D → 3.

The oxidation number of N-atom in given compounds are shown below :





37. Electronegativity of fluorine is more than that of oxygen atom, so F gains electron with negative charge.

In oxygen difluoride (OF<sub>2</sub>) and dioxygen difluoride (O<sub>2</sub>F<sub>2</sub>), oxygen transfers electron to fluorine atom. Thus,


Oxidation number of oxygen in OF<sub>2</sub> = + 2.

Oxidation number of oxygen in O<sub>2</sub>F<sub>2</sub> = + 1.

38. Writing the oxidation number (O.N.) of Cr, Cl and Mn on each species in the four set of ions, then,



Only in the arrangement (a), the ON of central atom increases from left to right, therefore, option (a) is correct.




This reaction is represented in basic medium

because in basic medium  $\overset{+7}{\text{Mn}}\text{O}_4^-$  is reduced to


$\overset{+4}{\text{Mn}}\text{O}_2$  (i.e. Mn to Mn), while in acidic medium,  $\overset{+7}{\text{Mn}}\text{O}_4^-$  is reduced from  $\text{Mn}^{7+}$  to  $\text{Mn}^{2+}$ .

40. The balanced chemical reaction is given as



Hence, the value of x, y and z are respectively 3, 8 and 2.


41. When a mixture of potassium chlorate, oxalic acid and sulphuric acid is heated, the following reaction occurs :



Thus, Cl is the element which undergoes maximum change in the oxidation state from +5 to -1.

42. Both Assertion and Reason are correct explanation and Reason is the correct explanation for Assertion.

43. The reaction of potassium permanganate and potassium iodide is as follows :



Oxidation state of Mn decreases from +7 to +2.

Thus, Assertion is correct but Reason is incorrect.


44. KI<sub>3</sub> dissociates into KI and I<sub>2</sub>.

∴ Average oxidation number of I

$$= \frac{-1 \times 1 + 0 \times 2}{1 + 2} = \frac{-1}{3}$$

Thus, either two values are reported separately or one value is reported.

45. PbCl<sub>2</sub> is more stable than PbCl<sub>4</sub> or Pb<sup>2+</sup> is more stable than Pb<sup>4+</sup> (due to the inert pair effect)



Assertion is correct but Reason is incorrect statement.

46. Oxygen is the most electronegative element after fluorine. Therefore, in the compounds between oxygen and fluorine, oxygen is found to show positive oxidation state.


Both Assertion and Reason are correct statements and Reason is the correct explanation of the Assertion.

47. Maximum oxidation state of S is + 6, it cannot exceed beyond, it. Therefore it, cannot be further oxidised.


Both Assertion and Reason are correct statements and Reason is the correct explanation of the Assertion.

48. Among halogen F<sub>2</sub> is the best oxidant because it has the highest E° value.

Both Assertion and Reason are correct, but Reason is not the correct explanation of Assertion.




The oxidation states of Fe in FeO and Fe<sub>2</sub>O<sub>3</sub> are + 2 and + 3.



Both Assertion and Reason are correct statements, but Reason is not the correct explanation of the Assertion.

50. The structure of  $\text{Br}_3\text{O}_8$  (tribromooctaoxide) is



Thus, oxidation state of two corner Br atoms is +6 and of middle one is +4. The difference in oxidation states is due to difference in bonding situations.

$$\text{Average oxidation state} = \frac{+6 + 4 + 6}{3} = \frac{16}{3}$$

Thus, both Assertion and Reason are correct but Reason is not the correct explanation of Assertion.


51. Due to high electronegativity and high heat of association, fluorine oxidises the elements to their highest oxidation state.

Both Assertion and Reason are correct statements, but Reason is not the correct explanation of the Assertion.

52.  $\text{SO}_2$  can act both as an oxidising and a reducing agent.  $\text{O}_3$  can act only as an oxidising agent.

The oxidation number of O in  $\text{O}_3$  is zero. It can only decrease from zero to -1 or -2 but cannot increase to +2. Therefore, it can act as an oxidising agent only. In  $\text{SO}_2$ , the oxidation number of S is +4. It can have a minimum oxidation number of -2 and maximum of +6. Its oxidation number either decreases or increases and hence, it can act both as an oxidising and a reducing agent.

Assertion is correct, but Reason is incorrect statement.



Formal oxidation number of sulphur

$$= \frac{2 \times 5 + 2 \times 0}{4} = 2.5$$

Assertion is correct, but Reason is incorrect statement.

54.  $\text{MnO}_4^-$  is reduced to  $\text{MnO}_2$  in a mild basic medium or neutral medium whereas in an acidic medium,

$\text{MnO}_4^-$  is reduced to  $\text{Mn}^{2+}$  and in a strong basic medium, it is reduced to  $\text{MnO}_4^{2-}$ .

So, Assertion is incorrect but Reason is correct.

55.  $\text{Cl}_2$  is an oxidising agent. It bleaches the articles permanently by oxidation in presence of moisture. Assertion is correct, but Reason is incorrect statement.

56. (i)  $\text{K}_2\text{Cr}_2\text{O}_7 + \text{Excess KI} \longrightarrow \text{Cr}^{3+} + \text{H}_2\text{O} + n\text{I}_2^0$
- $$\begin{array}{c}
 \text{Cr}^{6+} \\
 \xrightarrow{+3e^-} \\
 \text{Cr}^{3+}
 \end{array}$$

$\text{Cr}^{6+}$  accepts  $3e^-$ , so mole of  $\text{I}_2 = 3$ .

- (ii) If zinc rod is dipped in copper sulphate solution, then due to transfer of electron from zinc to copper ion, heat is evolved.
- (iii) A negative  $E^\ominus$  means that the redox couple is a stronger reducing agent than  $\text{H}^+/\text{H}_2$  couple.
- $E^\ominus =$  strong reducing agent
- +  $E^\ominus =$  weak reducing agent

- (iv) Oxidant = oxidising agent = reduces itself by increasing oxidation number.

Reductant = reducing agent = oxidising itself, by decreasing oxidation number.

Or

In alkaline medium, the balanced equation is,

$$2\text{ClO}_2 + 5\text{H}_2\text{O}_2 + 2\text{OH}^- \longrightarrow 2\text{Cl}^- + 5\text{O}_2 + 6\text{H}_2\text{O}$$

2 moles of  $\text{ClO}_2$  oxidise 5 moles of  $\text{H}_2\text{O}_2$

$\therefore$  1 mole of  $\text{ClO}_2$  will oxidise =  $\frac{5}{2} \times 1 = 2.5$ .

57. (i) Let the oxidation state of Mn be x.

(a)  $\text{KMnO}_4$ ,

$$\begin{array}{l}
 +1 + x + (-2) \times 4 = 0 \\
 x - 7 = 0 \\
 x = +7
 \end{array}$$

(b)  $\text{K}_2\text{MnO}_4$ ,

$$\begin{array}{l}
 (+1) \times 2 + x + (-2) \times 4 = 0 \\
 x - 6 = 0 \\
 x = +6
 \end{array}$$

(c)  $\text{Mn}_2\text{O}_3$ ,

$$\begin{array}{l}
 x \times 2 + (-2) \times 3 = 0 \\
 2x = +6 \\
 x = +3
 \end{array}$$


$$(d) \text{MnO}_2, x + (-2) \times 2 = 0$$

$$x = 4$$

Thus, oxidation state of Mn is highest in  $\text{KMnO}_4$ .

- (ii) Fluorine is the most electronegative element.

Hence, does not possess positive oxidation number in any of its compound.






The oxidation number of Mn changes from +2 to +6.

- (iv) Let oxidation state of Fe in complex is  $x$ .


$$\therefore x + 5(0) + (-1) = -2$$

$$\text{or } x = +3$$


Or




(no change in valency)



58. (i) The redox change in the given reaction is as follows :




$\therefore$  Both Assertion and Reason are correct and Reason is the correct explanation of Assertion.



Thus, the above reaction is an example of disproportionation reaction.

Thus, both Assertion and Reason are correct and Reason is the correct explanation of Assertion.

- (iii) The structure of  $\text{Br}_3\text{O}_8$  (tribromooctaoxide) is




Thus, oxidation state of two corner Br atoms is +6 and of middle one is +4. The difference in oxidation states is due to difference in bonding situations.

$$\text{Average oxidation state} = \frac{+6 + 4 + 6}{3} = \frac{16}{3}$$

Thus, both Assertion and Reason are correct but Reason is not the correct explanation of Assertion.

- (iv) The reaction of potassium permanganate and potassium iodide is as follows :



Oxidation state of Mn decreases from +7 to +2.

Thus, Assertion is correct but Reason is incorrect.

Or

As we know  $\text{H}^+/\text{H}_2$  couple has zero standard reduction potential so, ions having positive  $E^\circ$  value are weaker reducing agent, while ions having negative  $E^\circ$  value are stronger reducing agent. Thus, Assertion is incorrect but Reason is correct.


59. (i) Let, oxidation state of N in  $\text{N}_3\text{H}$  be  $x$ .

$$3x + 1 = 0$$

$$x = -\frac{1}{3}$$

Nitrogen is more electronegative than hydrogen.

$\therefore$  Assertion is correct but Reason is incorrect.



Thus, the above reaction is an example of disproportionation reaction.

Both Assertion and Reason are correct and Reason is the correct explanation of Assertion.

- (iii) The electrons are transferred from Zn to  $\text{Cu}^{2+}$  through the metallic wire which connects the two rods.

While electricity flows through the salt-bridge by migration of ions from one beaker to other.

Both Assertion and Reason are correct statements but Reason is not the correct explanation of Assertion.

- (iv) Redox couple is the combination of oxidised and reduced form of a substance.

In the representation  $E^\circ_{\text{Fe}^{3+}/\text{Fe}^{2+}}$  and  $E^\circ_{\text{Cu}^{2+}/\text{Cu}}$ ,  $\text{Fe}^{3+}/\text{Fe}^{2+}$  and  $\text{Cu}^{2+}/\text{Cu}$  are redox couples.

Both Assertion and Reason are correct statements but Reason is not the correct explanation of Assertion.

*Or*

Oxidation number of elements in their compounds or ions is obtained using some rules, e.g. hydrogen is assigned oxidation number of +1 in general and -1 for metal hydrides.

Both Assertion and Reason are correct statements but Reason is not the correct explanation of Assertion.