

Chapter 5. States of Matter

Question-1

What will be the minimum pressure required to compress 500 dm³ of air at 1 bar to 200 dm³ at 30⁰ c?

Solution:

$$P_1 = 1 \text{ bar}$$

$$P_2 = ?$$

$$V_1 = 500 \text{ dm}^3$$

$$V_2 = 200 \text{ dm}^3$$

$$P_1 V_1 = P_2 V_2$$

$$1 \times 500 = P_2 \times 200 \text{ or}$$

$$P_2 = \frac{1 \times 500}{200} = 2.5 \text{ bar.}$$

Question-2

A vessel of 120 mL capacity contains a certain amount of gas at 35 °C and 1.2 bar pressure. The gas is transferred to another vessel of volume 180 mL at 35 °C. What would be its pressure?

Solution:

Let $V_1 = 120$ ml

$V_2 = 180$ ml

$P_1 = 1.2$ bar

$P_2 = ?$

$$P_1 V_1 = P_2 V_2$$

$$1.2 \times 120 = P_2 \times 180$$

$$P_2 = \frac{120}{180} \times 1.2$$

$$P_2 = 0.8 \text{ bar.}$$

Question-3

Using the equation of state $PV = nRT$, show that at a given temperature density of a gas is proportional to gas pressure, P .

Solution:

$$PV = nRT$$

$$(\text{or}) P (\text{Pressure of gas}) = \frac{n \cdot R T}{V}$$

We know $\frac{1}{V} = \rho$ (density)

$$\therefore P = n \cdot \rho R T$$

\therefore Pressure of gas (P) \propto density of the gas (d).

Question-4

At 0°C, the density of a gaseous oxide at 2 bar is same as that of nitrogen at 5 bar. What is the molecule mass of the oxide?

Solution:

Density of nitrogen, $\rho = \frac{P_m}{RT}$

$$= P = 5 \times 0.987 \text{ atm, } m = 28$$

$$= \rho = \frac{5 \times 0.987 \times 28}{0.0821 \times 273}$$

Density of gaseous oxide

$$\frac{5 \times 0.987 \times 28}{0.0821 \times 273} = \frac{2 \times 0.987 \times x}{0.0821 \times 273}$$

$$X = \frac{0.0821 \times 273 \times 5 \times 0.987 \times 28}{0.0821 \times 273 \times 2 \times 0.987}$$

$$\text{or } x = \frac{5 \times 28}{2} = 70 \text{ g/mol.}$$

Question-5

Pressure of 1g of an ideal gas A at 27⁰ C is found to be 2 bar when 2 g of another ideal gas B is introduced in the same flask at same temperature the pressure becomes 3 bar. Find a relationship between their molecular masses

Solution:

Amount of ideal gas, A = $W_A = 1$ g

Its pressure, $P_A = 2$ bar

Amount of another gas, B = $W_B = 2$ g

Total pressure of the flask = 3 bar

∴ Partial Pressure of gas B in the gas mixture = $3 - 2 = 1$ bar

$$PV = nRT$$

$$= \frac{W \cdot R \cdot T}{M}$$

$$\text{For gas A, } P_A = \frac{W_A \cdot R \cdot T}{M_A V}$$

$$2 = \frac{1g \cdot R \cdot T}{M_A \cdot V} \dots\dots\dots (i)$$

For gas B,

$$P_B = \frac{W_B \cdot R \cdot T}{M_B \cdot V}$$

$$1 = \frac{2g \cdot R \cdot T}{M_B \cdot V} \dots\dots\dots (ii)$$

Divide equation (i) by (ii)

$$\frac{2g}{1g} = \frac{1g \cdot R \cdot T}{M_A \cdot V} \times \frac{M_B \cdot V}{2g \cdot R \cdot T}$$

$$\text{or } \frac{M_B}{2 \times M_A} = 2$$


$$4M_A = M_B$$

The molecular mass of M_b is four times of M_a .

Question-6

The drain cleaner, Drainex contains small bits of aluminium which react with caustic soda to produce hydrogen. What volume of hydrogen at 20⁰c and one bar will be released when 0.15 g of aluminium reacts?

Solution:



$$\begin{array}{ll} 2 \times 27 \text{ g of Al} & 3 \times 22.4 \text{ litre of H}_2 \\ = 54 \text{ g} & \end{array}$$

54g Al gives = 3 × 22.4 litre H₂ at N.T.P.

$$0.15 \text{ g Al gives} = \frac{0.15 \times 3 \times 22.4}{54} = 0.186 \text{ litre at N.T.P.}$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Normal Pressure $P_1 = 1 \text{ atm}$

$$P_2 = 1 \text{ bar} = 0.987 \text{ atm}$$

$$V_1 = 0.186 \text{ L}$$

$$V_2 = ?$$

$$T_1 = 273 \text{ K}$$

$$T_2 = 273 + 20 = 293 \text{ K}$$

$$\frac{1 \times 0.186}{273} = \frac{0.987 \times V_2}{293}$$

$$V_2 = \frac{1 \times 0.186 \times 293}{0.987 \times 273} = 0.202 \text{ litre} = 202 \text{ ml.}$$

Question-7

What will be the pressure exerted by a mixture of 3.2 g of methane and 4.4 g of carbon dioxide contained in a 9 dm³ flask at 27°C?

Solution:

$$w_{\text{CH}_4} = 3.2 \text{ g}$$

$$\text{Mol. Weight of CH}_4 = 16$$

$$\text{Moles of CH}_4 = \frac{3.2}{16} = 0.2$$

$$w_{\text{CO}_2} = 4.4 \text{ g}$$

$$\text{Mol. Weight of CO}_2 = 44$$

$$\text{Moles of CO}_2 = \frac{4.4}{44} = 0.1 \text{ mol}$$

$$\text{Total moles present in mixture} = 0.2 + 0.1 = 0.3 \text{ mol}$$

$$\text{Volume of flask} = 9 \text{ dm}^3 = 9 \text{ L}$$

$$\text{Temperature} = 273 + 27 = 300 \text{ K}$$

$$\text{Now for pressure of mixture} = P \times V = nRT$$

$$P \times 9 = 0.3 \times 0.0821 \times 300$$

$$P = \frac{0.3 \times 0.0821 \times 300}{9} = 0.821 \text{ atm}$$

$$0.821 \times 1.01 \times 10^5 \text{ Pa} = 8.29 \times 10^4 \text{ Pa}.$$

Question-8

What will be the pressure of the gas mixture when 0.5 L of H_2 at 0.8 bars and 2.0 L of oxygen at 0.7 bar are introduced in a 1 L vessel at $27^\circ C$?

Solution:

Volume of H_2 , (V_1) 0.5 l

Pressure of H_2 , (P_1) = 0.8 bar

Volume of oxygen, (V_2) = 2.0 l

Pressure of O_2 , (P_2) = 0.7 bar

Total Volume of the flask ($V_1 + V_2$) = 1 l

Total pressure of the gaseous mixture is P

$$P (V_1 + V_2) = P_1 V_1 + P_2 V_2$$

$$P \times 1 = (0.8 \times 0.5) + (0.7 \times 2.0)$$

$$P \times 1 = (0.4 + 1.4)$$

Total Pressure of the mixture of gases , $P = 1.8/1 = 1.8$ bar.

Question-9

Density of a gas is found to be 5.46 g/dm^3 at 27°C at 2 bar pressure. What will be its density at STP?

Solution:

$$\frac{d_1 T_1}{P_1} = \frac{d_2 T_2}{P_2}$$

$$d_1 = 5.46 \text{ g/dm}^3 \quad d_2 = ?$$

$$T_1 = 273 + 27 = 300 \text{ K} \quad T_2 = 273 \text{ K}$$

$$P_1 = 2 \text{ bar}$$

$$\frac{5.46 \times 300}{2} = \frac{d_2 \times 273}{1}$$

$$d_2 = \frac{5.46 \times 300}{2 \times 273} = 3 \text{ g/dm}^3.$$

Question-10

34.05 mL of phosphorous vapour weighs 0.00625 g at 546° C and 0.1 bar pressure. What is the molar mass of phosphorous?

Solution:

$$P = 0.1 \text{ bar}, V = 34.05 \text{ mL} = 0.03405$$

$$R = 0.0821 \text{ litre-atm K}^{-1} \text{ mol}^{-1}$$

$$T = 273 + 546 = 819 \text{ K}$$

$$PV = nRT$$

$$n = \frac{PV}{RT} = \frac{0.1 \times 0.03405}{0.0821 \times 819} \\ = 0.00005 \text{ mol}^{-1}$$

$$\text{Weight of phosphorous vapour} = 0.00625 \text{ g}$$

$$\text{Molar mass of phosphorus} = \frac{0.00625 \text{ g}}{0.00005 \text{ mol}^{-1}} = 125 \text{ g.}$$

Question-11

A student forgot to add the reaction mixture to the round bottomed flask at 27°C but put it on the flame. After a lapse of time, he realized his mistake, using a pyrometer he found the temperature of the flask was 477°C. What fraction of air would have been expelled out?

Solution:

$$\frac{V_1}{V_2} = \frac{T_1}{T_2}$$

$$\text{Or } \frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$T_1 = 27 + 273 = 300 \text{ K}$$

$$T_2 = 477 + 273 = 750 \text{ K}$$

$$\frac{V_1}{300} = \frac{V_2}{750}$$

$$\frac{V_1}{V_2} = \frac{300}{750} = \frac{2}{5}$$

If the total volume of flask is taken as 1

$$\text{Fraction of air expelled} = 1 - \frac{2}{5} = \frac{3}{5}.$$

Question-12

Calculate the temperature of 4.0 moles of a gas occupying 5 dm³ at 3.32 bar (R=0.083 bar dm³ K⁻¹ mol⁻¹).

Solution:

$$PV = nRT$$

$$P = 3.32 \text{ bar}, V = 5 \text{ dm}^3$$

$$n = 4 \text{ moles}$$

$$R = 0.083 \text{ bar dm}^3 \text{ K}^{-1} \text{ mol}^{-1}$$

$$\therefore 3.32 \times 5 = 4 \times 0.083 \times T$$

$$T = \frac{3.32 \times 5}{4 \times 0.083} = 50 \text{ K}.$$

Question-13

Calculate the total number of electrons present in 1.4g of Nitrogen gas.

Solution:

Atomic weight of Nitrogen = 14 g

weight of Nitrogen gas = 1.4 g

$$\text{Number of gram Moles of Nitrogen atoms} = \frac{W}{M} = \frac{1.4}{14} = 0.1 \text{ mol}.$$

No. of atoms of Nitrogen, present in 1 mol of Nitrogen gas = 6.022×10^{23} atoms

1 atom of Nitrogen contains = 7 electrons

$$\text{No. of electrons, present in 1 mol of Nitrogen atoms} = 7 \times 6.022 \times 10^{23}$$

$$\text{No. of electrons in 0.1 mol} = 0.1 \times 7 \times 6.02 \times 10^{23} = 4.22 \times 10^{23}.$$

Question-14

How much time it would take to distribute one Avogadro number of wheat grains if 10^{10} grains are distributed each second?

Solution:

10^{10} grains are distributed 1 s Time in seconds, taken to distribute

$$6.023 \times 10^{23} \text{ grains} = \frac{6.023 \times 10^{23}}{10^{10}} \text{ s} = \frac{6.023 \times 10^{23}}{10^{10} \times 60 \times 60 \times 24 \times 365} \text{ years} = 1,90,800$$

years.

Question-15

Calculate the total pressure in a mixture of 8 g of oxygen and 4 g of hydrogen confined in a vessel of 1 dm^3 at 27°C , $R = 0.083 \text{ bar dm}^3 \text{ K}^{-1} \text{ mol}^{-1}$.

Solution:

Weight of oxygen $w_{\text{O}_2} = 8 \text{ g}$

$$\text{Moles of O}_2 = \frac{8}{32} = \frac{1}{4}$$

$$w_{\text{H}_2} = 4 \text{ g}$$

$$\text{Moles of H}_2 = \frac{4}{2} = 2$$

$$\text{Total moles present in mixture} = \frac{1}{4} + 2 = \frac{9}{4}$$

$$\text{Volume of vessel} = 1 \text{ dm}^3 = 1 \text{ L}$$

$$\text{Temperature} = 27^\circ\text{C} = 27 + 273 = 300 \text{ K}$$

$$\text{Now for pressure of mixture} = P \times V = n \times RT$$

$$P \times 1 = \frac{9}{4} \times 0.083 \times 300$$

$$P = 56.025 \text{ bar.}$$

Question-16

Pay load is defined as the difference between the mass of displaced air and the mass of the balloon. Calculate the pay load when a balloon of radius 10m, mass 100kg is filled with helium at 1.66 bar at 27°C. (Density of air 1.2 kg m⁻³ and R = 0.083 bar dm³ K⁻¹ mol⁻¹).

Solution:

Volume of balloon, $V = \frac{4}{3}\pi r^3$

$$r = 10 \text{ m}$$

$$V = \frac{4}{3} \times 3.14 \times 1(10)^3 = 4187 \text{ m}^3$$

Volume of air displaced = Volume of balloon = 4187 m³

$M = \text{Vol} \times \text{Density}$

Mass of displaced air = 4187 m³ × 1.2 kg m⁻³ = 5024.4 kg

Temp, T = 27+273=300 K

$$\begin{aligned} \text{Moles of gas present, } n &= \frac{PV}{RT} \\ &= \frac{1.66 \times 4187}{0.083 \times 300} = 279.1 \times 10^3 \text{ mol.} \end{aligned}$$

Mass = No. of moles × Molecular weight

$$\begin{aligned} \text{Mass of He present} &= 279 \times 10^3 \times 4 \\ &= 1116.4 \times 10^3 \text{ g} \\ &= 1116.4 \text{ kg} \end{aligned}$$

Mass of filled balloon = 100 + 1116.4 = 1216.4 Kg

$$\begin{aligned} \text{Pay load} &= \text{Mass of displaced air} - \text{Mass of balloon} \\ &= 5024.4 - 1216.4 \\ &= 3808 \text{ kg.} \end{aligned}$$

Question-17

Calculate the volume occupied by 8.8 g of CO_2 at 31.1°C and 1 bar pressure. $R = 0.083 \text{ bar l K}^{-1} \text{ mol}^{-1}$.

Solution:

$$P = 1 \text{ bar}, T = 31.1 + 273 = 304.1 \text{ K}$$

$$\text{Mol. wt of } \text{CO}_2 = 44$$

$$\text{Weight of } \text{CO}_2 = 8.8 \text{ g}$$

$$PV = (W/m)RT$$

$$1 \times V = \frac{8.8}{44} \times 0.083 \times 304.1$$

$$V = 5.05 \text{ L.}$$

Question-18

2.9 g of a gas at 95°C occupied the same volume as 0.184 g of hydrogen at 17°C at the same pressure. What is the molar mass of the gas?

Solution:

Ideal gas equation, $PV = nRT$

1st Case

Let the molar mass of gas (M) = $M(\text{g mol}^{-1})$

Mass of gas (m) = 2.9 g

No. of moles (n) = $\frac{\text{Mass}}{\text{Molarmass}} = \frac{2.9}{M} \text{ mol}$

$T = 273 + 95 = 368$

Pressure = P, Volume = V

$PV = nRT$

$$PV = \frac{2.9}{M} \times R \times 368 \dots\dots\dots (i)$$

2nd Case

Mass of hydrogen = 0.184 g

Number of moles of $\text{H}_2 = \frac{0.184}{2} \text{ mol}$

$T = 273 + 17 = 290$

$$PV = \frac{0.184}{2} \times R \times 290 \dots\dots\dots (ii)$$

In both gases PV is same

From (i) and (ii)

$$\frac{2.9}{M} \times 368 = \frac{0.184 \times 290}{2}$$

$$M = \frac{2.9 \times 368 \times 2}{0.184 \times 290} = 40 \text{ g mol}^{-1}.$$

Question-19

A mixture of dihydrogen and dioxygen at one bar pressure contains 20% by weight of dihydrogen. Calculate the partial pressure of dihydrogen.

Solution:

Percentage of $H_2 = 20\%$, say 20g

No. of moles of $H_2 = \frac{20}{2} = 10$ moles

Mass of oxygen = $(100-20) = 80$ g)

No. of moles of $O_2 = \frac{80}{32} = 2.5$ moles

Mole fraction of $H_2 = \frac{10}{(10+2.5)} = \frac{10}{12.5}$

Partial pressure of $H_2 = \text{Total Pressure} \times \text{Mole fraction of } H_2$

Total Pressure = 1 bar

$$= 1 \times \frac{10}{12.5} = 0.8 \text{ bar.}$$

Question-20

What would be the SI unit for the quantity $pV^2T^{2/n}$?

Solution:

$$\text{S.I. unit} = \frac{P \times (m^3)^2 K^2}{mol} = Pa \text{ m}^6 K^2 mol^{-1}.$$

Question-21

In terms of Charles' law explain why -273°C is the lowest possible temperature.

Solution:

Charles law states that Pressure, remaining constant the volume of a given mass of gas increases or decreases by $1/273$ of its volume at 0°C for every 1°C rise or fall in temperature .

$$V_t = V_0 \left(1 + \frac{t}{273} \right)$$

$$\text{If } t = -273^{\circ}\text{C}$$

$$\begin{aligned} V_t &= V_0 \left(1 - \frac{273}{273} \right) \\ &= 0 \end{aligned}$$

Thus at -273°C any volume of a given mass of gas becomes zero. Thus -273°C is said to be limiting temperature and it is equal to OK (absolute scale).

Question-22

Critical temperature for carbon dioxide and methane are 31.1°C and -81.9°C respectively. Which of these has stronger intermolecular forces and why?

Solution:

CO_2 can more easily be liquified than Methane which shows that intermolecular forces in CO_2 are more stronger than in Methane, because in CO_2 polarity exists.

Question-23

Explain the physical significance of van der Waals parameters.

Solution:

Vander Waals equation of state for mol of gas = $\left(p + \frac{a}{v^2}\right)(v-b) = RT$

For n moles of gas = $\left(p + \frac{n^2 a}{v^2}\right)(v-nb) = RT$

The constant 'a' tells the magnitude of attractive forces between the molecules of the gas. In other words larger the value of 'a', greater the intermolecular attraction.

The constant 'b' relates to non compressible volume of gas, as gas molecules occupy some volume, however small may be, at high pressures and low temperatures.∴ Actual volume of gas = Available volume - Non compressible volume = (V - b).