

Expansion of Determinants

1 Mark Questions

1. If $\begin{vmatrix} 2x & 5 \\ 8 & x \end{vmatrix} = \begin{vmatrix} 6 & -2 \\ 7 & 3 \end{vmatrix}$, then write the value of x . Delhi 2014



Firstly, expand both determinants, which gives equation in x and then solve that equation for finding the value of x .

$$\text{Given, } \begin{vmatrix} 2x & 5 \\ 8 & x \end{vmatrix} = \begin{vmatrix} 6 & -2 \\ 7 & 3 \end{vmatrix}$$

$$\Rightarrow 2x^2 - 40 = 18 - (-14)$$

$$\Rightarrow 2x^2 - 40 = 18 + 14$$

$$\Rightarrow 2x^2 - 40 = 32$$

$$\Rightarrow 2x^2 = 32 + 40$$

$$\Rightarrow 2x^2 = 72 \Rightarrow x^2 = 36$$

$$\therefore x = \pm 6 \quad (1)$$

2. If $\begin{vmatrix} 3x & 7 \\ -2 & 4 \end{vmatrix} = \begin{vmatrix} 8 & 7 \\ 6 & 4 \end{vmatrix}$, then find the value of x .
All India 2014

Given, $\begin{vmatrix} 3x & 7 \\ -2 & 4 \end{vmatrix} = \begin{vmatrix} 8 & 7 \\ 6 & 4 \end{vmatrix}$

On expanding the determinant of both sides,
we get

$$\begin{aligned} 3x \times 4 - (-2) \times 7 &= 8 \times 4 - 6 \times 7 \\ \Rightarrow 12x - (-14) &= 32 - 42 \\ \Rightarrow 12x + 14 &= -10 \\ \Rightarrow 12x &= -10 - 14 \\ \Rightarrow 12x &= -24 \Rightarrow x = -\frac{24}{12} \end{aligned}$$

$$\therefore x = -2$$

(1)

3. If A is a 3×3 matrix, $|A| \neq 0$ and $|3A| = k|A|$, then
write the value of k .
Foreign 2014

We know that, if A is a square matrix of order n .
Then, $|kA| = k^n |A|$.

Here, the matrix A is of order 3.

$$\therefore |3A| = (3)^3 |A| = 27 |A|$$

On comparing with $k|A|$, we get (1)

$$k = 27$$

4. Find $(\text{adj } A)$, if $A = \begin{bmatrix} 5 & 2 \\ 7 & 3 \end{bmatrix}$.

Delhi 2014C

Given, $A = \begin{bmatrix} 5 & 2 \\ 7 & 3 \end{bmatrix}$

$$\therefore |A| = \begin{vmatrix} 5 & 2 \\ 7 & 3 \end{vmatrix} = 15 - 14 = 1$$

We know that, $|\text{adj}(A)| = |A|^{n-1}$, where n is order of determinant.

$$\therefore |\text{adj}(A)| = |1|^{2-1} \Rightarrow |\text{adj}(A)| = 1 \quad (1)$$

5. Write the value of the determinant

$$\begin{vmatrix} p & p+1 \\ p-1 & p \end{vmatrix}$$

Delhi 2014C

Suppose $A = \begin{vmatrix} p & p+1 \\ p-1 & p \end{vmatrix}$

On expanding, we get

$$A = p^2 - (p-1)(p+1)$$

$$\Rightarrow A = p^2 - (p^2 - 1^2)$$

$$[\because a^2 - b^2 = (a+b)(a-b)]$$

$$\Rightarrow A = p^2 - p^2 + 1$$

$$\therefore A = 1$$

6. If A is a square matrix of order 3 such that $|\text{adj}(A)| = 64$, then find $|A|$. Delhi 2013C

We know that, for a square matrix of order n , $|\text{adj}(A)| = |A|^{n-1}$ have $n = 3$

$$\therefore |\text{adj}(A)| = |A|^{3-1} = |A|^2$$

$$\text{Given, } |\text{adj}A| = 64, \quad 64 = |A|^2 \Rightarrow (8)^2 = |A|^2$$

$$\Rightarrow |A| = \pm 8 \quad [\text{taking square root}] \quad (1)$$

7. If $\begin{vmatrix} 2x & x+3 \\ 2(x+1) & x+1 \end{vmatrix} = \begin{vmatrix} 1 & 5 \\ 3 & 3 \end{vmatrix}$, then find the value of x . Delhi 2013C



Expand both determinants which gives equation in x and then solve that equation for finding the value of x .

$$\text{Given, } \begin{vmatrix} 2x & x+3 \\ 2(x+1) & x+1 \end{vmatrix} = \begin{vmatrix} 1 & 5 \\ 3 & 3 \end{vmatrix}$$

$$\Rightarrow 2x(x+1) - (x+3)(2x+2) = 3 - 15$$

$$\Rightarrow 2x^2 + 2x - (2x^2 + 8x + 6) = -12$$

$$\Rightarrow -6x - 6 = -12$$

$$\Rightarrow 6x = 6$$

$$\therefore x = 1 \quad (1)$$

8. If $\begin{vmatrix} x+1 & x-1 \\ x-3 & x+2 \end{vmatrix} = \begin{vmatrix} 4 & -1 \\ 1 & 3 \end{vmatrix}$, then write the value of x . Delhi 2013

$$\text{Given, } \begin{vmatrix} x+1 & x-1 \\ x-3 & x+2 \end{vmatrix} = \begin{vmatrix} 4 & -1 \\ 1 & 3 \end{vmatrix}$$

$$\Rightarrow (x+1)(x+2) - (x-3)(x-1) = 12 + 1$$

$$\Rightarrow (x^2 + 3x + 2) - (x^2 - 4x + 3) = 13$$

$$\Rightarrow 7x - 1 = 13$$

$$\Rightarrow 7x = 14$$

$$\therefore x = 2 \quad (1)$$

9. If A_{ij} is the cofactor of the element a_{ij} of the

$$\text{determinant } \begin{vmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{vmatrix}, \text{ then write the}$$

value of $a_{32} \cdot A_{32}$.

All India 2013; HOTS

$$\text{Let } A = \begin{vmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{vmatrix} = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

Here, $a_{32} = 5$

Given, A_{ij} is the cofactor of the element a_{ij} of A .

$$\begin{aligned} \text{Then, } A_{32} &= (-1)^{3+2} \begin{vmatrix} 2 & 5 \\ 6 & 4 \end{vmatrix} \\ &= (-1)^5 (8 - 30) = -(-22) = 22 \end{aligned}$$

$$\therefore a_{32} \cdot A_{32} = 5 \times 22 = 110 \quad (1)$$

10. Let A be a square matrix of order 3×3 . Write the value of $|2A|$, where $|A| = 4$. All India 2012

We know that, for a square matrix A of order n ,

$$|kA| = k^n \cdot |A|$$

Here, $|2A| = 2^3 \cdot |A|$ [\because order of A is 3×3]

$$= 2^3 \times 4 = 8 \times 4 = 32 \quad [\because \text{put } |A| = 4] \quad (1)$$

11. If $\Delta = \begin{vmatrix} 5 & 3 & 8 \\ 2 & 0 & 1 \\ 1 & 2 & 3 \end{vmatrix}$, then write the minor of the

element a_{23} .

Delhi 2012

Minor of the element

$$a_{23} = \begin{vmatrix} 5 & 3 \\ 1 & 2 \end{vmatrix} = 10 - 3 = 7 \quad (1)$$

12. If the determinant of matrix A of order 3×3 is of value 4, then write the value of $|3A|$.

All India 2012C

Given, the order of matrix A is 3×3 and

$$|A| = 4$$

$$\Rightarrow |3A| = 3^3 \cdot |A| \quad [\because |KA| = K^n \cdot |A|]$$

$$= 3^3 \cdot 4 = 27 \cdot 4 = 108 \quad (1)$$

13. For what value of x , $A = \begin{bmatrix} 2(x+1) & 2x \\ x & x-2 \end{bmatrix}$ is a singular matrix?

All India 2011C



For a singular matrix, $|A| = 0$. Use this relation and solve it.

Matrix A is said to be singular, if $|A| = 0$

$$\therefore \begin{vmatrix} 2x+2 & 2x \\ x & x-2 \end{vmatrix} = 0$$

$$\Rightarrow (2x+2)(x-2) - 2x^2 = 0$$

$$\Rightarrow 2x^2 - 2x - 4 - 2x^2 = 0 \Rightarrow -2x = 4$$

$$\therefore x = -2 \quad (1)$$

- 14.** For what value of x , the matrix $\begin{bmatrix} 2x+4 & 4 \\ x+5 & 3 \end{bmatrix}$ is a singular matrix? All India 2011C

$$\text{Let } A = \begin{bmatrix} 2x+4 & 4 \\ x+5 & 3 \end{bmatrix}$$

If matrix A is singular, then

$$|A| = 0$$

$$\therefore \begin{vmatrix} 2x+4 & 4 \\ x+5 & 3 \end{vmatrix} = 0$$

$$\Rightarrow (2x+4) \times 3 - (x+5) \times 4 = 0$$

$$\Rightarrow 6x + 12 - 4x - 20 = 0 \Rightarrow 2x = 8$$

$$\therefore x = 4 \quad (1)$$

- 15.** For what value of x , the matrix $\begin{bmatrix} 2x & 4 \\ x+2 & 3 \end{bmatrix}$ is a singular matrix? Delhi 2011C

Do same as Que. 14.

[Ans. $x = 4$]

- 16.** For what value of x , matrix $\begin{bmatrix} 6-x & 4 \\ 3-x & 1 \end{bmatrix}$ is a singular matrix? Delhi 2011C

Do same as Que. 14.

[Ans. $x = 2$]

17. For what value of x , the matrix

$$\begin{bmatrix} 5-x & x+1 \\ 2 & 4 \end{bmatrix} \text{ is a singular?}$$

Delhi 2011

Do same as Que. 14.

[Ans. $x = 3$]

18. Evaluate $\begin{vmatrix} \cos 15^\circ & \sin 15^\circ \\ \sin 75^\circ & \cos 75^\circ \end{vmatrix}$.

All India 2011; HOTS



Firstly, expand the determinant and use the trigonometric relation to calculate the value of determinant.

$$A = \begin{vmatrix} \cos 15^\circ & \sin 15^\circ \\ \sin 75^\circ & \cos 75^\circ \end{vmatrix}$$

On expanding, we get

$$\begin{aligned} A &= (\cos 15^\circ \cos 75^\circ - \sin 15^\circ \sin 75^\circ) \\ &= \cos (15^\circ + 75^\circ) \end{aligned}$$

$$\begin{aligned} [\because \cos x \cos y - \sin x \sin y &= \cos (x + y)] \\ &= \cos 90^\circ = 0 \quad [\because \cos 90^\circ = 0] \quad (1) \end{aligned}$$

19. If $\begin{vmatrix} x & x \\ 1 & x \end{vmatrix} = \begin{vmatrix} 3 & 4 \\ 1 & 2 \end{vmatrix}$, then write the positive

value of x .

Foreign 2011; All India 2008C



Expand both determinants which gives equation in x and then solve that equation for finding the value of x .

Given, $\begin{vmatrix} x & x \\ 1 & x \end{vmatrix} = \begin{vmatrix} 3 & 4 \\ 1 & 2 \end{vmatrix}$

On expanding, we get

$$x^2 - x = 6 - 4$$

$$\Rightarrow x^2 - x - 2 = 0 \Rightarrow (x - 2)(x + 1) = 0$$

$$\therefore x = 2 \text{ or } -1$$

Hence, the positive value of x is 2. (1)

20. What is the value of determinant $\begin{vmatrix} 0 & 2 & 0 \\ 2 & 3 & 4 \\ 4 & 5 & 6 \end{vmatrix}$?

Delhi 2010



Determinant can be easily expand either corresponding to a row or column which have maximum zeroes.

Given, determinant

$$A = \begin{vmatrix} 0 & 2 & 0 \\ 2 & 3 & 4 \\ 4 & 5 & 6 \end{vmatrix}$$

$$\Rightarrow |A| = -2(12 - 16)$$

[\because expanding along R_1]

$$= -2(-4) = 8 \quad (1)$$

21. Find the minor of the element of second row and third column in the determinant

$$\begin{vmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{vmatrix}$$

Delhi 2010

Minor of the element of second row and third column is given by

$$M_{23} = \begin{vmatrix} 2 & -3 \\ 1 & 5 \end{vmatrix} = 10 + 3 = 13 \quad (1)$$

22. If $A = \begin{bmatrix} 3 & 1 \\ 2 & -3 \end{bmatrix}$, then find $|\text{adj } A|$.
Delhi 2010C; HOTS

Given, $A = \begin{bmatrix} 3 & 1 \\ 2 & -3 \end{bmatrix}$

Cofactors of A are

$$C_{11} = -3, C_{12} = -2, C_{21} = -1, C_{22} = 3$$

We know that, adjoint $A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$

$$\therefore \text{adj}(A) = \begin{bmatrix} -3 & -2 \\ -1 & 3 \end{bmatrix}^T = \begin{bmatrix} -3 & -1 \\ -2 & 3 \end{bmatrix}$$

$$\begin{aligned} \text{Now } |\text{adj}(A)| &= \begin{vmatrix} -3 & -1 \\ -2 & 3 \end{vmatrix} = -3 \times 3 - (-1 \times -2) \\ &= -9 - 2 = -11 \end{aligned}$$

$$\Rightarrow |\text{adj}(A)| = -11 \quad (1)$$

Alternate Method

$$\text{Here, } |A| = \begin{vmatrix} 3 & 1 \\ 2 & -3 \end{vmatrix} = -9 - 2 = -11$$

Using the result

$$|\text{adj}(A)| = |A|^{n-1}$$

where, n is order of a determinant, we get

$$|\text{adj}(A)| = (-11)^{2-1} = -11 \quad (1)$$

23. If $|A| = 2$, where A is a 2×2 matrix, then find $|\text{adj } A|$.
All India 2010C

Given, $|A| = 2$, where A is a 2×2 matrix.

We know that, $|\text{adj}(A)| = |A|^{n-1}$, where n is the order of matrix. Here, we have

$$n = 2 \text{ and } |A| = 2$$

$$\therefore |\text{adj}(A)| = (2)^{2-1}$$

$$\Rightarrow |\text{adj}(A)| = 2 \quad (1)$$

24. What positive value of x makes following pair of determinants equal?

$$\begin{vmatrix} 2x & 3 \\ 5 & x \end{vmatrix} = \begin{vmatrix} 16 & 3 \\ 5 & 2 \end{vmatrix}$$

All India 2010

$$\text{Given, } \begin{vmatrix} 2x & 3 \\ 5 & x \end{vmatrix} = \begin{vmatrix} 16 & 3 \\ 5 & 2 \end{vmatrix}$$

On expanding, we get

$$2x^2 - 15 = 32 - 15 \Rightarrow 2x^2 - 15 = 17$$

$$\Rightarrow 2x^2 = 32 \Rightarrow x^2 = 16$$

$$\Rightarrow x = \pm 4 \quad (1)$$

Hence, for $x = 4$, given pair of determinants is equal.

25. If A is a non-singular matrix of order 3 and $|\text{adj } A| = |A|^k$, then what is the value of k ?

All India 2009C; HOTS

We know that, for a square matrix of order n
 $|\text{adj}(A)| = |A|^{n-1}$

Here, the order of $A = 3 \times 3$, then $n = 3$

$$\therefore |\text{adj}(A)| = |A|^2 \quad \dots(i)$$

$$\text{But } |\text{adj}(A)| = |A|^k \quad [\text{given}] \dots(ii)$$

From Eqs. (i) and (ii), we get

$$k = 2 \quad (1)$$

26. Evaluate $2 \begin{vmatrix} 7 & -2 \\ -10 & 5 \end{vmatrix}$. Delhi 2009C

$$2 \begin{vmatrix} 7 & -2 \\ -10 & 5 \end{vmatrix} = 2 [35 - (20)] = 2 (35 - 20) \\ = 2 \times 15 = 30 \quad (1)$$

NOTE Suppose we want to multiply with 2 inside the determinant, then we do not multiply each element of determinant. Here, we multiply any one row or column by 2.

27. Find x from equation $\begin{vmatrix} x & 4 \\ 2 & 2x \end{vmatrix} = 0$. All India 2009

Given, $\begin{vmatrix} x & 4 \\ 2 & 2x \end{vmatrix} = 0$

$$\Rightarrow 2x^2 - 8 = 0 \Rightarrow 2x^2 = 8 \Rightarrow x^2 = 4$$

$$\therefore x = \pm 2 \quad (1)$$

28. If $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$, then find the value of k , if $|2A| = k \cdot |A|$. Foreign 2009

Given, $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$ and $|2A| = k \cdot |A|$

$$\Rightarrow 2^2 \cdot |A| = k \cdot |A|$$

$[\because \text{for a square matrix of order } 2 \mid kA \mid = k^2 \cdot |A|, k \text{ is any scalar}]$

$$\therefore k = 4 \quad (1)$$

29. Evaluate $\begin{vmatrix} 2 \cos \theta & -2 \sin \theta \\ \sin \theta & \cos \theta \end{vmatrix}$. Delhi 2008C

$$\text{Suppose } A = \begin{vmatrix} 2 \cos \theta & -2 \sin \theta \\ \sin \theta & \cos \theta \end{vmatrix}$$

On expanding, we get

$$\begin{aligned} A &= 2 \cos^2 \theta - (-2 \sin^2 \theta) \\ &= 2 \cos^2 \theta + 2 \sin^2 \theta \\ &= 2 (\cos^2 \theta + \sin^2 \theta) \\ &= 2 \quad [\because \cos^2 \theta + \sin^2 \theta = 1] \quad (1) \end{aligned}$$

30. Evaluate $\begin{vmatrix} a + ib & c + id \\ -c + id & a - ib \end{vmatrix}$. Delhi 2008; HOTS

$$\text{Suppose } A = \begin{vmatrix} a + ib & c + id \\ -c + id & a - ib \end{vmatrix}$$

On expanding, we get

$$\begin{aligned} A &= (a + ib)(a - ib) - (c + id)(-c + id) \\ &= (a^2 - i^2 b^2) - (-c^2 + i^2 d^2) \\ &\quad [\because (a + b)(a - b) = a^2 - b^2] \\ &= (a^2 + b^2) - (-c^2 - d^2) \quad [\because i^2 = -1] \\ &= a^2 + b^2 + c^2 + d^2 \quad (1) \end{aligned}$$

31. Find for what value of x , is the following matrix singular?

$$\begin{vmatrix} 3 - 2x & x + 1 \\ 2 & 4 \end{vmatrix}$$

Delhi 2008

Do same as Que. 14.

[Ans. $x = 1$]

32. If $\begin{vmatrix} 2x + 5 & 3 \\ 5x + 2 & 9 \end{vmatrix} = 0$, then find the value of x . Foreign 2008

Do same as Que. 27.

[Ans. $x = -13$]