
Sample Paper-01
Mathematics
Class – XI

Time allowed: 3 hours

Maximum Marks: 100

General Instructions:

- a) All questions are compulsory.
 - b) The question paper consists of 26 questions divided into three sections A, B and C. Section A comprises of 6 questions of one mark each, Section B comprises of 13 questions of four marks each and Section C comprises of 7 questions of six marks each.
 - c) All questions in Section A are to be answered in one word, one sentence or as per the exact requirement of the question.
 - d) Use of calculators is not permitted.
-

Section A

1. Find the number of subsets of a set A containing 10 elements
2. How many ways can you choose one or more students from 3 students?
3. In How many ways can one choose 3 cards from a pack of 52 cards in succession (1) with replacement (2) without replacement?
4. State the condition under which the product of two complex numbers is purely imaginary
5. In a circle of radius 1 unit what is the length of the arc that subtends an angle of 2 radians at the centre.
6. Is $\cos \theta$ positive or negative if $\theta = 500 \text{ radians}$

Section B

7. Prove by mathematical induction that $n(n+1)(2n+1)$ is divisible by 6 if n is a natural number
 8. Solve $\cos 2x - 5 \sin x - 3 = 0$
 9. For what values of m the equation $m^2 x^2 + 2(m+1)x + 4 = 0$ will have exactly one zero
 10. Three numbers are in AP. Another 3 numbers are in GP. The sum of first term of the AP and the first term of the GP is 85, the sum of second term of AP and the second term of the GP is 76 and that of the 3rd term of AP and 3rd term of GP is 84. The sum of the AP is 126. Find each term of AP and GP
 11. If $f(x) = 4^x$ find $f(x+1) - f(x)$ in terms of $f(x)$
-

-
12. If $f(x) = \log \frac{(1+x)}{(1-x)}$ Prove that $f\left(\frac{3x+x^3}{1+3x^2}\right) = 3f(x)$ when $-1 < x < 1$
 13. Find the value of $\sin 75^\circ$ and $\cos 75^\circ$
 14. Prove that $\frac{\sin 3\theta}{\sin \theta} - \frac{\cos 3\theta}{\cos \theta} = 2$
 15. If the line $y = mx + 1$ is a tangent to the ellipse $x^2 + 4y^2 = 1$ then find the value of m^2
 16. Reduce the equation $3x - 4y + 20 = 0$ in to normal form
 17. Solve the inequality $\frac{x+3}{x-7} \leq 0$
 18. Find $\lim_{x \rightarrow \infty} \frac{x^2 - ax + 4}{3x^2 - bx + 7}$
 19. Find $\lim_{x \rightarrow 0} \frac{\tan x}{\sin 3x}$

Section C

20. Evaluate $x^3 + x^2 - 4x + 13$ when $x = 1 + i$ and when $x = 1 - i$
 21. Prove that the roots of the equation $(x - \alpha)(x - \beta) = k^2$ is always real
 22. If the roots of the equation $lx^2 + nx + n = 0$ are in the ratio $p : q$ then prove that $\frac{\sqrt{p}}{\sqrt{q}} + \frac{\sqrt{q}}{\sqrt{p}} + \frac{\sqrt{n}}{\sqrt{l}} = 0$
 23. Find $\lim_{x \rightarrow \pi} (\pi - x) \tan \frac{x}{2}$
 24. If a, b, c are 3 consecutive integers prove that $(a-i)(a+i)(c+i)(c-i) = b^4 + 1$
 25. Prove that $\frac{(1+i)^n}{(1-i)^{n-2}} = 2i^{n-1}$
 26. Determine the coordinates of a point which is equidistant from the point $(1, 2)$ and $(3, 4)$ and the shortest distance from the line joining the points $(1, 2)$ and $(3, 4)$ to the required point is $\sqrt{2}$
-

Sample Paper-01
Mathematics
Class – XI

ANSWERS

Section A

1. **Solution**

Number of subsets

$${}^{10}C_0 + {}^{10}C_1 + {}^{10}C_2 + {}^{10}C_3 + {}^{10}C_4 + {}^{10}C_5 + {}^{10}C_6 + {}^{10}C_7 + {}^{10}C_8 + {}^{10}C_9 + {}^{10}C_{10} = 2^{10}$$

2. **Solution**

$${}^3C_1 + {}^3C_2 + {}^3C_3 = 2^3 - 1 = 7$$

3. **Solution**

1. Each card can be drawn in 52 ways and so the total number of ways =

$$52 \times 52 \times 52 = 52^3$$

2. If there is no replacement the first card can be drawn in 52 ways, the second by 51 ways and the third by 50 ways. Hence the total number of ways is

$$52 \times 51 \times 50 = 132600$$

4. **Solution:**

1. None of the factors are zero

2. Factors must be of the form $(a + ib); k(b + ia)$ where k is a real number

5. **Solution**

Length of arc = $r\theta$

Hence length of arc = 2 units

6. **Solution**

1 Full rotation is 2π radians

$$500 \text{ radians} = \frac{500}{2\pi} \text{ rotations}$$

$$\frac{500}{2\pi} = 79.57 \text{ rotations}$$

79 full rotations and 0.57 of a rotation

$$0.5 < 0.57 < 0.75$$

The incomplete rotation is between $\frac{1}{2}$ and $\frac{3}{4}$ of a rotation . Hence 500 radians is in third quadrant. So $\cos \theta$ is negative

Section B

7. Solution

Let

$$n = 1$$

Then $n(n+1)(2n+1) = 6$ and divisible by 6

Let it be divisible by 6 for

$$n = m$$

Then

$$m(m+1)(2m+1) = 6k \text{ Where } k \text{ is an integer}$$

For $n = m+1$ the expression is

$$\begin{aligned} (m+1)(m+2)(2m+2+1) &= (m+2)(m+1)(2m+1) + 2(m+1)(m+2) \\ &= m(m+1)(2m+1) + 2(m+1)(2m+1) + 2(m+1)(m+2) \\ &= m(m+1)(2m+1) + 2(m+1)(3m+3) \\ &= m(m+1)(2m+1) + 6(m+1)^2 \\ &= 6k + 6(m+1)^2, \text{ This is divisible by 6} \end{aligned}$$

8. Solution

$$1 - 2\sin^2 x - 5\sin x - 3 = 0$$

$$2\sin^2 x + 5\sin x + 2 = 0$$

Let $\sin x = t$

$$\text{Then, } 2t^2 + 5t + 2 = 0$$

Solving this quadratic

$$2t(t+2) + (t+2) = 0$$

$$(2t+1)(t+2) = 0$$

$$t = -2, t = -\frac{1}{2}$$

$$\sin x = \frac{-1}{2}$$

First value of t is rejected as $\sin x$ should lie between $(-1 \text{ and } 1)$

General solution is $x = (-1)^{n+1} \frac{\pi}{6} + n\pi$

9. **Solution**

When

$$m = 0$$

The given equation reduces to a first degree and it will have only one solution

Also when the discriminant is zero it will have only one solution

Discriminant is

$$4(m+1)^2 - 4m^2 \cdot 4 = 0$$

$$4(m^2 + 1 + 2m) - 16m^2 = 0$$

On simplifying and solving,

$$(m-1)(3m+1) = 0$$

$$m = 1, m = -\frac{1}{3}$$

Hence the three values of m for which the equation will have only one solution is

$$m = 0, m = 1, m = -\frac{1}{3}$$

10. **Solution**

$$A.P \quad a-d, a, a+d$$

$$GP \quad \frac{b}{g}, b, bg$$

$$a-d+a+a+d=3a$$

$$3a=126$$

$$a=42$$

$$a+b=76$$

$$b=34$$

$$a-d+\frac{b}{g}=85 \dots (1)$$

$$a+d+bg=84 \dots (2)$$

$$2a+\frac{b}{g}+bg=169$$

$$34g^2-85g+34=0$$

$$g = \frac{85 \pm \sqrt{85^2 - 4 \times 34 \times 34}}{2 \times 34}$$

$$g = 2 \quad \text{or} \quad \frac{1}{2}$$

$$\text{When } g = 2$$

$$42 - d + \frac{34}{2} = 85$$

$$d = -26$$

$$a = 42, \quad d = -26, \quad g = 2, \quad b = 34$$

AP

$$68, \quad 42, \quad 16$$

GP

$$17, \quad 34, \quad 68$$

$$m = 1, m = -\frac{1}{3}$$

11. **Solution**

$$f(x+1) = 4^{x+1}$$

$$f(x) = 4^x$$

$$f(x+1) - f(x) = 4^{x+1} - 4^x$$

$$= 4^x \cdot 4 - 4^x$$

$$= 4^x (3)$$

$$= 3f(x)$$

12. **Solution**

$$\log \frac{1 + \frac{3x + x^3}{1 + 3x^2}}{1 - \frac{3x + x^3}{1 + 3x^2}}$$

$$= \log \frac{1 + 3x^2 + 3x + x^3}{1 + 3x^2 - 3x - x^3}$$

$$= \log \frac{(1+x)^3}{(1-x)^3}$$

$$= 3 \log \frac{(1+x)}{(1-x)}$$

$$= 3f(x)$$

13. **Solution**

$$\sin(45 + 30) = \sin 45 \cos 30 + \cos 45 \sin 30$$

$$= \frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \times \frac{1}{2}$$

$$= \frac{\sqrt{3} + 1}{2\sqrt{2}}$$

$$= \frac{\sqrt{6} + \sqrt{2}}{4}$$

$$\cos(45 + 30) = \cos 45 \cos 30 - \sin 45 \sin 30$$

$$= \frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \times \frac{1}{2}$$

$$= \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

14. **Solution**

$$\frac{\sin 3\theta}{\sin \theta} - \frac{\cos 3\theta}{\cos \theta} = \frac{\sin 3\theta \cos \theta - \cos 3\theta \sin \theta}{\sin \theta \cos \theta}$$

$$= \frac{\sin(3\theta - \theta)}{\sin \theta \cos \theta}$$

$$= \frac{2 \sin 2\theta}{2 \sin \theta \cos \theta}$$

$$= \frac{2 \sin 2\theta}{\sin 2\theta} = 2$$

15. **Solution**

$$x^2 + 4(mx + 1)^2 = 1$$

$$x^2 + 4(m^2x^2 + 2mx + 1) = 1$$

$$x^2 + 4m^2x^2 + 8mx + 4 = 1$$

$$x^2(1 + 4m^2) + 8mx + 3 = 0$$

The line being a tangent, it touches the ellipse at two coincident points, and so Discriminant must be zero,

$$(8m)^2 - 4(3)(1 + 4m^2) = 0$$

$$64m^2 - 12 - 48m^2 = 0$$

$$16m^2 = 12$$

$$m^2 = \frac{12}{16}$$

$$m^2 = \frac{3}{4}$$

16. **Solution**

Divide the equation by

$$-\sqrt{3^2 + -4^2} = -5$$

$$\text{Hence, } -\frac{3}{5}x + \frac{4}{5}y - 4 = 0$$

$$\text{Where, } \cos \alpha = \frac{-3}{5} \text{ and } \sin \alpha = \frac{4}{5} \text{ and } p = 4$$

17. **Solution**

Multiply both numerator and denominator with $x - 7$. Then denominator becomes a perfect square and it is always positive

Now

$$(x + 3)(x - 7) \leq 0$$

Critical points are

$$(-3, 7)$$

$$\text{Hence, } -3 \leq x < 7$$

18. **Solution**

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^2 - ax + 4}{3x^2 - bx + 7} &= \lim_{x \rightarrow \infty} \frac{x^2(1 - \frac{a}{x} + \frac{4}{x^2})}{x^2(3 - \frac{b}{x} + \frac{7}{x^2})} \\ &= \frac{1}{3} \end{aligned}$$

19. **Solution**

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\tan x}{\sin 3x} &= \lim_{x \rightarrow 0} \frac{\sin x}{x} \times \frac{1}{\cos x} \times \frac{1}{\frac{3 \sin 3x}{3x}} \\ &= 1 \times 1 \times \frac{1}{3} = \frac{1}{3} \end{aligned}$$

Section C

20. **Solution**

Form a quadratic equation whose roots are

$$1 + i \text{ and } 1 - i$$

The equation is

$$x^2 - 2x + 2 = 0$$

The given expression

$$x^3 + x^2 - 4x + 13 = x(x^2 - 2x + 2) + 3(x^2 - 2x + 2) + 7$$

$$x^3 + x^2 - 4x + 13 = x(0) + (0) + 7$$

$$x^3 + x^2 - 4x + 13 = 7$$

21. **Solution**

$$x^2 - (\alpha + \beta)x + \alpha\beta - k^2 = 0$$

Discriminant of the above quadratic is

$\{(\alpha + \beta)\}^2 - 4(\alpha\beta - k^2) = (\alpha - \beta)^2 + k^2$ is always positive and hence the roots are real.

22. **Solution**

Let the roots be

$p\alpha$ and $q\beta$

Then

$$p\alpha + q\alpha = -\frac{n}{l} \dots (1)$$

$$pq\alpha^2 = \frac{n}{l}$$

$$\alpha = \frac{\sqrt{n}}{\sqrt{l}} \times \frac{1}{\sqrt{pq}} \dots (2)$$

Hence substituting equation 2 in equation 1

$$(p + q) \frac{\sqrt{n}}{\sqrt{l}} \times \frac{1}{\sqrt{pq}} + \frac{n}{l} = 0$$

On simplifying ,

$$\frac{\sqrt{p}}{\sqrt{q}} + \frac{\sqrt{q}}{\sqrt{p}} + \frac{\sqrt{n}}{\sqrt{l}} = 0$$

23. **Solution**

$$\begin{aligned} \lim_{x \rightarrow \pi} (\pi - x) \tan \frac{x}{2} &= \lim_{x \rightarrow \pi} \frac{2(\pi - x)}{2} \cot \frac{\pi - x}{2} \\ &= \lim_{x \rightarrow \pi} \frac{2(\pi - x)}{2} \frac{\cos \frac{\pi - x}{2}}{\sin \frac{\pi - x}{2}} \end{aligned}$$

$$\begin{aligned}
&= \lim_{x \rightarrow \pi} 2 \frac{\cos \frac{\pi-x}{2}}{\frac{\sin \frac{\pi-x}{2}}{\frac{\pi-x}{2}}} \\
&= \lim_{\frac{x-\pi}{2} \rightarrow 0} 2 \frac{\cos \frac{\pi-x}{2}}{\frac{\sin \frac{\pi-x}{2}}{\frac{\pi-x}{2}}} \\
&= \lim_{\frac{x-\pi}{2} \rightarrow 0} 2 \frac{\cos \frac{x-\pi}{2}}{\frac{\sin \frac{x-\pi}{2}}{\frac{x-\pi}{2}}} = 2 \quad \text{since the limit of } \frac{\sin \frac{x-\pi}{2}}{\frac{x-\pi}{2}} = 1
\end{aligned}$$

24. **Solution**

Let

$$a = x-1$$

$$b = x$$

$$c = x+1$$

Then

$$\begin{aligned}
&(x-1-i)((x-1+i)(x+1+i)(x+1-i)) = \{(x-1)^2 - i^2\} \{(x+1)^2 - i^2\} \\
&= \{(x-1)^2 + 1\} \{(x+1)^2 + 1\} \\
&= \{(x-1)(x+1)\}^2 + (x-1)^2 + (x+1)^2 + 1 \\
&= (x^2 - 1)^2 + (x-1)^2 + (x+1)^2 + 1 \\
&= x^4 + 1 \\
&= b^4 + 1
\end{aligned}$$

25. **Solution**

Multiply both Numerator and denominator with $(1-i)^2$ Then

$$\frac{(1+i)^n}{(1-i)^{n-2}} = \frac{(1+i)^n(1-i)^2}{(1-i)^n}$$

multiplying both Numerator & denominator with $(1+i)^n$

$$= \frac{(1+i)^n(-2i)(1+i)^n}{(1-i)^n(1+i)^n}$$

Simplifying

$$= \frac{\{(1+i)^2\}^n(-2i)}{(1-i^2)^n}$$

On expanding and simplifying

$$= \frac{2^n i^n (-2)i}{2^n}$$

$$= -2i^{n+1}$$

$$= \frac{2(i)^{n+1}}{i^2} = 2i^{n-1}$$

26. **Solution**

Let the point be $A(1,2)$ and $B(3,4)$

The mid-point of the line joining A and B is $C(2,3)$

$$\text{Slope of line AB} = \frac{4-2}{3-1} = 1$$

Let the required point be $D(\alpha, \beta)$

Then D must be a point on the line perpendicular to the line AB and passing through point C

$$\therefore \text{Slope of } CD = -1$$

Equation of CD

$$y-3 = -1(x-2)$$

$$x+y=5$$

Equation of AB

$$y-2 = 1(x-1)$$

$$x-y+1=0$$

The point $D(\alpha, \beta)$ must satisfy the equation

$$x+y=5$$

$$\therefore \alpha + \beta = 5 \dots (1)$$

The perpendicular distance from (α, β) to AB is

$$\frac{\alpha - \beta + 1}{\sqrt{2}} = \sqrt{2}$$

$$\alpha - \beta = 1 \dots (2)$$

Solving equations 1 and 2

$$\alpha = 3, \beta = 2$$
