

Cross Product of vectors

Q.1. Given $\vec{a} = i - 2j + k$, $\vec{b} = 2i + j + k$ and $\vec{c} = i + 2j - k$. Find : $\vec{a} \times (\vec{b} \times \vec{c})$.

Solution : 1

$$\begin{aligned}\vec{a} \times (\vec{b} \times \vec{c}) &= (i - 2j + k) \times (2i + j + k) \times (i + 2j - k) \\ &= (a \cdot c)b - (a \cdot b)c \\ &= (1 - 4 - 1)(2i + j + k) - (2 - 2 + 1)(i + 2j - k) \\ &= -8i - 4j - 4k - i - 2j + k \\ &= -9i - 6j - 3k.\end{aligned}$$

Q.2. Find a unit vector perpendicular to the vectors $4i + 3j + k$ and $2i - j + 2k$. Determine the sine angle between these two vectors.

Solution : 2

Unit vector perpendicular to $\vec{a}$ and $\vec{b}$ is given by :

$$[\vec{a} \times \vec{b}] / |\vec{a} \times \vec{b}|$$

Therefore, unit vector perpendicular to $4i + 3j + k$ and $2i - j + k$ is

$$\begin{aligned}&= [(4i + 3j + k) \times (2i - j + 2k)] / |(4i + 3j + k) \times (2i - j + 2k)| \\ &= [7i - 6j - 10k] / \sqrt{49 + 36 + 100} \\ &= (7/\sqrt{185})i - (6/\sqrt{185})j - 10/\sqrt{185}k.\end{aligned}$$

Q.3. The vectors $-2i + 4j + 4k$ and $-4i - 2k$ represent the diagonals BD and AC of a parallelogram ABCD. Find the area of the parallelogram.

Solution : 3

$$\begin{aligned}
\text{Area of parallelogram} &= \frac{1}{2} |\vec{BD} \times \vec{AC}| = \frac{1}{2} |(-2\mathbf{i} + 4\mathbf{j} + 4\mathbf{k}) \times (-4\mathbf{i} - 2\mathbf{k})| \\
&= \frac{1}{2} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 4 & 4 \\ -4 & 0 & -2 \end{vmatrix} \\
&= \frac{1}{2} |\mathbf{i}(-8) - \mathbf{j}(4 + 16) + \mathbf{k}(16)| \\
&= \frac{1}{2} |-8\mathbf{i} - 20\mathbf{j} + 16\mathbf{k}| \\
&= \frac{1}{2} \sqrt{8^2 + 20^2 + 16^2} \\
&= \frac{1}{2} \sqrt{720} = \frac{1}{2} \times 12\sqrt{5} = 6\sqrt{5}.
\end{aligned}$$

Q.4. Find the unit vector perpendicular to the two vectors : $\mathbf{i} \rightarrow + \mathbf{j} \rightarrow - \mathbf{k} \rightarrow$ and $2\mathbf{i} \rightarrow + 3\mathbf{j} \rightarrow + \mathbf{k} \rightarrow$.

Solution : 4

Let $\vec{a} = \mathbf{i} \rightarrow + \mathbf{j} \rightarrow - \mathbf{k} \rightarrow$ and $\vec{b} = 2\mathbf{i} \rightarrow + 3\mathbf{j} \rightarrow + \mathbf{k} \rightarrow$

Unit vector perpendicular to $\vec{a}$ and $\vec{b} = [\vec{a} \times \vec{b}] / |\vec{a} \times \vec{b}|$

$$[\vec{a} \times \vec{b}] = \mathbf{i} \rightarrow (2 + 3) - \mathbf{j} \rightarrow (1 + 2) + \mathbf{k} \rightarrow (3 - 4) = 5\mathbf{i} \rightarrow - 3\mathbf{j} \rightarrow - \mathbf{k} \rightarrow$$

$$\text{and } |\vec{a} \times \vec{b}| = |5\mathbf{i} \rightarrow - 3\mathbf{j} \rightarrow - \mathbf{k} \rightarrow| = \sqrt{25 + 9 + 1} = \sqrt{35}]$$

Therefore, unit vector perpendicular to $\vec{a}$ and $\vec{b}$

$$= [5\mathbf{i} \rightarrow - 3\mathbf{j} \rightarrow - \mathbf{k} \rightarrow] / \sqrt{35}.$$

Q.5. Find the area of the parallelogram ABCD whose diagonals AC and BD are represented by the vectors $3\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ and $\mathbf{i} - 3\mathbf{j} - 4\mathbf{k}$ respectively.

Solution : 5

We are given , $\vec{AC} = 3\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ and $\vec{BD} = \mathbf{i} - 3\mathbf{j} - 4\mathbf{k}$.

$$\text{Area of parallelogram} = \frac{1}{2} |\vec{AC} \times \vec{BD}|$$

$$\vec{AC} \times \vec{BD} = (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) \times (\mathbf{i} - 3\mathbf{j} - 4\mathbf{k})$$

$$= i(4 - 6) - j(12 + 2) + k(-9 - 1)$$

$$= -2i - 14j - 10k$$

$$|\vec{AC} \times \vec{BD}| = \sqrt{(4 + 196 + 100)} = \sqrt{(300)} = 10\sqrt{3}$$

$$\text{Area of parallelogram} = \frac{1}{2} |\vec{AC} \times \vec{BD}| = \frac{1}{2} \times 10\sqrt{3} = 5\sqrt{3}.$$

Q.6. Find the area of a parallelogram whose diagonals are determined by the vectors :
 $\vec{a} = 3i + j - 2k$ and $\vec{b} = i - 3j + 4k$.

Solution : 6

$$\text{Area} = \frac{1}{2} |\vec{d}_1 \times \vec{d}_2|$$

$$|\vec{i} \vec{j} \vec{k}|$$

$$= \frac{1}{2} |3 \ 1 \ -2|$$

$$|1 \ -3 \ 4|$$

$$= \frac{1}{2} |i(4 - 6) - j(12 + 2) + k(-9 - 1)|$$

$$= \frac{1}{2} |(-2i - 14j - 10k)|$$

$$= \frac{1}{2} \sqrt{(4 + 196 + 100)}$$

$$= \frac{1}{2} \sqrt{(300)} = 5\sqrt{3} \text{ sq. units.}$$


Q.7. If $\vec{a}$, $\vec{b}$ and $\vec{c}$ represents the position vectors of the points with co-ordinates (2, -10, 2), (3, 1, 2) and (2, 1, 3) respectively, find the value of $\vec{a} \times (\vec{b} \times \vec{c})$.

Solution : 7

$$\vec{a} \times (\vec{b} \times \vec{c}) = (2i - 10j + 2k) \times \{(3i + j + 2k) \times (2i + j + 3k)\}$$

$$= (2i - 10j + 2k) \times (i - 5j + k) = 0.$$

Q.8. Using vectors, show that the medians of a triangle meet at a point.

Solution : 8

Fig

Let D, E and F be mid-points of sides BC, CA and AB of ΔABC and O the point of intersection of medians AD and BE. Let position vectors of A, B and C be $\vec{a}$, $\vec{b}$ and $\vec{c}$ respectively.

$$\vec{OD} = \vec{OB} - \vec{DB}$$

$$= \vec{OB} - \frac{1}{2}\vec{CB}$$

$$= \vec{OB} - \frac{1}{2}(\vec{OB} - \vec{OC})$$

$$= \frac{1}{2}(\vec{OB} + \vec{OC})$$

$$= \frac{1}{2}(\vec{b} + \vec{c})$$

$$\text{Similarly, } \vec{OE} = \frac{1}{2}(\vec{a} + \vec{c})$$

$$\text{and } \vec{OF} = \frac{1}{2}(\vec{a} + \vec{b})$$

$\vec{OA}$ and $\vec{OD}$ being in opposite direction,

$$\vec{OA} \times \vec{OD} = 0$$

$$\text{Or, } \vec{a} \times \frac{1}{2}(\vec{b} + \vec{c}) = 0$$

$$\text{Or, } \frac{1}{2}[\vec{a} \times (\vec{b} + \vec{c})] = 0 \text{ ----- (1)}$$

$$\text{Similarly, } \frac{1}{2}[\vec{b} \times (\vec{a} + \vec{c})] = 0 \text{ ----- (2)}$$

Adding (1) and (2), we get

$$\frac{1}{2}[\vec{a} \times (\vec{b} + \vec{c})] + \frac{1}{2}[\vec{b} \times (\vec{a} + \vec{c})] = 0$$

$$\text{Or, } \frac{1}{2}[\vec{a} \times \vec{b} + \vec{a} \times \vec{c} + \vec{b} \times \vec{a} + \vec{b} \times \vec{c}] = 0$$

$$\text{Or, } \vec{a} \times \vec{c} + \vec{b} \times \vec{c} = 0 \text{ [} \vec{a} \times \vec{b} = -\vec{b} \times \vec{a} \text{]}$$

$$\text{Or, } (\vec{a} + \vec{b}) \times \vec{c} = 0$$

$$\text{Or, } \frac{1}{2}(\vec{a} + \vec{b}) \times \vec{c} = 0$$

$$\text{Or, } \vec{OC} \times \vec{OF} = 0$$

Thus we see that $\vec{OC}$ and $\vec{OF}$ are in opposite in direction i.e., median CF also passes through the point 'O' which is the intersection point of median AD and BE. Therefore, medians of a triangle meet at a point.

Q.9. Given that $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} + 2\hat{j} - \hat{k}$. Find the vector $\vec{a} \times (\vec{b} \times \vec{c})$.

Solution : 9

We have, $\vec{b} \times \vec{c} = (2\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} + 2\hat{j} - \hat{k})$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 1 \\ 1 & 2 & -1 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & 1 & 1 \\ 1 & 2 & -1 \end{vmatrix}$$

$$= \hat{i} [(1)(-1) - (1)(2)] - \hat{j} [(2)(-1) - (1)(1)] + \hat{k} [(2)(2) - (1)(1)]$$

$$= -3\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\hat{i} - 2\hat{j} + \hat{k}) \times (-3\hat{i} + 3\hat{j} + 3\hat{k})$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 1 \\ -3 & 3 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & -2 & 1 \\ -3 & 3 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & -2 & 1 \\ -3 & 3 & 3 \end{vmatrix}$$

$$= \hat{i} [(-2)(3) - (1)(3)] - \hat{j} [(1)(3) - (1)(-3)] + \hat{k} [(1)(3) - (-2)(-3)]$$


$$= \hat{i} (-9) - \hat{j} (6) + \hat{k} (-3)$$

$$= -9\hat{i} - 6\hat{j} - 3\hat{k}$$

$$= -9\hat{i} - 6\hat{j} - 3\hat{k}$$

Q.10. If D, E, F are the mid-points of the sides BC, CA, AB respectively of a triangle ABC. Show that the area of triangle DEF = $\frac{1}{4}$ [area of Δ ABC].

Solution : 10



Fig

Let A be the origin and AB and AC represents $\vec{b}$ and $\vec{c}$.

Then, $\vec{DE} = -\frac{1}{2}\vec{b}$; $\vec{DF} = -\frac{1}{2}\vec{c}$.

Vector area of $\triangle DEF = \frac{1}{2}\vec{DE} \times \vec{DF}$

$$= \frac{1}{2} \left[-\frac{1}{2}\vec{b} \times -\frac{1}{2}\vec{c} \right]$$

$$= \frac{1}{8}(\vec{b} \times \vec{c})$$

$$= \frac{1}{4} \left(\frac{1}{2} \vec{b} \times \vec{c} \right)$$

$$= \frac{1}{4} \left[\frac{1}{2} \vec{AB} \times \vec{AC} \right]$$

$$= \frac{1}{4} [\text{Vector area of } \triangle ABC]$$

Q.11. Show that : $\vec{i} \times (\vec{a} \times \vec{i}) + \vec{j} \times (\vec{a} \times \vec{j}) + \vec{k} \times (\vec{a} \times \vec{k}) = 2\vec{a}$ where, $\vec{a} = a_1\vec{i} + a_2\vec{j} + a_3\vec{k}$.

Solution : 11

We have, $\vec{i} \times (\vec{a} \times \vec{i}) + \vec{j} \times (\vec{a} \times \vec{j}) + \vec{k} \times (\vec{a} \times \vec{k})$

$$= [(\vec{i} \cdot \vec{i})\vec{a} - (\vec{i} \cdot \vec{a})\vec{i}] + [(\vec{j} \cdot \vec{j})\vec{a} - (\vec{j} \cdot \vec{a})\vec{j}] + [(\vec{k} \cdot \vec{k})\vec{a} - (\vec{k} \cdot \vec{a})\vec{k}]$$

$$= \vec{a} - (\vec{i} \cdot \vec{a})\vec{i} + \vec{a} - (\vec{j} \cdot \vec{a})\vec{j} + \vec{a} - (\vec{k} \cdot \vec{a})\vec{k}$$

$$= 3\vec{a} - [(\vec{i} \cdot \vec{a})\vec{i} + (\vec{j} \cdot \vec{a})\vec{j} + (\vec{k} \cdot \vec{a})\vec{k}]$$

$$= 3\vec{a} - [(a_1\vec{i}) + (a_2\vec{j}) + (a_3\vec{k})]$$

$$= 3\vec{a} - \vec{a}$$

$$= 2\vec{a}.$$