

Topics : Vector, Three Dimensional Geometry

Type of Questions

M.M., Min.

Single choice Objective (no negative marking) Q.1, 2, 3	(3 marks, 3 min.)	[9, 9]
Multiple choice objective (no negative marking) Q.4, 5	(5 marks, 4 min.)	[10, 8]
Subjective Questions (no negative marking) Q.6 to Q.8	(4 marks, 5 min.)	[12, 15]
Match the Following (no negative marking) Q.9	(8 marks, 8 min.)	[8, 8]

1. If the line  $\frac{x-2}{5} = \frac{y+10}{2} = \frac{z+3}{2}$  meets the curve

$xy = a^2, z = 1$ , then number of values of  $a$  is

- (A) 0 (B) 1 (C) 2 (D) More than 2

2. If  $|\vec{a}| = 2, |\vec{b}| = 3$  and  $\vec{a} \cdot \vec{b} = 0$  then  $\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times \vec{b})))$  is equal to

- (A)  $48 \vec{b}$  (B)  $-48 \vec{b}$  (C)  $48 \vec{a}$  (D)  $16 \vec{b}$

3. The equation of a line  $4x - 4y - z + 11 = 0 = x + 2y - z - 1$  can be put as

- (A)  $\frac{x}{2} = \frac{y-2}{1} = \frac{z-3}{4}$  (B)  $\frac{x-2}{2} = \frac{y-2}{1} = \frac{z}{4}$  (C)  $\frac{x-2}{2} = \frac{y}{1} = \frac{z-3}{4}$  (D) None of these

4. A ray M is sent along the line  $\frac{x-0}{2} = \frac{y-2}{2} = \frac{z-1}{0}$  and is reflected by the plane  $x = 0$  at point A. The reflected ray is again reflected by the plane  $x + 2y = 0$  at point B. The initial ray and final reflected ray meets at point J. Then

- (A) the co-ordinates of point B is (4, -2, 1) (B) the co-ordinates of point J is (-3, -1, 1)  
(C) the centroid of  $\triangle ABJ$  is (0, 0, 0) (D) the co-ordinates of point J is (2, -1, 1)

5. The line which intersects each of the two lines  $L_1 : 2x + y - 1 = 0 = 3x - 2y + z$ ,

$L_2 : 3x - y - z + 1 = 0 = 4x + y + 5z - 3$  and is parallel to the line  $\frac{x}{1} = \frac{y}{2} = \frac{z}{-1}$

- (A) has direction ratio (1, 2, -1)  
(B) has equation  $8x - 3y + 2z - 1 = 0 = 5x + 3y + 11z - 7$

(C) having angle with  $L_2$  equal to  $\cos^{-1} \left( \sqrt{\frac{3}{7}} \right)$

(D) is perpendicular to the plane  $3x + 6y - 3z = 7$

6. Let image of the line  $\frac{x-1}{3} = \frac{y-3}{5} = \frac{z-4}{2}$  in the plane  $2x - y + z + 3 = 0$  be L. A plane  $7x + By + Cz + D = 0$  is such that it contains the line L and perpendicular to the plane  $2x - y + z + 3 = 0$  then find the value of  $B + C + D$

7. P is a point and PM, PN are perpendicular from P to the ZX and XY planes. If OP makes angle  $\theta, \alpha, \beta, \gamma$  with the plane OMN and the XY, YZ, ZX plane respectively, then prove that  $\operatorname{cosec}^2 \theta = \operatorname{cosec}^2 \alpha + \operatorname{cosec}^2 \beta + \operatorname{cosec}^2 \gamma$ .

8. Find the sum of n terms of the series  $\frac{3}{1.2} \cdot \frac{1}{2} + \frac{4}{2.3} \cdot \frac{1}{2^2} + \frac{5}{3.4} \cdot \frac{1}{2^3} + \frac{6}{4.5} \cdot \frac{1}{2^4} + \dots$

9. **Match the column**

**Column - I**

**Column - II**

- (A) If  $\vec{a}, \vec{b}, \vec{c}$  non-coplanar vectors, then  $(\vec{a} + \vec{b} + \vec{c}) \cdot ((\vec{a} + \vec{b}) \times (2\vec{a} + \vec{b}))$

(p)  $\frac{1}{4} a^2 b^2$

is equal to

- (B) If  $\vec{b}$  and  $\vec{c}$  are any two non-collinear perpendicular unit vectors and  $\vec{a}$  is (q)

–  $[\vec{a} \vec{b} \vec{c}]$

any vector, then  $(\vec{a} \cdot \vec{b})\vec{b} + (\vec{a} \cdot \vec{c})\vec{c} + \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{|\vec{b} \times \vec{c}|^2} (\vec{b} \times \vec{c})$  is equal to

- (C) If  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar vectors then  $[\vec{a} + \vec{b} + \vec{c} \quad \vec{a} - \vec{c} \quad \vec{a} - \vec{b}]$

(r)  $\vec{a}$

is equal to

- (D) Let  $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$ ,  $\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$ ,  $\vec{c} = c_1 \hat{i} + c_2 \hat{j} + c_3 \hat{k}$

(s)  $-3[\vec{a} \vec{b} \vec{c}]$

non-zero vectors such that  $\vec{c}$  is a unit vector perpendicular to both  $\vec{a}$  and  $\vec{b}$  (and angle between  $\vec{a}$  and  $\vec{b}$  is  $(\pi/6)$ ), then

$\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}^2$  is equal to

# Answers Key

1. A

2. D

3. A

4. AB

5. ABD

6. 30

8.  $1 - \frac{1}{2^n(n+1)}$

9.  $(A) \rightarrow q ; (B) \rightarrow r ; (C) \rightarrow s ; (D) \rightarrow p$