Class 11

Important Formulas

Limits and Derivatives

Limits:

1.
$$\lim_{x \to a} f(x) \text{ exists} \Leftrightarrow \lim_{x \to a^{-}} f(x) = \lim_{x \to a^{+}} f(x)$$

- 2. For a function f(x) and a real number a, $\lim_{x \to a} f(x)$ and f(a) may not be same. In fact:
 - (i) $\lim_{x \to a} f(x)$ exists but f(a) (the value of f(x) at x = a) may not exists
 - (ii) The value f(a) exists but $\lim_{x \to a} f(x)$ does not exist
 - (iii) $\lim_{x \to a} f(x)$ and f(a) both exist but are unequal
 - (iv) $\lim_{x \to a} f(x)$ and f(a) both exist and are equal.
- 3. Let $\lim_{x \to a} f(x) = l$ and $\lim_{x \to a} g(x) = m$. If l and m both exist, then
 - (i) $\lim_{x \to a} k f(x) = k \lim_{x \to a} f(x)$
 - (ii) $\lim_{x \to a} (f \pm g)(x) = \lim_{x \to a} f(x) \pm \lim_{x \to a} g(x) = l + m$
 - (iii) $\lim_{x \to a} (fg)(x) = \lim_{x \to a} f(x) \lim_{x \to a} g(x) = lm$

(iv)
$$\lim_{x \to a} \left(\frac{f}{g} \right) (x) = \frac{\lim_{x \to a} f(x)}{\lim_{x \to a} g(x)} = \frac{l}{m}$$

(v)
$$\lim_{x \to a} \{f(a)\}^{g(x)} = l^m$$

- 4. Following are some standard limits:
 - (i) $\lim_{x \to a} \frac{x^n a^n}{x a} = na^{n-1}$

(ii)
$$\lim_{x \to 0} \frac{\sin x}{x} = 1$$

(iii) $\lim_{x \to 0} \frac{\tan x}{x} = 1$

(iv)
$$\lim_{x \to a} \frac{\sin(x-a)}{x-a} = 1$$

(v) $\lim_{x \to a} \frac{\tan (x-a)}{x-a} = 1$

(vi)
$$\lim_{x \to 0} \frac{\log (1+x)}{x} = 1$$

(vii)
$$\lim_{x \to 0} \frac{a^x - 1}{x} = \log_e a, a \neq 0, a > 1$$

(viii)
$$\lim_{x \to 0} \frac{e^x - 1}{x} = 1$$

Derivatives:

1. A function f(x) is differentiable at x = c iff $\lim_{x \to c} \frac{f(x) - f(c)}{x - c}$ exists finitely.

This limit is called the derivative or differentiation of f(x) at x = c and is denoted by f'(c).

- 2. Geometrically the derivative of a function f(x) at a point x = c is the slope of the tangent to the curve y = f(x) at the point (c, f(c)).
- 3. If f(x) is a differentiable function, then $\lim_{h\to 0} \frac{f(x+h)-f(x)}{h}$ is called the differentiation of f(x) or differentiation of f(x) with respect to x.
- 4. Mechanically, $\frac{d}{dx}(f(x))$ measures the rate of change of f(x) with respect to x.
- 5. Following are some standard derivatives:

(i)
$$\frac{d}{dx}(x^n) = n x^{n-1}$$

(ii)
$$\frac{d}{dx}(a^x) = a^x \log_e a, a > 0, a \ne 1$$

(iii)
$$\frac{d}{dx}(e^x) = e^x$$

(iii)
$$\frac{dx}{dx} (e^x) = e^x$$
 (iv) $\frac{d}{dx} (\log_e x) = \frac{1}{x}$

(v)
$$\frac{d}{dx} (\sin x) = \cos x$$

(vi)
$$\frac{d}{dx}(\cos x) = -\sin x$$

(vii)
$$\frac{d}{dx}$$
 (tan x) = $\sec^2 x$

(viii)
$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

(ix)
$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

(x)
$$\frac{d}{dx}(\csc x) = -\csc x \cot x$$

- 5. Following are the fundamental rules for differentiation:
 - (i) Differentiation of a constant function is zero i.e., $\frac{d}{dx}(c) = 0$
- (ii) Differentiation of a constant and a function is equal to constant times the differentiation of the function.
- (iii) If f(x) and g(x) are differentiable functions, then

(a)
$$\frac{d}{dx} \left\{ f(x) \pm g(x) \right\} = \frac{d}{dx} \left(f(x) \right) \pm \frac{d}{dx} \left(g(x) \right)$$

(b)
$$\frac{d}{dx} \left\{ f(x) \times g(x) \right\} = g(x) \times \frac{d}{dx} \left(f(x) \right) + f(x) \times \frac{d}{dx} \left(g(x) \right)$$

(b)
$$\frac{d}{dx} \{f(x) \times g(x)\} = g(x) \times \frac{d}{dx} (f(x)) + f(x) \times \frac{d}{dx} (g(x)) \}$$

(c) $\frac{d}{dx} \{\frac{f(x)}{g(x)}\} = \frac{g(x) \times \frac{d}{dx} (f(x)) - f(x) \times \frac{d}{dx} \{g(x)\}}{\{g(x)\}^2}$