


Trigonometry

- **Trigonometric Ratio**



$$\sin \theta = \frac{\text{Opposite side}}{\text{Hypotenuse}} = \frac{AB}{AC}$$

$$\cos \theta = \frac{\text{Adjacent side}}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$\tan \theta = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{AB}{BC}$$

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta} = \frac{\text{Hypotenuse}}{\text{Opposite side}} = \frac{AC}{AB}$$

$$\sec \theta = \frac{1}{\cos \theta} = \frac{\text{Hypotenuse}}{\text{Adjacent side}} = \frac{AC}{BC}$$

$$\cot \theta = \frac{1}{\tan \theta} = \frac{\text{Adjacent side}}{\text{Opposite side}} = \frac{BC}{AB}$$


$$\text{Also, } \tan \theta = \frac{\sin \theta}{\cos \theta}, \cot \theta = \frac{\cos \theta}{\sin \theta}$$

If one of the trigonometric ratios of an acute angle is known, then the remaining trigonometric ratios of the angle can be calculated.

Example:

If $\sin \theta = \frac{7}{25}$, then find the value of $\sec \theta(1 + \tan \theta)$.

Solution:



It is given that $\sin \theta = \frac{7}{25}$

$$\sin \theta = \frac{AB}{AC} = \frac{7}{25}$$

$\Rightarrow AB = 7x$ and $AC = 25x$, where x is some positive integer

By applying Pythagoras theorem in $\triangle ABC$, we get:

$$AB^2 + BC^2 = AC^2$$

$$\Rightarrow (7x)^2 + BC^2 = (25x)^2$$

$$\Rightarrow 49x^2 + BC^2 = 625x^2$$

$$\Rightarrow BC^2 = 625x^2 - 49x^2$$

$$\Rightarrow BC = \sqrt{576}x = 24x$$

$$\therefore \sec \theta = \frac{\text{Hypotenuse}}{\text{Adjacent side}} = \frac{25}{24}$$

$$\tan \theta = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{7}{24}$$

$$\therefore \sec \theta(1 + \tan \theta) = \frac{25}{24} \left(1 + \frac{7}{24} \right) = \frac{25}{24} \times \frac{31}{24} = \frac{775}{576}$$

- Use trigonometric ratio in solving problem.

Example:

If $\tan \theta = \frac{3}{5}$, then find the value of $\frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta}$

Solution:

$$\frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta}$$

Take $\cos \theta$ common from numerator and denominator both

$$= \frac{\frac{\sin \theta}{\cos \theta} + 1}{\frac{\sin \theta}{\cos \theta} - 1}$$

$$= \frac{\tan \theta + 1}{\tan \theta - 1}$$

$$= \frac{\frac{3}{5} + 1}{\frac{3}{5} - 1}$$

$$= \frac{\frac{3+5}{5}}{\frac{3-5}{5}}$$

$$= \frac{8}{-2}$$

$$= -4$$

- Trigonometric Ratios of some specific angles


q	0	30°	45°	60°	90°
$\sin q$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos q$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan q$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined

cosec q	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
sec q	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
cot q	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

Example 1:

$\triangle ABC$ is right-angled at B and $AB = 6$ m, **$BC = \sqrt{12}$** m. Find the measure of $\angle A$ and $\angle C$.

Solution:



$AB = 6$ m,

$BC = \sqrt{12}$ m $= 2\sqrt{3}$ m

$$\tan C = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{AB}{BC} = \frac{6}{2\sqrt{3}} = \sqrt{3}$$

$$\Rightarrow \tan C = \tan 60^\circ \quad \left[\because \tan 60^\circ = \sqrt{3} \right]$$

$$\Rightarrow \angle C = 60^\circ$$

$$\therefore \angle A = 180^\circ - (90 + 60) = 30^\circ$$

Example 2:

Evaluate the expression

$$4(\cos^3 60^\circ - \sin^3 30^\circ) + 3(\sin 30^\circ - \cos 60^\circ)$$

Solution:

$$\begin{aligned}
& 4(\cos^3 60^\circ - \sin^3 30^\circ) + 3(\sin 30^\circ - \cos 60^\circ) \\
&= 4\left[\left(\frac{1}{2}\right)^3 - \left(\frac{1}{2}\right)^3\right] + 3\left(\frac{1}{2} - \frac{1}{2}\right) \\
&= 4 \times 0 + 3 \times 0 = 0 + 0 = 0
\end{aligned}$$

- **Trigonometric Identities**

1. $\cos^2 A + \sin^2 A = 1$
2. $1 + \tan^2 A = \sec^2 A$
3. $1 + \cot^2 A = \operatorname{cosec}^2 A$

Example:

If $\cos \theta = \frac{5}{7}$, find the value of $\cot \theta + \operatorname{cosec} \theta$

Solution:

We have, $\cos \theta = \frac{5}{7}$

$$\text{Now, } \sin^2 \theta + \cos^2 \theta = 1$$

$$\therefore \sin \theta = \sqrt{1 - \cos^2 \theta}$$

$$= \sqrt{1 - \left(\frac{5}{7}\right)^2}$$

$$= \sqrt{\frac{49-25}{49}} = \frac{2\sqrt{6}}{7}$$

$$\therefore \operatorname{cosec} \theta = \frac{7}{2\sqrt{6}}$$

$$\text{Also, } \cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$= \frac{\frac{5}{7}}{\frac{2\sqrt{6}}{7}} = \frac{5}{2\sqrt{6}}$$


$$\therefore \cot \theta + \operatorname{cosec} \theta = \frac{5}{2\sqrt{6}} + \frac{7}{2\sqrt{6}}$$

$$= \frac{12}{2\sqrt{6}} = \frac{6}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}}$$

$$= \sqrt{6}$$

- **Some Applications of Trigonometry**


- **Line of sight:** It is the line drawn from the eye of an observer to a point on the object viewed by the observer.
- **Angle of Elevation:**



Let P be the position of the eye of the observer. Let Q be the object above the horizontal line PR.

Angle of elevation of the object Q with respect to the observer P is the angle made by the line of sight PQ with the horizontal line PR. That is, $\angle QPR$ is the angle of elevation.

◦ Angle of Depression



Let P be the position of the eye of the observer. Let Q be the object below the horizontal line PX.

Angle of depression of the object Q with respect to the observer P is the angle made by the line of sight PQ with the horizontal line PX. That is, $\angle XPQ$ is the angle of depression. It can be seen that


$$\angle PQR = \angle XPQ \quad [\text{Alternate interior angles}]$$

The height or length of an object or the distance between two distant objects can be calculated by using trigonometric ratios.

Example:

The angle of elevation of the top of a tower from the foot of a building is 60° and the angle of elevation of the top of the building from the foot of the tower is 30° . If the building is 16 m tall, then what is the height of the tower?

Solution:



Let AB and CD be the building and the tower respectively.
 It is given that, angles of elevation $\angle ADB = 30^\circ$, $\angle CBD = 60^\circ$
 In $\triangle ABD$,

$$\begin{aligned}\frac{AB}{BD} &= \tan 30^\circ \\ \Rightarrow \frac{16}{BD} &= \frac{1}{\sqrt{3}} \\ \Rightarrow BD &= 16\sqrt{3} \text{ m} \quad \text{---(1)}\end{aligned}$$

Now, in $\triangle CBD$

$$\begin{aligned}\frac{CD}{BD} &= \tan 60^\circ \\ \Rightarrow \frac{CD}{16\sqrt{3}} &= \sqrt{3} \quad \text{[using (1)]} \\ \Rightarrow CD &= 16\sqrt{3} \times \sqrt{3} \text{ m} = 48 \text{ m}\end{aligned}$$

Thus, the height of the tower is 48 m.


Example:

Two wells are located on the opposite sides of a 18 m tall building. As observed from the top of the building, the angles of depression of the two

wells are 30° and 45° . Find the distance between the wells. [Use $\sqrt{3} = 1.732$]

Solution:

The given situation can be represented as



Here, PQ is the building. A and B are the positions of the two wells such that:

$$\angle XPB = 30^\circ, \angle XPA = 45^\circ$$

$$\text{Now, } \angle PAQ = \angle XPA = 45^\circ$$

$$\angle PBQ = \angle XPB = 30^\circ$$

In $\triangle PAQ$, we have

$$\frac{PQ}{AQ} = \tan 45^\circ$$

$$\Rightarrow \frac{18}{AQ} = 1$$

$$\Rightarrow AQ = 18\text{m}$$

In $\triangle PBQ$, we have

$$\frac{PQ}{QB} = \tan 30^\circ$$

$$\Rightarrow \frac{18}{QB} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow QB = 18\sqrt{3}$$

$$\begin{aligned}\therefore AB &= AQ + QB = (18 + 18\sqrt{3}) \text{ m} \\ &= 18(1 + \sqrt{3}) \text{ m} \\ &= 18(1 + 1.732) \text{ m} \\ &= 18 \times 2.732 \text{ m} \\ &= 49.176 \text{ m}\end{aligned}$$

- Trigonometric Ratios of Complementary Angles

$$\begin{array}{ll}\sin(90^\circ - \theta) = \cos \theta & \cos(90^\circ - \theta) = \sin \theta \\ \tan(90^\circ - \theta) = \cot \theta & \cot(90^\circ - \theta) = \tan \theta \\ \operatorname{cosec}(90^\circ - \theta) = \sec \theta & \sec(90^\circ - \theta) = \operatorname{cosec} \theta\end{array}$$

Where θ is an acute angle.

Example 1: Evaluate the expression

$$\sin 28^\circ \sin 30^\circ \sin 54^\circ \sec 36^\circ \sec 62^\circ$$

Solution:

$$\begin{aligned}& \sin 28^\circ \sin 30^\circ \sin 54^\circ \sec 36^\circ \sec 62^\circ \\ &= (\sin 28^\circ \sec 62^\circ)(\sin 54^\circ \sec 36^\circ) \sin 30^\circ \\ &= \{\sin 28^\circ \operatorname{cosec}(90^\circ - 62^\circ)\} \{\sin 54^\circ \operatorname{cosec}(90^\circ - 36^\circ)\} \sin 30^\circ \\ &= (\sin 28^\circ \operatorname{cosec} 28^\circ)(\sin 54^\circ \operatorname{cosec} 54^\circ) \sin 30^\circ \\ &= \left(\sin 28^\circ \frac{1}{\sin 28^\circ} \right) \left(\sin 54^\circ \frac{1}{\sin 54^\circ} \right) \times \frac{1}{2} \\ &= \frac{1}{2}\end{aligned}$$

Example 2: Evaluate the expression

$$4\sqrt{3}(\sin 40^\circ \sec 30^\circ \sec 50^\circ) + \frac{\sin^2 34^\circ + \sin^2 56^\circ}{\sec^2 31^\circ - \cot^2 59^\circ}$$

Solution:

$$\begin{aligned} & 4\sqrt{3}(\sin 40^\circ \sec 30^\circ \sec 50^\circ) + \frac{\sin^2 34^\circ + \sin^2 56^\circ}{\sec^2 31^\circ - \cot^2 59^\circ} \\ &= 4\sqrt{3}[\sec 30^\circ (\sin 40^\circ \sec 50^\circ)] + \frac{\sin^2 34^\circ + \sin^2 (90^\circ - 56^\circ)}{\sec^2 31^\circ - \tan^2 (90^\circ - 59^\circ)} \\ & \quad [\because \cos(90^\circ - \theta) = \sin \theta, \tan(90^\circ - \theta) = \cot \theta] \\ &= 4\sqrt{3}[\sec 30^\circ \sin 40^\circ \operatorname{cosec}(90^\circ - 50^\circ)] + \frac{\sin^2 34^\circ + \cos^2 34^\circ}{\sec^2 31^\circ - \tan^2 31^\circ} \\ &= 4\sqrt{3}\left[\frac{2}{\sqrt{3}} \sin 40^\circ \operatorname{cosec} 40^\circ\right] + \frac{1}{1} \\ &= 8 + 1 = 9 \end{aligned}$$