

POLYNOMIALS

2

CHAPTER

CONTENTS

- Introduction
- Constants & Variables
- Algebraic Expression
- Factors & Coefficients
- Degree of a Polynomial
- Types of Polynomial & Polynomial in one variable
- Remainder Theorem
- Values & Zeroes of a Polynomial
- Geometric Meaning of the Zeroes of a Polynomial
- Relation between Zeroes & Coefficients
- Formation of Quadratic Polynomial

➤ INTRODUCTION

Algebra is that branch of mathematics which treats the relation of numbers.

➤ CONSTANTS AND VARIABLES

In algebra, two types of symbols are used: constants and variable (literals).

◆ Constant :

It is a symbol whose value always remains the same, whatever the situation be.

For example: 5, -9, $\frac{3}{8}$, π , $\frac{7}{15}$, etc.

◆ Variable :

It is a symbol whose value changes according to the situation.

For example : x, y, z, ax, a + x, 5y, -7x, etc.

➤ ALGEBRAIC EXPRESSION

- (a) An algebraic expression is a collection of terms separated by plus (+) or minus (-) sign. For example : $3x + 5y$, $7y - 2x$, $2x - ay + az$, etc.
- (b) The various parts of an algebraic expression that are separated by '+' or '-' sign are called terms.

For example :

Algebraic expression	No. of terms	Terms
(i) $-32x$	1	$-32x$
(ii) $2x + 3y$	2	$2x$ and $3y$
(iii) $ax - 5y + cz$	3	ax , $-5y$ and cz
(iv) $\frac{3}{x} + \frac{y}{7} - \frac{xy}{8} + 9$	4	$\frac{3}{x}$, $\frac{y}{7}$, $-\frac{xy}{8}$ and 9 & so on.

Types of Algebraic Expressions :

- (i) **Monomial** : An algebraic expression having only one term is called a monomial. For ex. $8y$, $-7xy$, $4x^2$, abx , etc. 'mono' means 'one'.
- (ii) **Binomial** : An algebraic expression having two terms is called a binomial. For ex. $8x + 3y$, $8x + 3$, $8 + 3y$, $a + bz$, $9 - 4y$, $2x^2 - 4z$, $6y^2 - 5y$, etc. 'bi' means 'two'.
- (iii) **Trinomial** : An algebraic expression having three terms is called a trinomial. For ex. $ax - 5y + 8z$, $3x^2 + 4x + 7$, $9y^2 - 3y + 2x$, etc. 'tri' means 'three'.
- (iv) **Multinomial** : An algebraic expression having two or more terms is called a multinomial.

➤ FACTORS AND COEFFICIENTS

◆ Factor :

Each combination of the constants and variables, which form a term, is called a factor.

For examples :

- (i) 7, x and 7x are factors of 7x, in which 7 is constant (numerical) factor and x is variable (literal) factor.
- (ii) In $-5x^2y$, the numerical factor is -5 and literal factors are : x, y, xy , x^2 and x^2y .

◆ Coefficient :

Any factor of a term is called the coefficient of the remaining term.

For example :

- (i) In $7x$; 7 is coefficient of x
- (ii) In $-5x^2y$; 5 is coefficient of $-x^2y$; -5 is coefficient of x^2y .

Ex. 1 Write the coefficient of :

- (i) x^2 in $3x^3 - 5x^2 + 7$
- (ii) xy in $8xyz$
- (iii) $-y$ in $2y^2 - 6y + 2$
- (iv) x^0 in $3x + 7$

Sol.

- (i) -5
- (ii) $8z$
- (iii) 6
- (iv) Since $x^0 = 1$, Therefore
 $3x + 7 = 3x + 7x^0$
coefficient of x^0 is 7.

➤ DEGREE OF A POLYNOMIAL

The greatest power (exponent) of the terms of a polynomial is called degree of the polynomial.

For example :

- (a) In polynomial $5x^2 - 8x^7 + 3x$:
 - (i) The power of term $5x^2 = 2$
 - (ii) The power of term $-8x^7 = 7$
 - (iii) The power of $3x = 1$

Since, the greatest power is 7, therefore degree of the polynomial $5x^2 - 8x^7 + 3x$ is 7

(b) The degree of polynomial :

- (i) $4y^3 - 3y + 8$ is 3
- (ii) $7p + 2$ is 1 ($p = p^1$)
- (iii) $2m - 7m^8 + m^{13}$ is 13 and so on.

❖ EXAMPLES ❖

Ex.2 Find which of the following algebraic expression is a polynomial.

- (i) $3x^2 - 5x$
- (ii) $x + \frac{1}{x}$
- (iii) $\sqrt{y} - 8$
- (iv) $z^5 - \sqrt[3]{z} + 8$

Sol.

- (i) $3x^2 - 5x = 3x^2 - 5x^1$

It is a polynomial.

- (ii) $x + \frac{1}{x} = x^1 + x^{-1}$

It is not a polynomial.

- (iii) $\sqrt{y} - 8 = y^{1/2} - 8$

Since, the power of the first term ($\sqrt{y}$) is $\frac{1}{2}$, which is not a whole number.

- (iv) $z^5 - \sqrt[3]{z} + 8 = z^5 - z^{1/3} + 8$

Since, the exponent of the second term is $1/3$, which is not a whole number. Therefore, the given expression is not a polynomial.

Ex.3 Find the degree of the polynomial :

- (i) $5x - 6x^3 + 8x^7 + 6x^2$
- (ii) $2y^{12} + 3y^{10} - y^{15} + y + 3$
- (iii) x
- (iv) 8

Sol.

- (i) Since the term with highest exponent (power) is $8x^7$ and its power is 7.

$\therefore$ The degree of given polynomial is 7.

- (ii) The highest power of the variable is 15

$\Rightarrow$ degree = 15.

- (iii) $x = x^1 \Rightarrow$ degree is 1.

- (iv) $8 = 8x^0 \Rightarrow$ degree = 0

➤ TYPES OF POLYNOMIALS

(A) Based on degree :

If degree of polynomial is

			Examples
1.	One	Linear	$x + 3, y - x + 2, \sqrt{3}x - 3$
2.	Two	Quadratic	$2x^2 - 7, \frac{1}{3}x^2 + y^2 - 2xy, x^2 + 1 + 3y$
3.	Three	Cubic	$x^3 + 3x^2 - 7x + 8, 2x^2 + 5x^3 + 7,$
4.	Four	bi-quadratic	$x^4 + y^4 + 2x^2y^2, x^4 + 3, \dots$

(B) Based on Terms :

If number of terms in polynomial is

			Examples
1.	One	Monomial	$7x, 5x^9, \frac{7}{3}x^{16}, xy, \dots$
2.	Two	Binomial	$2 + 7y^6, y^3 + x^{14}, 7 + 5x^9, \dots$
3.	Three	Trinomial	$x^3 - 2x + y, x^{31} + y^{32} + z^{33}, \dots$

Note : (1) Degree of constant polynomials

(Ex.5, 7, -3, 8/5, ...) is zero.

(2) Degree of zero polynomial (zero = 0 = zero polynomial) is not defined.

➤ POLYNOMIAL IN ONE VARIABLE

If a polynomial has only one variable then it is called polynomial in one variable.

Ex. $P(x) = 2x^3 + 5x - 3$ Cubic trinomial

$Q(x) = 7x^7 - 5x^5 - 3x^3 + x + 3$ polynomial of degree 7

$R(y) = y$ Linear, monomial

$S(t) = t^2 + 3$ Quadratic Binomial

Note : General form of a polynomial in one variable x of degree ' n ' is $a_nx^n + a_{n-1}x^{n-1} + a_{n-2}x^{n-2} + \dots + a_2x^2 + a_1x + a_0$, $a_n \neq 0$, where $a_n, a_{n-1}, \dots, a_2, a_1, a_0$ all are constants.

$\therefore$ for linear $ax + b$, $a \neq 0$

for quadratic $ax^2 + bx + c$, $a \neq 0$

for cubic $ax^3 + bx^2 + cx + d$, $a \neq 0$

➤ REMAINDER THEOREM

- Remainder obtained on dividing polynomial $p(x)$ by $x - a$ is equal to $p(a)$.
- If a polynomial $p(x)$ is divided by $(x + a)$ the remainder is the value of $p(x)$ at $x = -a$.
- $(x - a)$ is a factor of polynomial $p(x)$ if $p(a) = 0$
- $(x + a)$ is a factor of polynomial $p(x)$ if $p(-a) = 0$
- $(x - a)(x - b)$ is a factor of polynomial $p(x)$, if $p(a) = 0$ and $p(b) = 0$.

❖ EXAMPLES ❖

Ex.4 Find the remainder when $4x^3 - 3x^2 + 2x - 4$ is divided by

(a) $x - 1$ (b) $x + 2$ (c) $x + \frac{1}{2}$

Sol. Let $p(x) = 4x^3 - 3x^2 + 2x - 4$

- (a) When $p(x)$ is divided by $(x - 1)$, then by remainder theorem, the required remainder will be $p(1)$

$$\begin{aligned} p(1) &= 4(1)^3 - 3(1)^2 + 2(1) - 4 \\ &= 4 \times 1 - 3 \times 1 + 2 \times 1 - 4 \\ &= 4 - 3 + 2 - 4 = -1 \end{aligned}$$

- (b) When $p(x)$ is divided by $(x + 2)$, then by remainder theorem, the required remainder will be $p(-2)$.

$$\begin{aligned} p(-2) &= 4(-2)^3 - 3(-2)^2 + 2(-2) - 4 \\ &= 4 \times (-8) - 3 \times 4 - 4 - 4 \\ &= -32 - 12 - 8 = -52 \end{aligned}$$

- (c) When $p(x)$ is divided by, $\left(x + \frac{1}{2}\right)$ then by remainder theorem, the required remainder will be

$$\begin{aligned} p\left(-\frac{1}{2}\right) &= 4\left(-\frac{1}{2}\right)^3 - 3\left(-\frac{1}{2}\right)^2 + 2\left(-\frac{1}{2}\right) - 4 \\ &= 4 \times \left(-\frac{1}{8}\right) - 3 \times \frac{1}{4} - 2 \times \frac{1}{2} - 4 \\ &= -\frac{1}{2} - \frac{3}{4} - 1 - 4 = -\frac{1}{2} - \frac{3}{4} - 5 \\ &= \frac{-2-3-20}{4} = \frac{-25}{4} \end{aligned}$$

VALUES OF A POLYNOMIAL

For a polynomial $f(x) = 3x^2 - 4x + 2$.

To find its value at $x = 3$;

replace x by 3 everywhere.

So, the value of $f(x) = 3x^2 - 4x + 2$ at $x = 3$ is

$$f(3) = 3 \times 3^2 - 4 \times 3 + 2$$

$$= 27 - 12 + 2 = 17.$$

Similarly, the value of polynomial

$$f(x) = 3x^2 - 4x + 2,$$

$$(i) \text{ at } x = -2 \text{ is } f(-2) = 3(-2)^2 - 4(-2) + 2 \\ = 12 + 8 + 2 = 22$$

$$(ii) \text{ at } x = 0 \text{ is } f(0) = 3(0)^2 - 4(0) + 2 \\ = 0 - 0 + 2 = 2$$

$$(iii) \text{ at } x = \frac{1}{2} \text{ is } f\left(\frac{1}{2}\right) = 3\left(\frac{1}{2}\right)^2 - 4\left(\frac{1}{2}\right) + 2 \\ = \frac{3}{4} - 2 + 2 = \frac{3}{4}$$

Ex.5 Find the value of the polynomial $5x - 4x^2 + 3$ at:

$$(i) x = 0 \quad (ii) x = -1$$

Sol. Let $p(x) = 5x - 4x^2 + 3$.

$$(i) \text{ At } x = 0, p(0) = 5 \times 0 - 4 \times (0)^2 + 3 \\ = 0 - 0 + 3 = 3$$

$$(ii) \text{ At } x = -1, p(-1) = 5(-1) - 4(-1)^2 + 3 \\ = -5 - 4 + 3 = -6$$

ZEROES OF A POLYNOMIAL

If for $x = a$, the value of the polynomial $p(x)$ is 0 i.e., $p(a) = 0$; then $x = a$ is a zero of the polynomial $p(x)$.

For example :

$$(i) \text{ For polynomial } p(x) = x - 2; p(2) = 2 - 2 = 0 \\ \therefore x = 2 \text{ or simply } 2 \text{ is a zero of the polynomial } \\ p(x) = x - 2.$$

$$(ii) \text{ For the polynomial } g(u) = u^2 - 5u + 6; \\ g(3) = (3)^2 - 5 \times 3 + 6 = 9 - 15 + 6 = 0 \\ \therefore 3 \text{ is a zero of the polynomial } g(u) \\ = u^2 - 5u + 6.$$

$$\text{Also, } g(2) = (2)^2 - 5 \times 2 + 6 = 4 - 10 + 6 = 0$$

$\therefore 2$ is also a zero of the polynomial

$$g(u) = u^2 - 5u + 6$$

- (a) Every linear polynomial has one and only one zero.
- (b) A given polynomial may have more than one zeroes.
- (c) If the degree of a polynomial is n ; the largest number of zeroes it can have is also n .

For example :

If the degree of a polynomial is 5, the polynomial can have at the most 5 zeroes; if the degree of a polynomial is 8; largest number of zeroes it can have is 8.

- (d) A zero of a polynomial need not be 0.

For example : If $f(x) = x^2 - 4$,
then $f(2) = (2)^2 - 4 = 4 - 4 = 0$

Here, zero of the polynomial $f(x) = x^2 - 4$ is 2 which itself is not 0.

- (e) 0 may be a zero of a polynomial.

For example : If $f(x) = x^2 - x$,
then $f(0) = 0^2 - 0 = 0$

Here 0 is the zero of polynomial

$$f(x) = x^2 - x.$$

❖ EXAMPLES ❖

Ex.6 Verify whether the indicated numbers are zeroes of the polynomial corresponding to them in the following cases :

$$(i) p(x) = 3x + 1, x = -\frac{1}{3}$$

$$(ii) p(x) = (x + 1)(x - 2), x = -1, 2$$

$$(iii) p(x) = x^2, x = 0$$

$$(iv) p(x) = \lambda x + m, x = -\frac{m}{\lambda}$$

$$(v) p(x) = 2x + 1, x = \frac{1}{2}$$

Sol. (i) $p(x) = 3x + 1$

$$\Rightarrow p\left(-\frac{1}{3}\right) = 3 \times -\frac{1}{3} + 1 = -1 + 1 = 0$$

$$\therefore x = -\frac{1}{3} \text{ is a zero of } p(x) = 3x + 1.$$

$$(ii) \quad p(x) = (x + 1)(x - 2)$$

$$\Rightarrow p(-1) = (-1 + 1)(-1 - 2) = 0 \times -3 = 0$$

and, $p(2) = (2 + 1)(2 - 2) = 3 \times 0 = 0$

$\therefore x = -1$ and $x = 2$ are zeroes of the given polynomial.

$$(iii) \quad p(x) = x^2 \Rightarrow p(0) = 0^2 = 0$$

$\therefore x = 0$ is a zero of the given polynomial

$$(iv) \quad p(x) = \lambda x + m \Rightarrow p\left(-\frac{m}{\lambda}\right) = \lambda\left(-\frac{m}{\lambda}\right) + m$$

$$= -m + m = 0$$

$\therefore x = -\frac{m}{\lambda}$ is a zero of the given polynomial.

$$(v) \quad p(x) = 2x + 1 \Rightarrow p\left(\frac{1}{2}\right) = 2 \times \frac{1}{2} + 1$$

$$= 1 + 1 = 2 \neq 0$$

$\therefore x = \frac{1}{2}$ is not a zero of the given polynomial.

Ex.7 Find the zero of the polynomial in each of the following cases :

$$(i) \quad p(x) = x + 5 \quad (ii) \quad p(x) = 2x + 5$$

$$(iii) \quad p(x) = 3x - 2$$

Sol. To find the zero of a polynomial $p(x)$ means to solve the polynomial equation $p(x) = 0$.

(i) For the zero of polynomial $p(x) = x + 5$

$$p(x) = 0 \Rightarrow x + 5 = 0 \Rightarrow x = -5$$

$\therefore x = -5$ is a zero of the polynomial $p(x) = x + 5$.

(ii) $p(x) = 0 \Rightarrow 2x + 5 = 0$

$$\Rightarrow 2x = -5 \text{ and } x = \frac{-5}{2}$$

$\therefore x = \frac{-5}{2}$ is a zero of $p(x) = 2x + 5$.

(iii) $p(x) = 0 \Rightarrow 3x - 2 = 0$

$$\Rightarrow 3x = 2 \text{ and } x = \frac{2}{3}.$$


$$\therefore x = \frac{2}{3} \text{ is zero of } p(x) = 3x - 2$$

➤ GEOMETRIC MEANING OF THE ZEROES OF A POLYNOMIAL

Let us consider linear polynomial $ax + b$. The graph of $y = ax + b$ is a straight line.


For example : The graph of $y = 3x + 4$ is a straight line passing through $(0, 4)$ and $(2, 10)$.

x	0	2
y = 3x + 4	4	10
Points	A	B



(i) Let us consider the graph of $y = 2x - 4$ intersects the x-axis at $x = 2$. The zero $2x - 4$ is 2. Thus, the zero of the polynomial $2x - 4$ is the x-coordinate of the point where the graph $y = 2x - 4$ intersects the x-axis.

x	2	0
y = 2x - 4	0	-4
Points	A	B




(ii) A general equation of a linear polynomial is $ax + b$. The graph of $y = ax + b$ is a straight line which intersects the x-axis at $\left(\frac{-b}{a}, 0\right)$.

Zero of the polynomial $ax + b$ is the x-coordinate of the point of intersection of the graph with x-axis.

- (iii) Let us consider the quadratic polynomial $x^2 - 4x + 3$. The graph of $x^2 - 4x + 3$ intersects the x-axis at the point (1, 0) and (3, 0). Zeroes of the polynomial $x^2 - 4x + 3$ are the x-coordinates of the points of intersection of the graph with x-axis.


x	1	2	3	4	5
$y = x^2 - 4x + 3$	0	-1	0	3	8
Points	A	B	C	D	E

The shape of the graph of the quadratic polynomials is $\cup$ and the curve is known as parabola.



- (iv) Now let us consider one more polynomial $-x^2 + 2x + 8$. Graph of this polynomial intersects the x-axis at the points (4, 0), (-2, 0). Zeroes of the polynomial $-x^2 + 2x + 8$ are the x-coordinates of the points at which the graph intersects the x-axis. The shape of the graph of the given quadratic polynomial is $\cap$ and the curve is known as parabola.

x	-2	-1	0	1	2	3	4
y	0	5	8	9	8	7	0
Points	A	B	C	D	E	F	G



The zeroes of a quadratic polynomial $ax^2 + bx + c$ are the x-coordinates of the points where the graph of $y = ax^2 + bx + c$ intersects the x-axis.

Cubic polynomial : Let us find out geometrically how many zeroes a cubic has.


Let consider cubic polynomial

$$x^3 - 6x^2 + 11x - 6.$$

x	0	0.5	1	1.5	2	2.5	3	3.5	4
$y = x^3 - 6x^2 + 11x - 6$	-6	-1.875	0	0.375	0	-0.375	0	1.875	6
Points	A	B	C	D	E	F	G	H	I


Case 1 :

The graph of the cubic equation intersects the x-axis at three points (1, 0), (2, 0) and (3, 0). Zeroes of the given polynomial are the x-coordinates of the points of intersection with the x-axis.



Case 2 :

The cubic equation $x^3 - x^2$ intersects the x-axis at the point (0, 0) and (1, 0). Zero of a polynomial $x^3 - x^2$ are the x-coordinates of the point where the graph cuts the x-axis.




Zeroes of the cubic polynomial are 0 and 1.

Case 3 :

$$y = x^3$$

Cubic polynomial has only one zero.













In brief : A cubic equation can have 1 or 2 or 3 zeroes or any polynomial of degree three can have at most three zeroes.

Remarks : In general, polynomial of degree n , the graph of $y = p(x)$ passes x-axis at most at n points. Therefore, a polynomial $p(x)$ of degree n has at most n zeroes.

❖ EXAMPLES ❖

Ex.8 Which of the following correspond to the graph to a linear or a quadratic polynomial and find the number of zeroes of polynomial.



Sol. (i) The graph is a straight line so the graph is of a linear polynomial. The number of zeroes is one as the graph intersects the x-axis at one point only.

(ii) The graph is a parabola. So, this is the graph of quadratic polynomial. The number of zeroes is zero as the graph does not intersect the x-axis.

(iii) Here the polynomial is quadratic as the graph is a parabola. The number of zeroes is one as the graph intersects the x-axis at one point only (two coincident points).

(iv) Here, the polynomial is quadratic as the graph is a parabola. The number of zeroes is two as the graph intersects the x-axis at two points.

(v) The polynomial is linear as the graph is straight line. The number of zeroes is one as the graph intersects the x-axis at one point only.

(vi) The polynomial is quadratic as the graph is a parabola. The number of zeroes is one as the graph intersects the x-axis at one point (two coincident points) only.

(vii) The polynomial is quadratic as the graph is a parabola. The number of zeroes is zero, as the graph does not intersect the x-axis.

(viii) Polynomial is neither linear nor quadratic as the graph is neither a straight line nor a parabola is one as the graph intersects the x-axis at one point only.

(ix) Here, the polynomial is quadratic as the graph is a parabola. The number of zeroes is one as the graph intersects the x-axis at one point only (two coincident points).

(x) The polynomial is linear as the graph is a straight line. The number of zeroes is one as the graph intersects the x-axis at only one point.



RELATIONSHIP BETWEEN THE ZEROES AND THE COEFFICIENTS OF A POLYNOMIAL.

Consider quadratic polynomial

$$P(x) = 2x^2 - 16x + 30.$$

$$\begin{aligned} \text{Now, } 2x^2 - 16x + 30 &= (2x - 6)(x - 3) \\ &= 2(x - 3)(x - 5) \end{aligned}$$

The zeroes of $P(x)$ are 3 and 5.

Sum of the zeroes

$$= 3 + 5 = 8 = \frac{-(-16)}{2} = -\left[\frac{\text{coefficient of } x}{\text{coefficient of } x^2}\right]$$

Product of the zeroes

$$= 3 \times 5 = 15 = \frac{30}{2} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

So if $ax^2 + bx + c$, $a \neq 0$ is a quadratic polynomial and α, β are two zeroes of polynomial

$$\text{then } \boxed{\alpha + \beta = -\frac{b}{a}}, \boxed{\alpha\beta = \frac{c}{a}}$$

❖ EXAMPLES ❖

Ex.9 Find the zeroes of the quadratic polynomial $6x^2 - 13x + 6$ and verify the relation between the zeroes and its coefficients.

$$\begin{aligned} \text{Sol. We have, } 6x^2 - 13x + 6 &= 6x^2 - 4x - 9x + 6 \\ &= 2x(3x - 2) - 3(3x - 2) \\ &= (3x - 2)(2x - 3) \end{aligned}$$

So, the value of $6x^2 - 13x + 6$ is 0, when $(3x - 2) = 0$ or $(2x - 3) = 0$ i.e.,

$$\text{When } x = \frac{2}{3} \text{ or } \frac{3}{2}$$

Therefore, the zeroes of $6x^2 - 13x + 6$ are

$$\frac{2}{3} \text{ and } \frac{3}{2}.$$

Sum of the zeroes

$$= \frac{2}{3} + \frac{3}{2} = \frac{13}{6} = \frac{-(-13)}{6} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

Product of the zeroes

$$= \frac{2}{3} \times \frac{3}{2} = \frac{6}{6} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Ex.10 Find the zeroes of the quadratic polynomial $4x^2 - 9$ and verify the relation between the zeroes and its coefficients.

Sol. We have,

$$4x^2 - 9 = (2x)^2 - 3^2 = (2x - 3)(2x + 3)$$

So, the value of $4x^2 - 9$ is 0, when

$$2x - 3 = 0 \text{ or } 2x + 3 = 0$$

$$\text{i.e., when } x = \frac{3}{2} \text{ or } x = -\frac{3}{2}.$$

Therefore, the zeroes of $4x^2 - 9$ are $\frac{3}{2}$ & $-\frac{3}{2}$.

Sum of the zeroes

$$= \frac{3}{2} - \frac{3}{2} = 0 = \frac{-(0)}{4} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

Product of the zeroes

$$= \left(\frac{3}{2}\right) \left(-\frac{3}{2}\right) = \frac{-9}{4} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Ex.11 Find the zeroes of the quadratic polynomial $9x^2 - 5$ and verify the relation between the zeroes and its coefficients.

Sol. We have,

$$9x^2 - 5 = (3x)^2 - (\sqrt{5})^2 = (3x - \sqrt{5})(3x + \sqrt{5})$$

So, the value of $9x^2 - 5$ is 0,

$$\text{when } 3x - \sqrt{5} = 0 \text{ or } 3x + \sqrt{5} = 0$$

$$\text{i.e., when } x = \frac{\sqrt{5}}{3} \text{ or } x = -\frac{\sqrt{5}}{3}.$$

Sum of the zeroes

$$= \frac{\sqrt{5}}{3} - \frac{\sqrt{5}}{3} = 0 = \frac{-(0)}{9} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

Product of the zeroes

$$= \left(\frac{\sqrt{5}}{3}\right) \left(-\frac{\sqrt{5}}{3}\right) = \frac{-5}{9} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Ex.12 If α and β are the zeroes of $ax^2 + bx + c$, $a \neq 0$ then verify the relation between the zeroes and its coefficients.

Sol. Since α and β are the zeroes of polynomial $ax^2 + bx + c$.

Therefore, $(x - \alpha)$, $(x - \beta)$ are the factors of the polynomial $ax^2 + bx + c$.

$$\Rightarrow ax^2 + bx + c = k(x - \alpha)(x - \beta)$$

$$\Rightarrow ax^2 + bx + c = k\{x^2 - (\alpha + \beta)x + \alpha\beta\}$$

$$\Rightarrow ax^2 + bx + c = kx^2 - k(\alpha + \beta)x + k\alpha\beta \dots(1)$$

Comparing the coefficients of x^2 , x and constant terms of (1) on both sides, we get

$$a = k, b = -k(\alpha + \beta) \text{ and } c = k\alpha\beta$$

$$\Rightarrow \alpha + \beta = -\frac{b}{k} \text{ and } \alpha\beta = \frac{c}{k}$$

$$\alpha + \beta = \frac{-b}{a} \quad \text{and} \quad \alpha\beta = \frac{c}{a} \quad [\Theta k = a]$$

$$\text{Sum of the zeroes} = \frac{-b}{a} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

$$\text{Product of the zeroes} = \frac{c}{a} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Ex. 13 Prove relation between the zeroes and the coefficient of the quadratic polynomial $ax^2 + bx + c$.

Sol. Let α and β be the zeroes of the polynomial $ax^2 + bx + c$

$$\therefore \alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \dots(1)$$

$$\beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \dots(2)$$

By adding (1) and (2), we get

$$\alpha + \beta = \frac{-b + \sqrt{b^2 - 4ac}}{2a} + \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-2b}{2a} = -\frac{b}{a} = \frac{\text{coefficient of } x}{\text{coefficient of } x^2}$$

Hence, sum of the zeroes of the polynomial $ax^2 + bx + c$ is $-\frac{b}{a}$

By multiplying (1) and (2), we get

$$\begin{aligned} \alpha\beta &= \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{(-b)^2 - \sqrt{(b^2 - 4ac)^2}}{4a^2} = \frac{b^2 - b^2 + 4ac}{4a^2} \\ &= \frac{4ac}{4a^2} = \frac{c}{a} \\ &= \frac{\text{constant term}}{\text{coefficient of } x^2} \end{aligned}$$

$$\text{Hence, product of zeroes} = \frac{c}{a}$$

In general, it can be proved that if α , β , γ are the zeroes of a cubic polynomial $ax^3 + bx^2 + cx + d$, then

$$\alpha + \beta + \gamma = \frac{-b}{a}$$

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$$

$$\alpha\beta\gamma = \frac{-d}{a}$$

Note, $\frac{b}{a}$, $\frac{c}{a}$ and $\frac{d}{a}$ are meaningful because $a \neq 0$.

Ex.14 find the zeroes of the quadratic polynomial $x^2 - 2x - 8$ and verify a relationship between zeroes and its coefficients.

$$\begin{aligned} \text{Sol.} \quad x^2 - 2x - 8 &= x^2 - 4x + 2x - 8 \\ &= x(x - 4) + 2(x - 4) = (x - 4)(x + 2) \end{aligned}$$

So, the value of $x^2 - 2x - 8$ is zero when $x - 4 = 0$ or $x + 2 = 0$ i.e., when $x = 4$ or $x = -2$.

So, the zeroes of $x^2 - 2x - 8$ are 4, -2.

Sum of the zeroes

$$= 4 - 2 = 2 = \frac{-(-2)}{1} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

Product of the zeroes

$$= 4(-2) = -8 = \frac{-8}{1} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Ex.15 Verify that the numbers given along side of the cubic polynomials are their zeroes. Also

verify the relationship between the zeroes and the coefficients. $2x^3 + x^2 - 5x + 2$; $\frac{1}{2}, 1, -2$

Sol. Here, the polynomial $p(x)$ is

$$2x^3 + x^2 - 5x + 2$$

Value of the polynomial $2x^3 + x^2 - 5x + 2$

when $x = 1/2$

$$= 2\left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^2 - 5\left(\frac{1}{2}\right) + 2 = \frac{1}{4} + \frac{1}{4} - \frac{5}{2} + 2 = 0$$

So, $1/2$ is a zero of $p(x)$.

On putting $x = 1$ in the cubic polynomial

$$2x^3 + x^2 - 5x + 2$$

$$= 2(1)^3 + (1)^2 - 5(1) + 2 = 2 + 1 - 5 + 2 = 0$$

On putting $x = -2$ in the cubic polynomial

$$2x^3 + x^2 - 5x + 2$$

$$= 2(-2)^3 + (-2)^2 - 5(-2) + 2$$

$$= -16 + 4 + 10 + 2 = 0$$

Hence, $\frac{1}{2}, 1, -2$ are the zeroes of the given polynomial.

Sum of the zeroes of $p(x)$

$$= \frac{1}{2} + 1 - 2 = -\frac{1}{2} = \frac{-\text{coefficient of } x^2}{\text{coefficient of } x^3}$$

Sum of the products of two zeroes taken at a time

$$= \frac{1}{2} \times 1 + \frac{1}{2} \times (-2) + 1 \times (-2)$$

$$= \frac{1}{2} - 1 - 2 = -\frac{5}{2} = \frac{\text{coefficient of } x}{\text{coefficient of } x^3}$$

Product of all the three zeroes

$$= \left(\frac{1}{2}\right) \times (1) \times (-2) = -1$$

$$= \frac{-(2)}{2} = \frac{-\text{constant term}}{\text{coefficient of } x^3}$$

➤ SYMMETRIC FUNCTIONS OF ZEROS OF A QUADRATIC POLYNOMIAL.

◆ Symmetric function :

An algebraic expression in α and β , which remains unchanged, when α and β are interchanged is known as symmetric function in α and β .

For example, $\alpha^2 + \beta^2$ and $\alpha^3 + \beta^3$ etc. are symmetric functions. Symmetric function is to be expressed in terms of $(\alpha + \beta)$ and $\alpha\beta$. So, this can be evaluated for a given quadratic equation.

◆ Some useful relations involving α and β :

- $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
- $(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$
- $\alpha^2 - \beta^2 = (\alpha + \beta)(\alpha - \beta) = (\alpha + \beta)\sqrt{(\alpha + \beta)^2 - 4\alpha\beta}$
- $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$
- $\alpha^3 - \beta^3 = (\alpha - \beta)^3 + 3\alpha\beta(\alpha - \beta)$
- $\alpha^4 + \beta^4 = [(\alpha + \beta)^2 - 2\alpha\beta]^2 - 2(\alpha\beta)^2$
- $\alpha^4 - \beta^4 = (\alpha^2 + \beta^2)(\alpha^2 - \beta^2)$ then use (1) and (3)

◆ EXAMPLES ◆

Ex.16 If α and β are the zeroes of the polynomial $ax^2 + bx + c$. Find the value of

- (i) $\alpha - \beta$ (ii) $\alpha^2 + \beta^2$.

Sol. Since α and β are the zeroes of the polynomial $ax^2 + bx + c$.

$$\therefore \alpha + \beta = -\frac{b}{a}; \quad \alpha\beta = \frac{c}{a}$$

$$(i) (\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

$$= \left(-\frac{b}{a}\right)^2 - \frac{4c}{a} = \frac{b^2}{a^2} - \frac{4c}{a} = \frac{b^2 - 4ac}{a^2}$$

$$\alpha - \beta = \frac{\sqrt{b^2 - 4ac}}{a}$$

$$(ii) \alpha^2 + \beta^2 = \alpha^2 + \beta^2 + 2\alpha\beta - 2\alpha\beta$$

$$= (\alpha + \beta)^2 - 2\alpha\beta$$

$$= \left(-\frac{b}{a}\right)^2 - 2\left(\frac{c}{a}\right) = \frac{b^2 - 2ac}{a^2}$$

Ex.17 If α and β are the zeroes of the quadratic polynomial $ax^2 + bx + c$. Find the value of

- (i) $\alpha^2 - \beta^2$ (ii) $\alpha^3 + \beta^3$.

Sol. Since α and β are the zeroes of $ax^2 + bx + c$

$$\therefore \alpha + \beta = \frac{-b}{a}, \quad \alpha\beta = \frac{c}{a}$$

$$(i) \alpha^2 - \beta^2 = (\alpha + \beta)(\alpha - \beta)$$

$$\begin{aligned} &= -\frac{b}{a} \sqrt{(\alpha + \beta)^2 - 4\alpha\beta} \\ &= -\frac{b}{a} \sqrt{\left(\frac{-b}{a}\right)^2 - 4\frac{c}{a}} = -\frac{b}{a} \sqrt{\frac{b^2 - 4ac}{a^2}} \\ &= -\frac{b\sqrt{b^2 - 4ac}}{a^2} \end{aligned}$$

$$(ii) \alpha^3 + \beta^3 = (\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta)$$

$$\begin{aligned} &= (\alpha + \beta)[(\alpha^2 + \beta^2 + 2\alpha\beta) - 3\alpha\beta] \\ &= (\alpha + \beta)[(\alpha + \beta)^2 - 3\alpha\beta] \\ &= \frac{-b}{a} \left[\left(\frac{-b}{a}\right)^2 - \frac{3c}{a} \right] \\ &= \frac{-b}{a} \left[\frac{b^2}{a^2} - \frac{3c}{a} \right] = \frac{-b}{a} \left(\frac{b^2 - 3ac}{a^2} \right) \\ &= \frac{-b^3 + 3abc}{a^3} \end{aligned}$$

➤ TO FORM A QUADRATIC POLYNOMIAL WITH THE GIVEN ZEROES

Let zeroes of a quadratic polynomial be α and β .

$$\therefore x = \alpha, \quad x = \beta$$

$$x - \alpha = 0, \quad x - \beta = 0$$

The obviously the quadratic polynomial is

$$(x - \alpha)(x - \beta)$$

$$\text{i.e., } x^2 - (\alpha + \beta)x + \alpha\beta$$

$$x^2 - (\text{Sum of the zeroes})x + \text{Product of the zeroes}$$

❖ EXAMPLES ❖

Ex.18 Form the quadratic polynomial whose zeroes are 4 and 6.

Sol. Sum of the zeroes = 4 + 6 = 10

$$\text{Product of the zeroes} = 4 \times 6 = 24$$

Hence the polynomial formed

$$= x^2 - (\text{sum of zeroes})x + \text{Product of zeroes}$$

$$= x^2 - 10x + 24$$

Ex.19 Form the quadratic polynomial whose zeroes are -3, 5.

Sol. Here, zeroes are -3 and 5.

$$\text{Sum of the zeroes} = -3 + 5 = 2$$

$$\text{Product of the zeroes} = (-3) \times 5 = -15$$

Hence the polynomial formed

$$= x^2 - (\text{sum of zeroes})x + \text{Product of zeroes}$$

$$= x^2 - 2x - 15$$

Ex.20 Find a quadratic polynomial whose sum of zeroes and product of zeroes are respectively-

$$(i) \frac{1}{4}, -1 \quad (ii) \sqrt{2}, \frac{1}{3} \quad (iii) 0, \sqrt{5}$$

Sol. Let the polynomial be $ax^2 + bx + c$ and its zeroes be α and β .

$$(i) \text{ Here, } \alpha + \beta = \frac{1}{4} \text{ and } \alpha \cdot \beta = -1$$

Thus the polynomial formed

$$= x^2 - (\text{Sum of zeroes})x + \text{Product of zeroes}$$

$$= x^2 - \left(\frac{1}{4}\right)x - 1 = x^2 - \frac{x}{4} - 1$$

$$\text{The other polynomial are } k \left(x^2 - \frac{x}{4} - 1 \right)$$

$$\text{If } k = 4, \text{ then the polynomial is } 4x^2 - x - 4.$$

$$(ii) \text{ Here, } \alpha + \beta = \sqrt{2}, \alpha\beta = \frac{1}{3}$$

Thus the polynomial formed

$$= x^2 - (\text{Sum of zeroes})x + \text{Product of zeroes}$$

$$= x^2 - (\sqrt{2})x + \frac{1}{3} \text{ or } x^2 - \sqrt{2}x + \frac{1}{3}$$

$$\text{Other polynomial are } k \left(x^2 - \sqrt{2}x + \frac{1}{3} \right)$$

$$\text{If } k = 3, \text{ then the polynomial is}$$

$$3x^2 - 3\sqrt{2}x + 1$$

$$(iii) \text{ Here, } \alpha + \beta = 0 \text{ and } \alpha\beta = \sqrt{5}$$

Thus the polynomial formed

$$= x^2 - (\text{Sum of zeroes})x + \text{Product of zeroes}$$

$$= x^2 - (0)x + \sqrt{5} = x^2 + \sqrt{5}$$

Ex.21 Find a cubic polynomial with the sum of its zeroes, sum of the products of its zeroes taken

two at a time, and product of its zeroes as 2, -7 and -14, respectively.

Sol. Let the cubic polynomial be

$$ax^3 + bx^2 + cx + d$$

$$\Rightarrow x^3 + \frac{b}{a}x^2 + \frac{c}{a}x + \frac{d}{a} \quad \dots(1)$$

and its zeroes are α , β and γ , then

$$\alpha + \beta + \gamma = 2 = -\frac{b}{a}$$

$$\alpha\beta + \beta\gamma + \alpha\gamma = -7 = \frac{c}{a}$$

$$\alpha\beta\gamma = -14 = -\frac{d}{a}$$

Putting the values of $\frac{b}{a}$, $\frac{c}{a}$ and $\frac{d}{a}$ in (1),

we get

$$x^3 + (-2)x^2 + (-7)x + 14$$

$$\Rightarrow x^3 - 2x^2 - 7x + 14$$

Ex.22 Find the cubic polynomial with the sum, sum of the product of its zeroes taken two at a time and product of its zeroes as 0, -7 and -6 respectively.

Sol. Let the cubic polynomial be

$$ax^3 + bx^2 + cx + d$$

$$\Rightarrow x^3 + \frac{b}{a}x^2 + \frac{c}{a}x + \frac{d}{a} \quad \dots(1)$$

and its zeroes are α , β , γ . Then

$$\alpha + \beta + \gamma = 0 = -\frac{b}{a}$$

$$\alpha\beta + \beta\gamma + \alpha\gamma = -7 = \frac{c}{a}$$

$$\alpha\beta\gamma = -6 = -\frac{d}{a}$$

Putting the values of $\frac{b}{a}$, $\frac{c}{a}$ and $\frac{d}{a}$ in (1),

we get

$$x^3 - (0)x^2 + (-7)x + (-6)$$

$$\text{or } x^3 - 7x + 6$$

Ex.23 If α and β are the zeroes of the polynomials $ax^2 + bx + c$ then form the polynomial whose zeroes are $\frac{1}{\alpha}$ and $\frac{1}{\beta}$.

Sol. Since α and β are the zeroes of $ax^2 + bx + c$

$$\text{So } \alpha + \beta = \frac{-b}{a}, \alpha\beta = \frac{c}{a}$$

$$\text{Sum of the zeroes} = \frac{1}{\alpha} + \frac{1}{\beta} = \frac{\beta + \alpha}{\alpha\beta}$$

$$= \frac{-\frac{b}{a}}{\frac{c}{a}} = -\frac{b}{c}$$

Product of the zeroes

$$= \frac{1}{\alpha} \cdot \frac{1}{\beta} = \frac{1}{\alpha\beta} = \frac{a}{c}$$

But required polynomial is

$$x^2 - (\text{sum of zeroes})x + \text{Product of zeroes}$$

$$\Rightarrow x^2 - \left(\frac{-b}{c}\right)x + \left(\frac{a}{c}\right)$$

$$\text{or } x^2 + \frac{b}{c}x + \frac{a}{c}$$

$$\text{or } c\left(x^2 + \frac{b}{c}x + \frac{a}{c}\right)$$

$$\Rightarrow cx^2 + bx + a$$

Ex.24 If α and β are the zeroes of the polynomial $x^2 + 4x + 3$, form the polynomial whose zeroes are $1 + \frac{\beta}{\alpha}$ and $1 + \frac{\alpha}{\beta}$.

Sol. Since α and β are the zeroes of the polynomial $x^2 + 4x + 3$.

$$\text{Then, } \alpha + \beta = -4, \alpha\beta = 3$$

Sum of the zeroes

$$= 1 + \frac{\beta}{\alpha} + 1 + \frac{\alpha}{\beta} = \frac{\alpha\beta + \beta^2 + \alpha\beta + \alpha^2}{\alpha\beta}$$

$$= \frac{\alpha^2 + \beta^2 + 2\alpha\beta}{\alpha\beta} = \frac{(\alpha + \beta)^2}{\alpha\beta} = \frac{(-4)^2}{3} = \frac{16}{3}$$

Product of the zeroes

$$\begin{aligned}
 &= \left(1 + \frac{\beta}{\alpha}\right) \left(1 + \frac{\alpha}{\beta}\right) = 1 + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{\alpha\beta}{\alpha\beta} \\
 &= 2 + \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{2\alpha\beta + \alpha^2 + \beta^2}{\alpha\beta} \\
 &= \frac{(\alpha + \beta)^2}{\alpha\beta} = \frac{(-4)^2}{3} = \frac{16}{3}
 \end{aligned}$$

But required polynomial is

$x^2 - (\text{sum of zeroes})x + \text{product of zeroes}$

$$\text{or } x^2 - \frac{16}{3}x + \frac{16}{3} \quad \text{or } k \left(x^2 - \frac{16}{3}x + \frac{16}{3} \right)$$

$$\text{or } 3 \left(x^2 - \frac{16}{3}x + \frac{16}{3} \right) \quad (\text{if } k = 3)$$

$$\Rightarrow 3x^2 - 16x + 16$$

➤ WORKING RULE TO DIVIDE A POLYNOMIAL BY ANOTHER POLYNOMIAL

Step 1:

First arrange the term of dividend and the divisor in the decreasing order of their degrees.

Step 2 :

To obtain the first term of quotient divide the highest degree term of the dividend by the highest degree term of the divisor.

Step 3 :

To obtain the second term of the quotient, divide the highest degree term of the new dividend obtained as remainder by the highest degree term of the divisor.

Step 4 :

Continue this process till the degree of remainder is less than the degree of divisor.

◆ Division Algorithm for Polynomial

If $p(x)$ and $g(x)$ are any two polynomials with

$g(x) \neq 0$, then we can find polynomials $q(x)$ and $r(x)$ such that

$$p(x) = q(x) \times g(x) + r(x)$$

where $r(x) = 0$ or degree of $r(x) < \text{degree of } g(x)$.

The result is called Division Algorithm for polynomials.

$$\text{Dividend} = \text{Quotient} \times \text{Divisor} + \text{Remainder}$$

❖ EXAMPLES ❖

Ex.25 Divide $3x^3 + 16x^2 + 21x + 20$ by $x + 4$.

Sol.

$$\begin{array}{r}
 3x^2 + 4x + 5 \\
 x+4 \overline{) 3x^3 + 16x^2 + 21x + 20} \\
 \underline{3x^3 + 12x^2} \\
 4x^2 + 21x + 20 \\
 \underline{4x^2 + 16x} \\
 5x + 20 \\
 \underline{5x + 20} \\
 0
 \end{array}$$

First term of $q(x) = \frac{3x^3}{x} = 3x^2$
 Second term of $q(x) = \frac{4x^2}{x} = 4x$
 Third term of $q(x) = \frac{5x}{x} = 5$

$$\text{Quotient} = 3x^2 + 4x + 5$$

$$\text{Remainder} = 0$$

Ex.26 Apply the division algorithm to find the quotient and remainder on dividing $p(x)$ by $g(x)$ as given below :

$$p(x) = x^3 - 3x^2 + 5x - 3, g(x) = x^2 - 2$$

Sol. We have,

$$p(x) = x^3 - 3x^2 + 5x - 3 \text{ and } g(x) = x^2 - 2$$

$$\begin{array}{r}
 x - 3 \\
 x^2-2 \overline{) x^3 - 3x^2 + 5x - 3} \\
 \underline{x^3 - 2x^2} \\
 -x^2 + 5x - 3 \\
 \underline{-x^2 + 2x} \\
 -3x - 3 \\
 \underline{-3x - 6} \\
 3
 \end{array}$$

First term of quotient is $\frac{x^3}{x^2} = x$
 Second term of quotient is $\frac{-3x^2}{x^2} = -3$

We stop here since

$$\text{degree of } (7x - 9) < \text{degree of } (x^2 - 2)$$

$$\text{So, quotient} = x - 3, \text{ remainder} = 7x - 9$$

Therefore,

$$\text{Quotient} \times \text{Divisor} + \text{Remainder}$$

$$= (x - 3)(x^2 - 2) + 7x - 9$$

$$= x^3 - 2x - 3x^2 + 6 + 7x - 9$$

$$= x^3 - 3x^2 + 5x - 3 = \text{Dividend}$$

Therefore, the division algorithm is verified.

Ex.27 Apply the division algorithm to find the quotient and remainder on dividing $p(x)$ by $g(x)$ as given below

$$p(x) = x^4 - 3x^2 + 4x + 5, g(x) = x^2 + 1 - x$$

Sol. We have,

$$p(x) = x^4 - 3x^2 + 4x + 5, g(x) = x^2 + 1 - x$$

$$\begin{array}{r} x^2 + x - 3 \\ x^2 - x + 1 \overline{) x^4 - 3x^2 + 4x + 5} \\ \underline{- + } \\ x^3 - 4x^2 + 4x + 5 \\ \underline{- + } \\ x^3 - x^2 + x \\ \underline{- + } \\ -3x^2 + 3x + 5 \\ \underline{- + } \\ -3x^2 + 3x - 3 \\ \underline{+ - } \\ 8 \end{array}$$

We stop here since

degree of (8) < degree of $(x^2 - x + 1)$.

So, quotient = $x^2 + x - 3$, remainder = 8

Therefore,

Quotient $\times$ Divisor + Remainder

$$= (x^2 + x - 3)(x^2 - x + 1) + 8$$

$$= x^4 - x^3 + x^2 + x^3 - x^2 + x - 3x^2 + 3x - 3 + 8$$

$$= x^4 - 3x^2 + 4x + 5 = \text{Dividend}$$

Therefore the Division Algorithm is verified.

Ex.28 Check whether the first polynomial is a factor of the second polynomial by applying the division algorithm. $t^2 - 3$; $2t^4 + 3t^3 - 2t^2 - 9t - 12$.

Sol. We divide $2t^4 + 3t^3 - 2t^2 - 9t - 12$ by $t^2 - 3$

$$\begin{array}{r} 2t^2 + 3t + 4 \\ t^2 - 3 \overline{) 2t^4 + 3t^3 - 2t^2 - 9t - 12} \\ \underline{- + } \\ 3t^3 + 4t^2 + 9t - 12 \\ \underline{- + } \\ 4t^2 - 12 \\ \underline{- + } \\ 4t^2 - 12 \\ \underline{- + } \\ 0 \end{array}$$

Here, remainder is 0, so $t^2 - 3$ is a factor of $2t^4 + 3t^3 - 2t^2 - 9t - 12$.

$$2t^4 + 3t^3 - 2t^2 - 9t - 12 = (2t^2 + 3t + 4)(t^2 - 3)$$

Ex.29 Obtain all the zeroes of

$$3x^4 + 6x^3 - 2x^2 - 10x - 5, \text{ if two of its zeroes are } \sqrt{\frac{5}{3}} \text{ and } -\sqrt{\frac{5}{3}}.$$

Sol. Since two zeroes are $\sqrt{\frac{5}{3}}$ and $-\sqrt{\frac{5}{3}}$,

$$x = \sqrt{\frac{5}{3}}, x = -\sqrt{\frac{5}{3}}$$

$$\Rightarrow \left(x - \sqrt{\frac{5}{3}}\right)\left(x + \sqrt{\frac{5}{3}}\right) = x^2 - \frac{5}{3} \text{ or } 3x^2 - 5$$

is a factor of the given polynomial.

Now, we apply the division algorithm to the given polynomial and $3x^2 - 5$.

$$\begin{array}{r} x^2 + 2x + 1 \\ 3x^2 - 5 \overline{) 3x^4 + 6x^3 - 2x^2 - 10x - 5} \\ \underline{- + } \\ 6x^3 + 3x^2 - 10x - 5 \\ \underline{- + } \\ 3x^2 - 5 \\ \underline{- + } \\ 0 \end{array}$$

$$\text{So, } 3x^4 + 6x^3 - 2x^2 - 10x - 5$$

$$= (3x^2 - 5)(x^2 + 2x + 1) + 0$$

$$\text{Quotient} = x^2 + 2x + 1 = (x + 1)^2$$

Zeroes of $(x + 1)^2$ are $-1, -1$.

Hence, all its zeroes are $\sqrt{\frac{5}{3}}, -\sqrt{\frac{5}{3}}, -1, -1$.

Ex.30 On dividing $x^3 - 3x^2 + x + 2$ by a polynomial $g(x)$, the quotient and remainder were $x - 2$ and $-2x + 4$, respectively. Find $g(x)$.

Sol. $p(x) = x^3 - 3x^2 + x + 2$

$$q(x) = x - 2 \text{ and } r(x) = -2x + 4$$

By Division Algorithm, we know that

$$p(x) = q(x) \times g(x) + r(x)$$

Therefore,

$$x^3 - 3x^2 + x + 2 = (x - 2) \times g(x) + (-2x + 4)$$

$$\Rightarrow x^3 - 3x^2 + x + 2 + 2x - 4 = (x - 2) \times g(x)$$

$$\Rightarrow g(x) = \frac{x^3 - 3x^2 + 3x - 2}{x - 2}$$

On dividing $x^3 - 3x^2 + 3x - 2$ by $x - 2$, we get $g(x)$

$$\begin{array}{r} x-2 \overline{) \begin{array}{r} x^3 - 3x^2 + 3x - 2 \\ x^3 - 2x^2 \\ \hline -x^2 + 3x - 2 \\ -x^2 + 2x \\ \hline +x - 2 \\ x - 2 \\ \hline 0 \end{array}} \end{array}$$

First term of quotient is $\frac{x^3}{x} = x$

Second term of quotient is $\frac{-x^2}{x} = -x$

Third term of quotient is $\frac{x}{x} = 1$

$$\text{Hence, } g(x) = x^2 - x + 1.$$

Ex.31 Give examples of polynomials $p(x)$, $q(x)$ and $r(x)$, which satisfy the division algorithm and

(i) $\deg p(x) = \deg q(x)$

(ii) $\deg q(x) = \deg r(x)$

(iii) $\deg q(x) = 0$

Sol. (i) Let $q(x) = 3x^2 + 2x + 6$, degree of $q(x) = 2$

$p(x) = 12x^2 + 8x + 24$, degree of $p(x) = 2$

Here, $\deg p(x) = \deg q(x)$

(ii) $p(x) = x^5 + 2x^4 + 3x^3 + 5x^2 + 2$

$q(x) = x^2 + x + 1$, degree of $q(x) = 2$

$g(x) = x^3 + x^2 + x + 1$

$r(x) = 2x^2 - 2x + 1$, degree of $r(x) = 2$

Here, $\deg q(x) = \deg r(x)$

(iii) Let $p(x) = 2x^4 + 8x^3 + 6x^2 + 4x + 12$

$q(x) = 2$, degree of $q(x) = 0$

$g(x) = x^4 + 4x^3 + 3x^2 + 2x + 6$

$r(x) = 0$

Here, $\deg q(x) = 0$

Ex.32 If the zeroes of polynomial $x^3 - 3x^2 + x + 1$ are $a - b$, a , $a + b$. Find a and b .

Sol. Θ $a - b$, a , $a + b$ are zeros

$\therefore$ product $(a - b)a(a + b) = -1$

$\Rightarrow (a^2 - b^2)a = -1 \quad \dots(1)$

and sum of zeroes is $(a - b) + a + (a + b) = 3$

$\Rightarrow 3a = 3 \Rightarrow a = 1 \quad \dots(2)$

by (1) and (2)

$(1 - b^2)1 = -1$

$\Rightarrow 2 = b^2 \Rightarrow b = \pm\sqrt{2}$

$\therefore a = -1$ & $b = \pm\sqrt{2}$ **Ans.**

Ex.33 If two zeroes of the polynomial

$x^4 - 6x^3 - 26x^2 + 138x - 35$ are $2 \pm \sqrt{3}$, find other zeroes.

Sol. Θ $2 \pm \sqrt{3}$ are zeroes.

$\therefore x = 2 \pm \sqrt{3}$

$\Rightarrow x - 2 = \pm\sqrt{3}$ (squaring both sides)

$\Rightarrow (x - 2)^2 = 3 \Rightarrow x^2 + 4 - 4x - 3 = 0$

$\Rightarrow x^2 - 4x + 1 = 0$, is a factor of given polynomial

$\therefore$ other factors

$$= \frac{x^4 - 6x^3 - 26x^2 + 138x - 35}{x^2 - 4x + 1}$$

$$\begin{array}{r} x^2 - 2x - 35 \\ x^2 - 4x + 1 \overline{) \begin{array}{r} x^4 - 6x^3 - 26x^2 + 138x - 35 \\ x^4 - 4x^3 + x^2 \\ \hline -2x^3 - 27x^2 + 138x - 35 \\ -2x^3 + 8x^2 + 2x \\ \hline -35x^2 + 140x - 35 \\ -35x^2 + 140x - 35 \\ \hline 0 \end{array}} \end{array}$$

$\therefore$ other factors $= x^2 - 2x - 35$

$= x^2 - 7x + 5x - 35 = x(x - 7) + 5(x - 7)$

$= (x - 7)(x + 5)$

$\therefore$ other zeroes are $(x - 7) = 0 \Rightarrow x = 7$

$x + 5 = 0 \Rightarrow x = -5$ **Ans.**

Ex.34 If the polynomial $x^4 - 6x^3 + 16x^2 - 25x + 10$ is divided by another polynomial $x^2 - 2x + k$, the remainder comes out to be $x + a$, find k & a .

Sol.

$$\begin{array}{r}
 x^2 - 2x + k \overline{) \begin{array}{l} x^2 - 4x + (8 - k) \\ x^4 - 6x^3 + 16x^2 - 25x + 10 \\ \underline{x^4 - 2x^3 + x^2k} \\ -4x^3 + x^2(16 - k) - 25x + 10 \\ \underline{-4x^3 + x^2(8) - 4xk} \\ x^2[8 - k] + x[4k - 25] + 10 \\ \underline{x^2[8 - k] - 2x[8 - k] + k(8 - k)} \\ x[4k - 25 + 16 - 2k] + 10 - 8k + k^2 \end{array}}
 \end{array}$$

According to questions, remainder is $x + a$

$\therefore$ coefficient of $x = 1$

$$\Rightarrow 2k - 9 = 1$$

$$\Rightarrow k = (10/2) = 5$$

Also constant term = a

$$\Rightarrow k^2 - 8k + 10 = a \Rightarrow (5)^2 - 8(5) + 10 = a$$

$$\Rightarrow a = 25 - 40 + 10$$

$$\Rightarrow a = -5$$

$$\therefore k = 5, a = -5$$

Ans.

EXERCISE # 1

A. Very Short Answer Type Questions

Factorize each of the following expression

Q.1 $x^2 - x - 42$

Q.2 $6 - 5y - y^2$

Q.3 $a^2 + 46a + 205$

Q.4 $ab + ac - b^2 - bc$

Q.5 $p^4 - 81q^4$

Use remainder theorem to find remainder, when $p(x)$ is divided by $q(x)$ in following questions.

Q.6 $p(x) = 2x^2 - 5x + 7$, $q(x) = x - 1$

Q.7 $p(x) = x^9 - 5x^4 + 1$, $q(x) = x + 1$

Q.8 $p(x) = 2x^3 - 3x^2 + 4x - 1$, $q(x) = x + 2$

B. Short Answer Type Questions

Q.9 Find positive square root of $36x^2 + 60x + 25$

Q.10 Simplify : $\sqrt{2a^2 + 2\sqrt{6ab} + 3b^2}$

Q.11 $(x^2 + 4y)^2 + 21(x^2 + 4y) + 98$

Q.12 Find the value of k if $(x - 2)$ is a factor of $2x^3 - 6x^2 + 5x + k$.

Q.13 Find the value of k if $(x + 3)$ is a factor of $3x^2 + kx + 6$.

Q.14 $p(x) = 3x^6 - 7x^5 + 7x^4 - 3x^3 + 2x^2 - 2$, $q(x) = x - 1$

Q.15 For what value of k is $y^3 + ky + 2k - 2$ exactly divisible by $(y + 1)$?

C. Long Answer Type Questions

Q.16 If $x + 1$ and $x - 1$ are factors of $mx^3 + x^2 - 2x + n$, find the value of m and n .

Q.17 Find the zeros of the polynomial $f(x) = 2x^2 + 5x - 12$ and verify the relation between its zeroes and coefficients.

Q.18 Find the zeroes of the polynomial $f(x) = x^2 - 2$ and verify the relation between its zeroes and coefficients.

Q.19 Obtain the zeroes of the quadratic polynomial $\sqrt{3}x^2 - 8x + 4\sqrt{3}$ and verify the relation between its zeroes and coefficients.

Q.20 Find a cubic polynomial with the sum of its zeroes, sum of the products of its zeroes taken two at a time and the product of its zeroes as 2, -7 and -14 respectively.

Q.21 Find a cubic polynomial whose zeroes are 3, 5 and -2 .

Q.22 Divide $5x^3 - 13x^2 + 21x - 14$ by $(3 - 2x + x^2)$ and verify the division algorithm.

Q.23 What real number should be subtracted from the polynomial $(3x^3 + 10x^2 - 14x + 9)$ so that $(3x - 2)$ divides it exactly?

Q.24 Find all the zeroes of $(2x^4 - 3x^3 - 5x^2 + 9x - 3)$, it being given that two of its zeroes are $\sqrt{3}$ and $-\sqrt{3}$.

ANSWER KEY

A. VERTY SHORT ANSWER TYPE :

1. $(x + 6)(x - 7)$ 2. $(6 + y)(1 - y)$ 3. $(a + 41)(a + 5)$ 4. $(a - b)(b + c)$
5. $(p + 3q)(p - 3q)(p^2 + 9q^2)$ 6. 4 7. -5 8. -37

B. SHORT ANSWER TYPE :

9. $6x + 5$ 10. $(\sqrt{2}a + \sqrt{3}b)$ 11. $(x^2 + 4y + 7)(x^2 + 4y + 14)$ 12. -2
13. 11 15. 3

C. LONG ANSWER TYPE :

16. $m = 2, n = -1$ 17. $-4, \frac{3}{2}$ 18. $-\sqrt{2}, \sqrt{2}$ 19. $2\sqrt{3}, \frac{2}{\sqrt{3}}$
20. $x^3 - 2x^2 - 7x + 14$ 21. $x^3 - 6x^2 - x + 30$ 22. quotient = $5x - 3$, Remainder = -5
23. 5 24. $\sqrt{3}, -\sqrt{3}, 1, \frac{1}{2}$

EXERCISE # 2

- Q.1** If $\left(x + \frac{1}{x}\right) = 3$, then find value of $\left(x^2 + \frac{1}{x^2}\right)$.
- Q.2** If $\left(x - \frac{1}{x}\right) = \frac{1}{2}$, then find $\left(4x^2 + \frac{4}{x^2}\right)$.
- Q.3** If $\left(x + \frac{1}{x}\right) = 4$, then find $\left(x^4 + \frac{1}{x^4}\right)$.
- Q.4** If $(x - 2)$ is a factor of $(x^2 + 3qx - 2q)$, then find the value of q .
- Q.5** If $x^3 + 6x^2 + 4x + k$ is exactly divisible by $(x + 2)$, then find the value of k .
- Q.6** Let $f(x) = x^3 - 6x^2 + 11x - 6$. Then, which one of the following is not factor of $f(x)$?
(A) $x - 1$ (B) $x - 2$
(C) $x + 3$ (D) $x - 3$
- Q.7** If $x^{100} + 2x^{99} + k$ is divisible by $(x + 1)$, then find the value of k .
- Q.8** On dividing $(x^3 - 6x + 7)$ by $(x + 1)$, find the remainder.
- Q.9** Find the value of expression $(16x^2 + 24x + 9)$ for $x = -\frac{3}{4}$.
- Q.10** If $2x^3 + 5x^2 - 4x - 6$ is divided by $2x + 1$, then find remainder.
- Q.11** If $p(x) = x^2 - 2x - 3$, then find
(i) $p(3)$; (ii) $p(-1)$
- Q.12** Find the zeros of the quadratic polynomial $(6x^2 - 7x - 3)$ and verify the relation between its zeros and coefficients.
- Q.13** Find the zeros of the quadratic polynomial $(5u^2 + 10u)$ and verify the relation between the zeros and the coefficients.
- Q.14** Find the quadratic polynomial whose zeros are $\frac{2}{3}$ and $\frac{-1}{4}$. Verify the relation between the coefficients and the zeros of the polynomial.
- Q.15** Find the quadratic polynomial, sum of whose zeros is 8 and their product is 12. Hence, find the zeros of the polynomial.
- Q.16** Find the quadratic polynomial, the sum of whose zeros is -5 and their product is 6. Hence, find the zeros of the polynomial.
- Q.17** Find the quadratic polynomial, the sum of whose zeros is 0 and their product is -1 . Hence, find the zeros of the polynomial.
- Q.18** Find a quadratic polynomial whose one zero is $5 + \sqrt{7}$.
- Q.19** On dividing $(x^3 - 3x^2 + x + 2)$ by a polynomial $g(x)$, the quotient and remainder are $(x - 2)$ and $(-2x + 4)$ respectively. Find $g(x)$.
- Q.20** If the polynomial $(x^4 + 2x^3 + 8x^2 + 12x + 18)$ is divided by another polynomial $(x^2 + 5)$, the remainder comes out to be $(px + q)$. Find the value of p and q .
- Q.21** Obtain all zeros of the polynomial $(2x^3 - 4x - x^2 + 2)$, if two of its zeros are $\sqrt{2}$ and $-\sqrt{2}$.
- Q.22** If 1 and -2 are two zeros of the polynomial $(x^3 - 4x^2 - 7x + 10)$, find its third zero.
- Q.23** Find all the zeros of the polynomial $(2x^4 - 11x^3 + 7x^2 + 13x - 7)$, it being given that two of its zeros are $(3 + \sqrt{2})$ and $(3 - \sqrt{2})$.
- Q.24** If α, β are the zeros of the polynomial $f(x) = x^2 - 5x + k$ such that $\alpha - \beta = 1$, find the value of k .
- Q.25** Show that the polynomial $f(x) = x^4 + 4x^2 + 6$ has no zero.
- Q.26** Use remainder theorem to find the value of k , it being given that when $x^3 + 2x^2 + kx + 3$ is divided by $(x - 3)$, then the remainder is 21.

ANSWER KEY

- | | | | | |
|--|---------------------------------|---|----------------------|--------|
| 1. 7 | 2. 9 | 4. -1 | 5. -8 | 6. (C) |
| 7. 1 | 8. 12 | 9. 0 | 10. -3 | |
| 11. (i) 0, (ii) 0 | 12. $\frac{3}{2}, -\frac{1}{3}$ | 13. $-2, 0$ | 14. $12x^2 - 5x - 2$ | |
| 15. $(x^2 - 8x + 12), \{6, 2\}$ | | 16. $(x^2 + 5x + 6), \{-3, -2\}$ | | |
| 17. $(x^2 - 1), \{1, -1\}$ | 18. $x^2 - 10x + 18$ | 19. $x^2 - x + 1$ | 20. $p = 2, q = 3$ | |
| 21. $\sqrt{2}, -\sqrt{2}, \frac{1}{2}$ | 22. 5 | 23. $(3 + \sqrt{2}), (3 - \sqrt{2}), \frac{1}{2}, -1$ | | |
| 24. $k = 6$ | 26. $k = -9$ | | | |