

Sample Question Paper - 11
Mathematics (041)
Class- XII, Session: 2021-22
TERM II

Time Allowed: 2 hours

Maximum Marks: 40

General Instructions:

1. This question paper contains three sections – A, B and C. Each part is compulsory.
2. Section - A has 6 short answer type (SA1) questions of 2 marks each.
3. Section – B has 4 short answer type (SA2) questions of 3 marks each.
4. Section - C has 4 long answer-type questions (LA) of 4 marks each.
5. There is an internal choice in some of the questions.
6. Q 14 is a case-based problem having 2 sub-parts of 2 marks each.

Section A

1. Evaluate: $\int \frac{x+1}{x(x+\log x)} dx$ [2]

OR

Evaluate the integral: $\int \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$

2. Find the general solution of the differential equation: $(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{(1+x^2)}$ [2]

3. For any vector \vec{a} , prove that $|\vec{a} \times \hat{i}|^2 + |\vec{a} \times \hat{j}|^2 + |\vec{a} \times \hat{k}|^2 = 2|\vec{a}|^2$ [2]

4. Find the direction cosines of the line $\frac{x-2}{2} = \frac{2y-5}{-3}$, $z = -1$. Also, find the vector equation of the line. [2]

5. The probability that a person will get an electric contract is $\frac{2}{5}$ and the probability that he will not get plumbing contract is $\frac{4}{7}$. If the probability of getting at least one contract is $\frac{2}{3}$, what is the probability that he will get both? [2]

6. There are three urns A, B, and C. Urn A contains 4 red balls and 3 black balls. Urn B contains 5 red balls and 4 black balls. Urn C contains 4 red and 4 black balls. One ball is drawn from each of these urns. What is the probability that 3 balls drawn consist of 2 red balls and a black ball? [2]

Section B

7. Evaluate $\int e^{-3x} \cos^3 x dx$ [3]

8. Solve the differential equation: $(x^2 + 3xy + y^2) dx - x^2 dy = 0$ [3]

OR

Verify that $y = Ae^{ax} \cos bx + Be^{ax} \sin bx$, where A and B are arbitrary constants, is the general solution of the differential equation $\frac{d^2y}{dx^2} - 2a \frac{dy}{dx} + (a^2 + b^2)y = 0$.

9. If $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j}$ and $\vec{c} = 3\hat{i} - 4\hat{j} - 5\hat{k}$, then find a unit vector perpendicular to both of the vectors $(\vec{a} - \vec{b})$ and $(\vec{c} - \vec{b})$. [3]

10. Find the intercepts made on the coordinate axes by the plane $2x + y - 2z = 3$ and find also the direction cosines of the normal to the plane. [3]

OR

Show that the line whose vector equation is $\vec{r} = 2\hat{i} + 5\hat{j} + 7\hat{k} + \lambda(\hat{i} + 3\hat{j} + 4\hat{k})$ is parallel to the plane whose vector equation is $\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 7$. Also, find the distance between them.

Section C

11. Evaluate: $\int \frac{x^2}{(x^2+6x-3)} dx$ [4]
12. Using integration, find the area of the region enclosed between the two circles $x^2 + y^2 = 1$ and $(x - 1)^2 + y^2 = 1$. [4]

OR

Using integration find the area of the region bounded by the curve $y = \sqrt{4 - x^2}$, $x^2 + y^2 - 4x = 0$ and the x-axis.

13. Show that the lines $\frac{x-a+d}{\alpha-\delta} = \frac{y-a}{\alpha} = \frac{z-a-d}{\alpha+\delta}$ and $\frac{x-b+c}{\beta-\gamma} = \frac{y-b}{\beta} = \frac{z-b-c}{\beta+\gamma}$ are coplanar. Also find the equation of the plane containing them. [4]

CASE-BASED/DATA-BASED

14. Kamal is a good card player. He plays many magic tricks on cards as well. He has a deck of 52 cards. He shuffled the cards well and drew five cards one by one, with replacement. [4]



Find the probability that

- i. all five cards are diamonds.
- ii. none is a diamond.

Solution

MATHEMATICS 041

Class 12 - Mathematics

Section A

1. Let $I = \int \frac{x+1}{x(x+\log x)} dx \dots(i)$

Now let $(x + \log x) = t$ then, we have

$$d(x + \log x) = dt$$

$$\Rightarrow \left(1 + \frac{1}{x}\right) dx = dt$$

$$\Rightarrow \left(\frac{x+1}{x}\right) dx = dt$$

$$\Rightarrow dx = \frac{x}{x+1} dt$$

Putting $(x + \log x) = t$ and $dx = \frac{x}{x+1} dt$ in equation (i), we get

$$I = \int \frac{x+1}{x \times t} \times \frac{x}{x+1} dt$$

$$= \int \frac{dt}{t}$$

$$= \log |t| + c$$

$$= \log |x + \log x| + c$$

$$\Rightarrow I = \log |x + \log x| + c$$

OR

$$\text{Let, } I = \int \left(\frac{\sin x + \cos x}{\sqrt{\sin 2x}} \right) dx$$

Put, $\sin x - \cos x = t$

$$\Rightarrow (\cos x + \sin x) dx = dt$$

$$\text{Also } (\sin x - \cos x)^2 = t^2$$

$$\Rightarrow \sin^2 x + \cos^2 x - 2 \sin x \cos x = t^2$$

$$\Rightarrow 1 - t^2 = \sin(2x)$$

$$\therefore I = \int \frac{dt}{\sqrt{1-t^2}}$$

$$= \sin^{-1} t + C \quad \dots \left[\int \frac{dt}{\sqrt{a^2-t^2}} = \sin^{-1} \frac{t}{a} + C \right]$$

$$= \sin^{-1}(\sin x - \cos x) + C \quad (\because t = \sin x - \cos x)$$

2. Given $(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{1+x^2}$

$$\Rightarrow \frac{dy}{dx} + \frac{2x}{1+x^2} \cdot y = \frac{1}{(1+x^2)^2}$$

This is of the form $\frac{dy}{dx} + Py = Q$

where $P = \frac{2x}{1+x^2}$ and $Q = \frac{1}{(1+x^2)^2}$

This the given differential equation is linear

$$\text{Now, } IF = e^{\int P dx} \Rightarrow I \cdot F = e^{\int \frac{2x dx}{1+x^2}}$$

$$= e^{\log(1+x^2)} = 1 + x^2$$

Therefore the solution is given by $y \cdot (1.F) = \int (1.F)Q + C$

$$\Rightarrow y(1+x^2) = \int (1+x^2) \frac{1}{(1+x^2)^2} dx + C$$

$$\Rightarrow y(1+x^2) = \int \frac{dx}{1+x^2} + C$$

$$\Rightarrow y \cdot (1+x^2) = \tan^{-1} x + C.$$

3. Let $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k} \dots(i)$

Then,

$$\vec{a} \times \hat{i} = (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \times \hat{i} = a_1(\hat{i} \times \hat{i}) + a_2(\hat{j} \times \hat{i}) + a_3(\hat{k} \times \hat{i}) = -a_2 \hat{k} + a_3 \hat{j}$$

$$\Rightarrow |\vec{a} \times \hat{i}|^2 = a_2^2 + a_3^2$$

$$\vec{a} \times \hat{j} = (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \times \hat{j} = a_1(\hat{i} \times \hat{j}) + a_2(\hat{j} \times \hat{j}) + a_3(\hat{k} \times \hat{j}) = a_1 \hat{k} - a_3 \hat{i}$$

$$\Rightarrow |\vec{a} \times \hat{j}|^2 = a_1^2 + a_3^2$$

$$\text{and } \vec{a} \times \hat{k} = (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \times \hat{k} = a_1(\hat{i} \times \hat{k}) + a_2(\hat{j} \times \hat{k}) + a_3(\hat{k} \times \hat{k}) = -a_1 \hat{j} + a_2 \hat{i}$$

$$\Rightarrow |\vec{a} \times \hat{k}|^2 = a_1^2 + a_2^2$$

$$\therefore |\vec{a} \times \hat{i}|^2 + |\vec{a} \times \hat{j}|^2 + |\vec{a} \times \hat{k}|^2 = a_2^2 + a_3^2 + a_1^2 + a_3^2 + a_1^2 + a_2^2$$

$$\Rightarrow |\vec{a} \times \hat{i}|^2 + |\vec{a} \times \hat{j}|^2 + |\vec{a} \times \hat{k}|^2 = 2(a_1^2 + a_2^2 + a_3^2) = 2|\vec{a}|^2 \dots \text{using (i)}$$

4. The Cartesian equations of the given line are

$$\frac{x-2}{2} = \frac{2y-5}{-3}, z = -1$$

These above equations can be re-written as

$$\frac{x-2}{2} = \frac{2y-5}{-3} = \frac{z+1}{0} \text{ or, } \frac{x-2}{2} = \frac{y-5/2}{-3/2} = \frac{z+1}{0}$$

This shows that the given line passes through the point $(2, \frac{5}{2}, -1)$ and has direction ratios proportional to $2, -\frac{3}{2}, 0$. So, its direction cosines are

$\frac{2}{5}, 0$. So, its direction cosines are

$$\frac{2}{\sqrt{2^2 + (-\frac{3}{2})^2 + 0^2}}, \frac{-3/2}{\sqrt{2^2 + (-\frac{3}{2})^2 + 0^2}}, \frac{0}{\sqrt{2^2 + (-\frac{3}{2})^2 + 0^2}} \text{ or, } \frac{2}{5/2}, \frac{-3/2}{5/2}, 0$$

$$\text{or, } \frac{4}{5}, -\frac{3}{5}, 0$$

The given line passes through the point having a position vector $\vec{a} = 2\hat{i} + \frac{5}{2}\hat{j} - \hat{k}$ and is parallel to the vector $\vec{b} = 2\hat{i} - \frac{3}{2}\hat{j} + 0\hat{k}$.

Therefore, it's vector equation is

$$\vec{r} = \left(2\hat{i} + \frac{5}{2}\hat{j} - \hat{k}\right) + \lambda \left(2\hat{i} - \frac{3}{2}\hat{j} + 0\hat{k}\right)$$

5. Consider the following events:

A = Person gets an electric contract, B = Person gets plumbing contract .

Therefore, we have,

$$P(A) = \frac{2}{5}, P(\bar{B}) = \frac{4}{7} \text{ and } P(A \cup B) = \frac{2}{3}$$

By addition theorem of probability, we have,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow \frac{2}{3} = \frac{2}{5} + \left(1 - \frac{4}{7}\right) - P(A \cap B)$$

$$\Rightarrow P(A \cap B) = \frac{2}{5} + \frac{3}{7} - \frac{2}{3} = \frac{17}{105}$$

6. We are given that,

Urn A (4R + 3B)

Urn B (5R + 4B)

Urn C (4R + 4B)

Required probability is given by,

$P(\text{two red and one black}) = P(\text{Red from urn A}) \times P(\text{Black from urn B}) \times P(\text{Red from urn C}) + P(\text{Red from urn A})$

$\times P(\text{Red from urn B}) \times P(\text{Black from urn C}) + P(\text{Black from urn A}) \times P(\text{Red from urn B}) \times P(\text{Red from urn C})$

$$= \frac{3}{7} \times \frac{5}{9} \times \frac{4}{8} + \frac{4}{7} \times \frac{4}{9} \times \frac{4}{8} + \frac{4}{7} \times \frac{5}{9} \times \frac{4}{8}$$

$$= \frac{5}{42} \times \frac{16}{126} \times \frac{20}{126}$$

$$= \frac{15+16+20}{126} = \frac{51}{126} = \frac{17}{42}$$

Section B

7. Given integral is: $\int e^{-3x} \cos^3 x dx$

Using trigonometric identity $\cos 3x = 4 \cos^3 x - 3 \cos x$

$$\Rightarrow \int e^{-3x} \cos^3 x dx = \frac{1}{4} \int e^{-3x} (\cos 3x + 3 \cos x) dx$$

$$\Rightarrow \frac{1}{4} \int e^{-3x} (\cos 3x + 3 \cos x) dx = \frac{1}{4} \int e^{-3x} \cos 3x dx + \frac{3}{4} \int e^{-3x} \cos x dx \dots (i)$$

Using a general formula i.e.

$$\Rightarrow \int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx)$$

$$\Rightarrow \int e^{-3x} \cos 3x dx = \frac{e^{-3x}}{(-3)^2+3^2} ((-3) \cos 3x + 3 \sin 3x)$$

$$\Rightarrow \frac{e^{-3x}}{(-3)^2+3^2} ((-3) \cos 3x + 3 \sin 3x) = \frac{e^{-3x}}{6} (\sin 3x - \cos 3x) \dots \text{(ii)}$$

$$\Rightarrow \int e^{-3x} \cos x dx = \frac{e^{-3x}}{(-3)^2+1^2} ((-3) \cos 3x + 3 \sin 3x) = \frac{e^{-3x}}{10} (\sin x - 3 \cos x)$$

$$= \frac{e^{-3x}}{10} (\sin x - 3 \cos x) \dots \text{(iii)}$$

On putting (ii) and (iii) in (i)

$$\Rightarrow \frac{1}{4} \int e^{-3x} \cos 3x dx + \frac{3}{4} \int e^{-3x} \cos x dx = \frac{e^{-3x}}{4 \times 6} (\sin 3x - \cos 3x) + \frac{3e^{-3x}}{4 \times 10} (\sin x - \cos x)$$

$$\Rightarrow \int e^{-3x} \cos^3 x dx = e^{-3x} \left\{ \frac{(\sin 3x - \cos 3x)}{24} + \frac{3(\sin x - 3 \cos x)}{40} \right\} + C$$

8. The given differential equation is,

$$(x^2 + 3xy + y^2) dx - x^2 dy = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{x^2 + 3xy + y^2}{x^2}$$

This is a homogeneous differential equation

Putting $y = vx$ and $\frac{dy}{dx} = v + x \frac{dv}{dx}$, we get

$$v + x \frac{dv}{dx} = \frac{x^2 + 3vx^2 + v^2x^2}{x^2}$$

$$\Rightarrow x \frac{dv}{dx} = 1 + 3v + v^2 - v$$

$$\Rightarrow x \frac{dv}{dx} = 1 + v^2 + 2v$$

$$\Rightarrow \frac{1}{1+v^2+2v} dv = \frac{1}{x} dx$$

Integrating both sides, we get

$$\int \frac{1}{1+v^2+2v} dv = \int \frac{1}{x} dx$$

$$\Rightarrow \int \frac{1}{(1+v)^2} dv = \int \frac{1}{x} dx$$

$$\Rightarrow -\frac{1}{(1+v)} = \log |x| + C$$

$$\Rightarrow \log |x| + \frac{1}{(1+v)} = -C$$

Putting $v = \frac{y}{x}$, we get

$$\therefore \log |x| + \frac{x}{(x+y)} = C_1$$

where, $C_1 = -C$

Hence, $\log |x| + \frac{x}{(x+y)} = C_1$ is the required solution.

OR

We have, $y = Ae^{ax} \cos bx + Be^{ax} \sin bx \dots \text{(i)}$

Differentiating (i) on both sides w.r.t. x, we have,

$$\frac{dy}{dx} = A \cdot [e^{ax} \{-b \sin bx\} + ae^{ax} \cos bx] + B \cdot [e^{ax} (b \cos bx) + ae^{ax} \sin bx]$$

$$\Rightarrow \frac{dy}{dx} = a (Ae^{ax} \cos bx + Be^{ax} \sin bx) + b \{-Ae^{ax} \sin bx + Be^{ax} \cos bx\}$$

$$\Rightarrow \frac{dy}{dx} = ay + be^{ax} \{B \cos bx - A \sin bx\} \dots \text{(ii) [using (i)]}$$

Differentiating (ii) on both sides w.r.t. x, we have,

$$\frac{d^2y}{dx^2} = a \frac{dy}{dx} + b [e^{ax} (-b B \sin bx - b A \cos bx) + (B \cos bx - A \sin bx) ae^{ax}]$$

$$\Rightarrow \frac{d^2y}{dx^2} = a \frac{dy}{dx} - b^2 \{Ae^{ax} \cos bx + Be^{ax} \sin bx\} + a \{be^{ax} (B \cos bx - A \sin bx)\}$$

$$\Rightarrow \frac{d^2y}{dx^2} = a \frac{dy}{dx} - b^2 y + a \left(\frac{dy}{dx} - ay \right) \text{ [using (i) and (ii)]}$$

$$\Rightarrow \frac{d^2y}{dx^2} - 2a \frac{dy}{dx} + (a^2 + b^2) y = 0, \text{ which is the required differential equation.}$$

Hence, the given functional relation is a solution of the given differential equation.

9. According to the question,

$$\vec{a} = \hat{i} + 2\hat{j} + \hat{k},$$

$$\vec{b} = 2\hat{i} + \hat{j} \text{ and}$$

$$\vec{c} = 3\hat{i} - 4\hat{j} - 5\hat{k}$$

$$\text{Now, } \vec{a} - \vec{b} = (i + 2\hat{j} + \hat{k}) - (2\hat{i} + \hat{j}) = -\hat{i} + \hat{j} + \hat{k}$$

$$\text{Now, } \vec{c} - \vec{b} = (3\hat{i} - 4\hat{j} - 5\hat{k}) - (2\hat{i} + \hat{j}) = \hat{i} - 5\hat{j} - 5\hat{k}$$

Now, a vector perpendicular to $(\vec{a} - \vec{b})$ and $(\vec{c} - \vec{b})$ is given by

$$\vec{(a - b)} \times \vec{(c - b)} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & 1 \\ 1 & -5 & -5 \end{vmatrix}$$

$$= \hat{i}(-5 + 5) - \hat{j}(5 - 1) + \hat{k}(5 - 1)$$

$$= \hat{i}(0) - \hat{j}(4) + \hat{k}(4)$$

$$= -4\hat{j} + 4\hat{k}$$

Unit vector along $(\vec{a} - \vec{b}) \times (\vec{c} - \vec{b})$ is given by

$$\begin{aligned} & \frac{-4\hat{j} + 4\hat{k}}{|-4\hat{j} + 4\hat{k}|} \\ &= \frac{-4\hat{j} + 4\hat{k}}{\sqrt{(-4)^2 + 4^2}} \\ &= \frac{-4\hat{j} + 4\hat{k}}{\sqrt{32}} \\ &= \frac{-4\hat{j} + 4\hat{k}}{4\sqrt{2}} \\ &= -\frac{\hat{j}}{\sqrt{2}} + \frac{\hat{k}}{\sqrt{2}} \end{aligned}$$

10. According to question the given equation of the plane is $2x + y - 2z = 3$

Dividing both sides by 3, we obtain

$$\begin{aligned} \frac{2x}{3} + \frac{y}{3} + \frac{-2z}{3} &= \frac{3}{3} \\ \Rightarrow \frac{x}{\left(\frac{3}{2}\right)} + \frac{y}{3} + \frac{z}{\left(\frac{-3}{2}\right)} &= 1 \dots (1) \end{aligned}$$

We know that the equation of the plane whose intercepts on the coordinate axes are

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1 \dots (2)$$

Comparing (1) and (2), we get

$$a = \frac{3}{2}; b = 3; c = \frac{-3}{2}$$

Finding the direction cosines of the normal

The given equation of the plane is

$$2x + y - 2z = 3$$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 3$$

$$\Rightarrow \vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 3 \text{ which is the vector equation of the plane.}$$

$$\text{Also, } |\vec{n}| = \sqrt{4 + 1 + 4} = 3$$

$$\text{So, the unit vector perpendicular to } \vec{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{2\hat{i} + \hat{j} - 2\hat{k}}{3} = \frac{2}{3}\hat{i} + \frac{1}{3}\hat{j} - \frac{2}{3}\hat{k}$$

$$\text{So, the direction cosines of the normal to the plane are, } \frac{2}{3}, \frac{1}{3}, \frac{-2}{3}$$

OR

We know that line $\vec{r} = \vec{a} + k\vec{b}$ and plane $\vec{r} \cdot \vec{n} = d$ is parallel if

$$\vec{b} \cdot \vec{n} = 0$$

Given, the equation of the line

$$\vec{r} = (2\hat{i} + 5\hat{j} + 7\hat{k}) + k(\hat{i} + 3\hat{j} + 4\hat{k}) \text{ and the equation of the plane is;}$$

$$\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 7,$$

$$\text{So, } \vec{b} = \hat{i} + 3\hat{j} + 4\hat{k} \text{ and } \vec{n} = \hat{i} + \hat{j} - \hat{k}$$

$$\text{Now, } \vec{b} \cdot \vec{n}$$

$$= (\hat{i} + 3\hat{j} + 4\hat{k}) \cdot (\hat{i} + \hat{j} - \hat{k})$$

$$= 1 + 3 - 4 = 0$$

So, the line and the plane are parallel

We know that the distance (D) of a plane $\vec{r} \cdot \vec{n} = d$ from a point \vec{a} is given by

$$D = \frac{|\vec{a} \cdot \vec{n} - d|}{|\vec{n}|}$$

$$\vec{a} = (2\hat{i} + 5\hat{j} + 7\hat{k})$$

$$D = \frac{(2\hat{i} + 5\hat{j} + 7\hat{k}) \cdot (\hat{i} + \hat{j} - \hat{k}) - 7}{\sqrt{1^2 + 1^2 + (-1)^2}}$$

$$D = \frac{2 + 5 - 7 - 7}{\sqrt{1 + 1 + 1}}$$

$$D = -\frac{7}{\sqrt{3}}$$

Since the distance is always positive,

$$\text{So, } D = \frac{7}{\sqrt{3}}$$

Section C

11. To find: $\int \frac{x^2}{(x^2 + 6x - 3)} dx$

We will use following Formula ;

$$\text{i. } \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C$$

$$\text{ii. } \int \frac{f'(x)}{f(x)} dx = \log |f(x)| + C$$

Given equation can be rewritten as following:

$$\Rightarrow \int \frac{x^2 + (6x - 3) - (6x - 3)}{(x^2 + 6x - 3)} dx$$

$$\Rightarrow \int \frac{(x^2 + 6x - 3) - (6x - 3)}{(x^2 + 6x - 3)} dx$$

$$\Rightarrow x - \int \frac{6x - 3}{x^2 + 6x - 3} dx$$

$$\text{Let } I = \int \frac{6x - 3}{x^2 + 6x - 3} dx \dots \text{(ii)}$$

Using partial fractions,

$$(6x - 3) = A \left(\frac{d}{dx} (x^2 + 6x - 3) \right) + B$$

$$6x - 3 = A(2x + 6) + B$$

Equating the coefficients of x,

$$6 = 2A$$

$$A = 3$$

$$\text{Also, } -3 = 6A + B$$

$$\Rightarrow B = -21$$

Substituting in (I),

$$\Rightarrow \int \frac{3(2x+6) - 21}{(x^2 + 6x - 3)} dx$$

$$\Rightarrow 3 \times \log |x^2 + 6x - 3| + C_1 - 21 \int \frac{1}{(x+3)^2 - (\sqrt{12})^2} dx$$

$$\Rightarrow 3 \times \log |x^2 + 6x - 3| + C_1 - 21 \times \frac{1}{2\sqrt{12}} \times \log \left| \frac{x+3-\sqrt{12}}{x+3+\sqrt{12}} \right| + C_2$$

$$I = 3 \log |x^2 + 6x - 3| - \frac{7\sqrt{3}}{4} \times \log \left| \frac{x+3-2\sqrt{3}}{x+3+2\sqrt{3}} \right| + C$$

Therefore,

$$\int \frac{x^2}{(x^2 + 6x - 3)} dx = x - 3 \log |x^2 + 6x - 3| + \frac{7\sqrt{3}}{4} \times \log \left| \frac{x+3-2\sqrt{3}}{x+3+2\sqrt{3}} \right| + c$$

12. We have


$$x^2 + y^2 = 1 \dots \text{(i)}$$

$$\text{and } (x - 1)^2 + y^2 = 1 \dots \text{(ii)}$$

From (i) and (ii) we get point of Intersection as

$$A \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right), D \left(\frac{1}{2}, -\frac{\sqrt{3}}{2} \right)$$

As shown in fig.



Required Area , (Area OABD)

$$\begin{aligned}
 &= 2 \left[\int_0^{1/2} y dx + \int_{1/2}^1 y dx \right] \\
 &= 2 \left[\int_0^{1/2} \sqrt{1 - (x-1)^2} dx + \int_{1/2}^1 \sqrt{1 - x^2} dx \right] \\
 &= 2 \left[\left(\frac{x-1}{2} \sqrt{1 - (x-1)^2} + \frac{1}{2} \sin^{-1} \left(\frac{x-1}{1} \right) \right) \Big|_0^{1/2} + 2 \left[\frac{x}{3} \sqrt{1 - x^2} + \frac{1}{2} \sin^{-1} \left(\frac{x}{1} \right) \right] \Big|_{1/2}^1 \right] \\
 &= \left[(x-1) \sqrt{1 - (x-1)^2} + \sin^{-1}(x-1) \right]_0^{1/2} + \left[x \sqrt{1 - x^2} + \sin^{-1}(x) \right]_{1/2}^1 \\
 &= \left[-\frac{\sqrt{3}}{4} + 8 \sin^{-1} \left(-\frac{1}{2} \right) - \sin^{-1}(-1) \right] + \left[\sin^{-1}(1) - \frac{\sqrt{3}}{4} - \sin^{-1} \left(\frac{1}{2} \right) \right] \\
 &= \left[-\frac{\sqrt{2}}{4} - \frac{\pi}{6} + \frac{\pi}{2} \right] + \left[\frac{\pi}{2} - \frac{\sqrt{3}}{4} - \frac{\pi}{6} \right] = \left(\frac{2x}{3} - \frac{\sqrt{3}}{2} \right) \text{sq. units}
 \end{aligned}$$

OR

The given curves are $y = \sqrt{4 - x^2}$ and $x^2 + y^2 - 4x = 0$,

Now, $y = \sqrt{4 - x^2} \Rightarrow x^2 + y^2 = 4 \dots (i)$

This represents a circle with centre O(0,0) and radius = 2 units.

Also,

$x^2 + y^2 - 4x = 0 \Rightarrow (x-2)^2 + y^2 = 4 \dots (ii)$

This represents a circle with centre B(2, 0) and radius = 2 units

Solving (i) and (ii), we get

$$(x-2)^2 = x^2$$


$$\Rightarrow x^2 - 4x + 4 = x^2$$

$$\Rightarrow x = 1$$

$$\therefore y^2 = 3 \Rightarrow y = \pm \sqrt{3}$$

Thus, the given circles intersect at $A(1, \sqrt{3})$ and $C(1, -\sqrt{3})$

A rough sketch of the curves is,



\therefore Required area

= Area of shaded region OABO

$$\begin{aligned}
 &= \int_0^1 \sqrt{4 - (x-2)^2} dx + \int_1^2 \sqrt{4 - x^2} dx \\
 &= \left[\frac{1}{2} (x-2) \sqrt{4 - (x-2)^2} + \frac{4}{2} \sin^{-1} \left(\frac{x-2}{2} \right) \right]_0^1 \\
 &+ \left[\frac{1}{2} x \sqrt{4 - x^2} + \frac{4}{2} \sin^{-1} \left(\frac{x}{2} \right) \right]_1^2 \\
 &= \left[-\frac{\sqrt{3}}{2} + 2 \sin^{-1} \left(-\frac{1}{2} \right) \right] - [0 + 2 \sin^{-1}(-1)] \\
 &+ \left(0 - \frac{1}{2} \sqrt{3} \right) + 2 \left[\sin^{-1}(1) - \sin^{-1} \left(\frac{1}{2} \right) \right]
 \end{aligned}$$

$$= -\frac{\sqrt{3}}{2} - 2 \times \frac{\pi}{6} + 2 \times \frac{\pi}{2} - \frac{\sqrt{3}}{2} + 2 \times \frac{\pi}{2} - 2 \times \frac{\pi}{6}$$

$$= -\sqrt{3} + 2\pi - \frac{2\pi}{3}$$

$$\therefore A = \left(\frac{4\pi}{3} - \sqrt{3}\right) \text{ Square units}$$

13. The given equations of lines are

$$\frac{x-(a-d)}{\alpha-\delta} = \frac{y-a}{\alpha} = \frac{z-(a+d)}{\alpha+\delta} \dots(i)$$

$$\frac{x-(b-c)}{\beta-\gamma} = \frac{y-b}{\beta} = \frac{z-(b+c)}{\beta+\gamma} \dots(ii)$$

We know that the lines

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and } \frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$$

$$\text{are coplanar, therefore } \Leftrightarrow \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Here, $x_1 = (a - d)$, $y_1 = a$, $z_1 = (a + d)$;

$x_2 = (b - c)$, $y_2 = b$, $z_2 = (b + c)$;

$a_1 = (\alpha - \delta)$, $b_1 = \alpha$, $c_1 = (\alpha + \delta)$ and

$a_2 = (\beta - \gamma)$, $b_2 = \beta$, $c_2 = (\beta + \gamma)$

$$\begin{aligned} \therefore & \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} \\ &= \begin{vmatrix} (b-c) - (a-d) & b-a & (b+c) - (a+d) \\ \alpha-\delta & \alpha & \alpha+\delta \\ \beta-\gamma & \beta & \beta+\gamma \end{vmatrix} \\ &= \begin{vmatrix} 2(b-a) & b-a & b+c-a+d \\ 2\alpha & \alpha & \alpha+\delta \\ 2\beta & \beta & \beta+\gamma \end{vmatrix} \quad \{C_1 \rightarrow C_1 + C_3\} \end{aligned}$$

Therefore, the given lines are coplanar.

Equation of the plane containing the given lines is given by

$$\begin{vmatrix} x - (a-d) & y - a & z - (a+d) \\ \alpha - \delta & \alpha & \alpha + \delta \\ \beta - \gamma & \beta & \beta + \gamma \end{vmatrix} = 0$$

Applying $C_1 \rightarrow C_1 + C_3 - 2C_2$, we get

$$\begin{vmatrix} x - z - 2y & y - a & z - (a+d) \\ 0 & \alpha & \alpha + \delta \\ 0 & \beta & \beta + \gamma \end{vmatrix} = 0$$

$$\Rightarrow (x + z - 2y) - [\alpha(\beta + \gamma) - \beta(\alpha + \delta)] = 0$$

$$\Rightarrow (x + z - 2y)(\alpha\gamma - \beta\delta) = 0$$

$$\Rightarrow x + z - 2y = 0$$

Therefore, the required equation of the plane is $x + z - 2y = 0$.

CASE-BASED/DATA-BASED

14. Let X represents the number of diamond cards among the five cards drawn. Since, the drawing card is with replacement, so the trials are Bernoulli trials. In a well-shuffled deck of 52 cards, there are 13 diamond cards.

Now, $p = P(\text{success}) = P(\text{a diamond card is drawn})$

$$= \frac{13}{52} = \frac{1}{4}$$

$$\text{and } q = p(\text{failure}) = 1 - p = 1 - \frac{1}{4} = \frac{3}{4}$$

Thus, X has a binomial distribution with

$$n = 5, p = \frac{1}{4} \text{ and } q = \frac{3}{4}$$

$$\text{Therefore, } P(X = r) = {}^n C_r p^r q^{n-r},$$

where $r = 0, 1, 2, 3, 4, 5$

$$P(X = r) = {}^5 C_r \left(\frac{1}{4}\right)^r \left(\frac{3}{4}\right)^{5-r}$$

i. P(all the five cards are diamonds) = P(X = 5)

$$= {}^5C_5 p^5 q^0 = 1p^5 = \left(\frac{1}{4}\right)^5 = \frac{1}{1024}$$

ii. P(none is a diamond) = P(X = 0)

$$= {}^5C_0 p^0 q^5 = (q)^5$$

$$= \left(\frac{3}{4}\right)^5 = \frac{243}{1024}$$