



# **CHEMICAL EQUILIBRIUM**

#### Types of Reaction

	Reversible	Irreversible
1	A reaction in which not only the reactants react to form products but also the products react to form reactants under the same conditions is called reversible reaction.	A reaction which cannot take place in the reverse direction is called an irreversible reaction
1	A few examples are given as $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ $N_2O_4(g) \rightleftharpoons 2NO_2(g)$	A few examples are given as $AgNO_{3}(aq) + NaCl(aq) \rightarrow AgCl(s) + NaNO_{3}(aq)$ $BaCl_{2}(aq) + Na_{2}SO_{4}(aq) \rightarrow BaSO_{4}(s) + 2NaCl(aq)$

## **Characteristics of Equilibrium State**

- \* A reaction (or a process) is said to be in equilibrium when the rate of forward reaction (process) becomes equal to the rate of backward reaction (process).
- At equilibrium the concentration of each of the reactants and products become constant.
- An equilibrium is dynamic in nature and not static i.e., even after equilibrium is attained, the forward as well as the backward reaction take place but at equal speeds.
- \* A chemical equilibrium can be established only if none of the products is allowed to escape out.
- Chemical equilibrium can be attained from either direction.
- When reaction attain equilibrium at certain temperature and pressure,  $\Delta_r G = 0$
- Catalyst increases rate of reaction in forward as well as backward direction upto same extent.

#### Law of Mass Action

It was put forward by Guldberg and Waage. It sates that the rate at which a substance reacts is directly proportional to its active mass and hence the rate at which substances react together is directly proportional to the product of their active masses. Active mass means molar concentration.

#### Law of Chemical Equilibrium

For the reaction  $aA(aq) + bB(aq) \Longrightarrow xX(aq) + yY(aq)$   $K_c = \frac{[X]^x [Y]^y}{[A]^a [B]^b}$ 

Where  $K_c$  is called equilibrium constant at constant temperature.

### Types of Equilibrium Constant

 $K_c$  = Equilibrium constant in terms of molar concentration =  $\frac{[X]^x[Y]^y}{[A]^a[B]^b}$ 

 $K_p$  = Equilibrium constant in terms of pressure =  $\frac{p_X^x \cdot p_Y^y}{p_A^a \cdot p_B^b}$ 

 $K_x$  = Equilibrium constant in terms of mole fraction =  $\frac{(X_x)^x (X_y)^y}{(X_A)^a (X_B)^b}$ 

#### **Unit of Equilibrium Constant**

Unit of  $K_p = (atm)^{\Delta n_g}$  (where  $\Delta n_g = change in gas mole of reaction)$ 

Unit of  $K_c = M^{\Delta n_g}$ 

Unit of  $K_x$  = Unitless

Relationship between  $K_p$  and  $K_c$ :  $K_p = K_c (RT)^{\Delta n_g}$ 

where R = universal gas constant, T = temperature

Relationship between  $K_p$  and  $K_x$ :  $K_p = K_x(P)^{\Delta n_g}$ 

where P = equilibrium pressure in container.

#### Types of Chemical Equilibria

Homogeneous equilibrium. When in an equilibrium reaction, all the reactants and products are present in the same phase (i.e. gaseous or liquid), it is called homogeneous equilibrium.

$$PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$$
;  $CH_3COOH(l) + C_2H_5OH(l) \rightleftharpoons CH_3COOC_2H_3(l) + H_2O(l)$  etc.

Heterogeneous equilibrium. When in an equilibrium reaction, the reactants and the products are present in two or more than two phases, it is called a heterogeneous equilibrium.

The second se

$$PbCl_2(s) \rightleftharpoons Pb^{2+}(aq) + 2Cl^{-}(aq)$$
;  $CaO(s) + CO_2(g) \rightleftharpoons CaCO_3(s)$  etc.

### Expression of K for Equilibrium Reaction

Homogeneous equilibria in gases

Example: (i)  $NH_3(aq) + H_2O(l) \rightleftharpoons NH_4^+(aq) + OH^-(aq)$ 

(ii) 
$$3O_2(g) \rightleftharpoons 2O_3(g)$$
  
(iii)  $C_3H_8(g) + 5O_2(g) \rightleftharpoons 3CO_2(g) + 4H_2O(g)$ 

Equilibrium constant expression for them are

(i) 
$$K_c = \frac{[NH_4^+(aq)][OH^-(aq)]}{[NH_3(aq)]}$$
 ; [] represents concentration in mol/ litre at equilibrium.

(ii) 
$$K_c = \frac{[{\rm O}_3]^2}{[{\rm O}_2]^3}$$
;  $K_p = \frac{P_{{\rm O}_3}^2}{P_{{\rm O}_2}^3}$  (iii)  $K_C = \frac{[{\rm CO}_2]^3 [{\rm H}_2 {\rm O}]^4}{[{\rm C}_3 {\rm H}_8] [{\rm O}_2]^5}$ ;  $K_p = \frac{P_{{\rm CO}_2}^3 \cdot P_{{\rm H}_2 {\rm O}}^4}{P_{{\rm C}_3 {\rm H}_8} \cdot P_{{\rm O}_2}^5}$ 

**Note:** Equilibrium constant for gaseous homogeneous equilibrium can be expressed in two ways  $K_p$  and  $K_C$ . This means value of equilibrium constant depends upon choice of standard state in which concentration of reactants and products are expressed.

#### \* Heterogeneous equilibria

If reactants and product are found in two or more phases, the equilibria describing them is called heterogeneous equilibrium.

Example: (i) 
$$AgI(s) \rightleftharpoons Ag^{+}(aq) + I^{-}(aq)$$
  
(ii)  $CaO(s) + CO_{2}(g) \rightleftharpoons CaCO_{3}(s)$   
(iii)  $Br_{2}(l) \rightleftharpoons Br_{2}(g)$ 

Equilibrium expression for them can be written as

(i) 
$$K_C = [Ag^+(aq)][I^-(aq)]$$
 (ii)  $K_P = \frac{1}{P_{CO_2}}$  ;  $K_C = \frac{1}{[CO_2(g)]}$ 

(iii) 
$$K_P = P_{\text{Br}_2}(g)$$
 ;  $K_C = [\text{Br}_2(g)]$ 

#### Predicting the Extent of a Reaction

High value of equilibrium constant indicates that product(s) concentration is high and its low value indicates that concentration of the product(s) in equilibrium mixture is low.

Large value of  $K_P$  or  $K_C$  (larger than about  $10^3$ ), favour the products strongly. For intermediate values of K (approximately in the range of  $10^{-3}$  to  $10^3$ ), the concentrations of reactants and products are comparable. Small values of equilibrium constant (smaller than  $10^{-3}$ ), favour the reactants strongly.

#### Predicting the Direction of the Reaction

For this purpose, we calculate the reaction quotient, Q. The reaction quotient is defined in the same way as the equilibrium constant (with molar concentrations to give  $Q_C$ , or with partial pressure to give  $Q_P$ ) at any stage of reaction. For a general reaction

$$aA + bB \rightleftharpoons cC + dD$$

$$Q_C = \frac{[C]^c[D]^d}{[A]^a[B]^d}$$

Then, if  $Q_C > K_C$ , the reaction will proceed towards direction of reactants.

If  $Q_C < K_c$ , the reaction will move towards direction of the products.

If  $Q_C = K_c$ , the reaction mixture is at equilibrium

#### Important Point about Equilibrium Constant

If the reaction is reversed, the value of equilibrium constant is inversed e.g.,

 $CH_3COOH(aq) + C_2H_5OH(aq) \rightleftharpoons CH_3COOC_2H_5(aq) + H_2O(l)$ , Equilibrium constant = K then for  $CH_3COOC_2H_5(aq) + H_2O(l) \rightleftharpoons CH_3COOH(aq) + C_2H_5OH(aq)$ , Equilibrium constant = 1/K

If the reaction is divided by 2, equilibrium constant is the square root of the original e.g.

for 
$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$
, Equilibrium constant =  $K$   
then for  $\frac{1}{2}N_2(g) + \frac{3}{2}H_2(g) \rightleftharpoons NH_3(g)$ , Equilibrium constant =  $\sqrt{K}$ 

If the reaction is multiplied by 2, equilibrium constant is the square of the original.

❖ If the reaction is written in two steps, equilibrium constant is equal to the product of the equilibrium constants of the step reactions e.g., if for

$$N_2(g) + 2O_2(g) \rightleftharpoons 2NO_2(g)$$
, Equilibrium constant = K

and for the same reaction taking place in steps i.e.,

$$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$$
, Equilibrium constant =  $K_1$ 

and  $NO(g) + O_2(g) \rightleftharpoons 2NO_2(g)$ , Equilibrium constant =  $K_2$ then  $K = K_1 \times K_2$ 

• Effect of temperature: According to van't Hoff equation,  $\log \frac{K_2}{K_1} = \frac{\Delta H}{2.303R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$ 

where  $K_1$  and  $K_2$  are the equilibrium constants at temperatures  $T_1$  and  $T_2$  respectively and  $\Delta H$  is the molar enthalpy change in the temperature range  $T_1$  to  $T_2$ .

For exothermic reaction, as temperature increases K decreases.

For endothermic reaction, as temperature increases K increases.

#### Relationship Between Degree of Dissociation (a) and Vapour Density

\* For dissociation reaction:  $A_n(g) \rightleftharpoons nA(g)$ 

Degree of dissociation (
$$\alpha$$
) =  $\frac{\text{No. of moles dissociated}}{\text{Total no. of moles taken}}$   
=  $\frac{D-d}{d(n-1)} = \frac{M_t - M_0}{M_0(n-1)}$ 

where

 $M_t$  = theoretical (calculated) molecular weight

 $M_0$  = observed (experimental) molecular weight

D = theoretical vapour density, d = observed vapour density.

n = number of moles of product formed from 1 mole reactant

Density of gas mixture = 
$$\frac{PM}{RT}$$

· For polymerization reaction

$$nA(g) \rightleftharpoons A_n(g)$$
, where  $n \ge 2$ 

$$\alpha = \frac{D-d}{d\left(\frac{1}{n}-1\right)} = \frac{M_t - M_0}{M_0\left(\frac{1}{n}-1\right)}$$

#### Le Chatelier's Principle

"If a system in equilibrium is subjected to a change of concentration, temperature or pressure, the equilibrium shifts in a direction so as to undo the effect of the change imposed."

· Effect of change in concentration on equilibrium

As we add or remove reactant (or product) the ratio of equilibrium concentration become 'Q' (reaction quotient)

- Q < K: equilibrium will shift in forward direction.
- Q > K: equilibrium will shift in backward direction.
- · Effect of change in pressure

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, the volume decreases proportionately. The equilibrium will shift in the direction in which there is decrease in number of moles *i.e.*, towards the direction in which there is decrease in volume.

- \* Effect of change in pressure on melting point: There are two types of solids:
  - (i) Solids whose volume decreases on melting, e.g., ice, diamond, carborundum, magnesium nitride and quartz.

The process of melting is facilitated at high pressure, thus melting point is lowered.

- (ii) Solid whose volume increase on melting, e.g., Fe, Cu, Ag, Au, etc.

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

- \* Solubility of substances: When solid substance are dissolved in water, either heat is evolved (exothermic) or heat is absorbed (endothermic).
  - (i) For endothermic solubility process, solubility increase with increase in temperature.
  - (ii) For exothermic solubility process, decrease with increase in temperature.
- Solubility of gases in liquids: When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.
- Effect of temperature: For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.
- Addition of inert gas
  - (i) For reactions in which  $n_p = n_r$  there is no effect of adding an inert gas at constant volume or at constant pressure on the equilibrium.
  - (ii) For reactions in which  $n_p \neq n_r$ , there is no effect of adding inert gas on the equilibrium at constant volume but at constant pressure, equilibrium shifts towards larger mole side.

Applying Le Chatelier's principle, the favourable conditions for the dissociation of NH<sub>3</sub> by Haber's process

i.e., 
$$N_2(g) + 3H_2(g) \implies 2NH_3(g), \Delta H = -92.5 \text{ kJ}$$

are (i) High temperature (ii) Low pressure (iii) Removal of  $N_2$  and  $H_2$  (iv) Addition of inert gas at constant pressure.



#### \* Physical Equilibrium

Only physical changes are involved.

Example: (i) Solid liquid equilibria

$$H_2O(s) \rightleftharpoons H_2O(l)$$

At equilibrium: Net rate of conversion of ice into water = Net rate of conversion of water into ice.

For  $H_2O(l)$  normal freezing point is 273 K.

(ii) Liquid vapour equilibria:

$$H_2O(l) \rightleftharpoons H_2O(g)$$

Equilibrium is characterized by constant value of vapour pressure of  $\mathrm{H}_2\mathrm{O}(l)$  at certain temperature.

For H<sub>2</sub>O(l) normal boiling point is 373 K.

# Level

1.	Α	reversible	reaction	is	one	which	
••	43	ICACIDIDIC	I Cacuon	10	0110	AA 111 C11	•

(a) Proceeds in one direction

(b) Proceeds in both directions

(c) Proceeds spontaneously

- (d) All the statements are wrong
- 2. The equilibrium constant  $K_c$  for the reaction

$$P_4(g) \rightleftharpoons 2P_2(g)$$

is 1.4 at 400°C. Suppose that 3 moles of  $P_4(g)$  and 2 moles of  $P_2(g)$  are mixed in 2 litre container at 400°C. What is the value of reaction quotient  $(Q_c)$ ?

(a)  $\frac{3}{2}$ 

(b)  $\frac{2}{3}$ 

(c) 1


- (d) None of these
- 3. In a chemical reaction equilibrium is established when:
  - (a) Opposing reaction ceases
  - (b) Concentrations of reactants and product are equal
  - (c) Velocity of opposing reaction is the same as that of forward reaction
  - (d) Reaction ceases to generate heat
- **4.** The equilibrium constant for a reaction is K, and the reaction quotient is Q. For a particular reaction mixture, the ratio  $\frac{K}{Q}$  is 0.33. This means that :
  - (a) the reaction mixture will equilibrate to form more reactant species
  - (b) the reaction mixture will equilibrate to form more product species
  - (c) the equilibrium ratio of reactant to product concentrations will be 3
  - (d) the equilibrium ratio of reactant to product concentrations will be 0.33
- 5. Consider the reaction  $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$  for which  $K_c = 278 \, M^{-1}$ . 0.001 mole of each of the reagents  $SO_2(g)$ ,  $O_2(g)$  and  $SO_3(g)$  are mixed in a 1.0 L flask. Determine the reaction quotient of the system and the spontaneous direction of the system:
  - (a)  $Q_c = 1000$ , the equilibrium shifts to the right
  - (b)  $Q_c = 1000$ ; the equilibrium shifts to the left
  - (c)  $Q_c = 0.001$ ; the equilibrium shifts to the left
  - (d)  $Q_c = 0.001$ ; the equilibrium shifts to the right
- 6. In Q. No. 5, if the mixture of gases was allowed to come to equilibrium. The volume of the reaction vessel was then rapidly increased by a factor of two. As a result of the change the reaction quotient (Q<sub>c</sub>) would:
  - (a) increase because of the pressure decrease
  - (b) decrease because of the pressure decrease
  - (c) remain the same because the equilibrium constant is independent of volume
  - (d) increase because the reaction is endothermic
- 7. For the reaction  $A(g) + 3B(g) \iff 2C(g)$  at 27°C, 2 moles of A, 4 moles of B and 6 moles of C are present in 2 litre vessel. If  $K_c$  for the reaction is 1.2, the reaction will proceed in :
  - (a) forward direction


(b) backward direction


(c) neither direction


(d) none of these

- **8.** For a reversible gaseous reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  at equilibrium, if some moles of  $H_2$  are replaced by same number of moles of  $T_2$  (T is tritium, isotope of H and assume isotopes do not have different chemical properties) without affecting other parameters, then:
  - (a) The sample of ammonia obtained after sometime will be radioactive.
  - (b) Moles of  $N_2$  after the change will be different as compared to moles of  $N_2$  present before the change
  - (c) The value of  $K_p$  or  $K_c$  will change
  - (d) The average molecular mass of new equilibrium will be same as that of old equilibrium
- **9.** For the synthesis of ammonia by the reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  in the Haber's process, the attainment of equilibrium is correctly predicted by the curve









**10.** The figure shows the change in concentration of species A and B as a function of time.


The equilibrium constant  $K_c$  for the reaction  $A(g) \iff 2B(g)$  is:




(b) 
$$K < 1$$

(c) 
$$K = 1$$

(d) data insufficient



11. Attainment of the equilibrium  $A(g) \rightleftharpoons 2C(g) + B(g)$  gave the following graph. Find the correct option. (% dissociation = fraction dissociated ×100)



(a) 6.7

 $\frac{K_p \text{ (atm)}}{K_c \text{ (M)}} = 3?$ 

(a) 0.027 K

(b) 0.61

(b) 0.36 K

	(a) At $t = 5$ sec equilibrium has been reached and $K_c = 40$ (mol/litre) <sup>2</sup>				
	(b) At $t = 5$ sec equilibrium has been reached and % dissociation of A is 20%				
	(c) At $t = 5$ sec equilibrium has been reached and % dissociation of A is 30%				
	(d) None of these			_	
12.	Using molar concent	rations, what is the unit	•	?	
	•	$CH_3OH(g) \iff$	$CO(g) + 2H_2(g)$		
	(a) $M^{-2}$	(b) M <sup>2</sup>	(c) $M^{-1}$	(d) M	
13.	What is the unit of K	•			
	•	$CS_2(g) + 4H_2(g) \rightleftharpoons$	$CH_4(g) + 2H_2S(g)$		
	(a) atm	(b) atm <sup>-2</sup>	(c) atm <sup>2</sup>	(d) atm <sup>-1</sup>	
14.	What is the equilibrium	um expression for the re	eaction $P_4(s) + 5O_2(g)$	$\rightleftharpoons P_4O_{10}(s)$	
	(a) $K_c = [O_2]^5$		(b) $K_c = [P_4O_{10}] / 5[1]$	P <sub>4</sub> ][O <sub>2</sub> ]	
	(c) $K_c = [P_4O_{10}]/[P_4O_{10}]$	<sub>4</sub> ][O <sub>2</sub> ] <sup>5</sup>	(d) $K_c = 1/[O_2]^5$	•	
15.	At 527°C, the reaction	n given below has $K_c =$	4		
	•	$NH_3(g) \rightleftharpoons \frac{1}{2}$	$\frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$		
	What is the $K_p$ for the	he reaction? $N_2(g) + 3H_2(g)$	$g) \rightleftharpoons 2NH_3(g)$	٠	
	•	(800R) <sup>-2</sup>	$\begin{pmatrix} 1 \end{pmatrix}^2$		
	(a) $16 \times (800  R)^2$	(b) $\left(\frac{800R}{4}\right)^{-2}$	(c) $\left(\frac{1}{4 \times 800R}\right)$	(d) None of these	
				g) at temperature (T) is	
	$4 \times 10^{-4}$ . The value	of $K_c$ for the reaction	on $NO(g) \rightleftharpoons \frac{1}{2}N_2(g)$	$+\frac{1}{2}O_2(g)$ at the same	
	temperature is:	(b) 50 ·	(c) $2.5 \times 10^2$	(4) 0.00	
17.	(a) $4 \times 10^{-4}$ The equilibrium cons	tant K. for the following	(c) 25×10 <sup>-</sup> g reaction at 842°C is 7	(d) $0.02$ .90 × $10^{-3}$ . What is $K_p$ at	
	same temperature?		,	,	
		$\frac{1}{2}\mathrm{F}_2(g)$	$\rightleftharpoons$ $F(g)$		
	(a) $8.64 \times 10^{-5}$	(b) 8.26 × 10 <sup>-4</sup>	(c) $7.90 \times 10^{-2}$	(d) $7.56 \times 10^{-2}$	
18.	The equilibrium cons	stant $K_p$ for the following	ng reaction at 191°C is	1.24. What is $K_c$ ?	
			$g) \iff BF_3(g)$		

19. For the equilibrium  $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$ , what is the temperature at which

(d) 7.6

(d) 273 K

(c) 8.30

(c) 36.54 K

20. For the reversible reaction,

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

at 500°C, the value of  $K_p$  is  $1.44 \times 10^{-5}$  when partial pressure is measured in atmospheres. The corresponding value of  $K_p$  with concentration in mole litre<sup>-1</sup>, is:

- (a)  $1.44 \times 10^{-5}/(0.082 \times 500)^{-2}$
- (b)  $1.44 \times 10^{-5} / (8.314 \times 773)^{-2}$

(c)  $1.44 \times 10^{-5} / (0.082 \times 773)^2$ 

- (d)  $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$
- **21.** For the reaction  $CO(g) + Cl_2(g) \iff COCl_2(g)$  the value of  $\frac{K_c}{K_p}$  is equal to :
  - (a)  $\sqrt{RT}$
- (b) RT
- (c)  $\frac{1}{RT}$
- (d) 1.0
- **22.** The concentration of a pure solid or liquid phase is not included in the expression of equilibrium constant because :
  - (a) solid and liquid concentrations are independent of their quantities.
  - (b) solids and liquids react slowly.
  - (c) solids and liquids at equilibrium do not interact with gaseous phase.
  - (d) the molecules of solids and liquids cannot migrate to the gaseous phase.
- 23. A catalyst is a substance which:
  - (a) increases the equilibrium concentration of the product.
  - (b) changes the equilibrium constant of the reaction.
  - (c) shortens the time to reach equilibrium.
  - (d) supplies energy to the reaction.
- **24.** What will be the effect on the equilibrium constant on increasing temperature, if the reaction neither absorbs heat nor releases heat?
  - (a) Equilibrium constant will remain constant.
  - (b) Equilibrium constant will decrease.
  - (c) Equilibrium constant will increase.
  - (d) Can not be predicted.
- **25.** The equilibrium constant for the reaction  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$  is  $4 \times 10^{-4}$  at 200 K. In presence of a catalyst, equilibrium is attained ten times faster. Therefore, the equilibrium constant in presence of the catalyst at 200 K is:
  - (a)  $40 \times 10^{-4}$

(b)  $4 \times 10^{-4}$ 

(c)  $4 \times 10^{-3}$ 

- (d) difficult to compute without more data
- **26.** For the reaction  $H_2(g) + I_2(g) \implies 2HI(g)$ , the equilibrium constant changes with :
  - (a) total pressure

- (b) catalyst
- (c) concentration of H2 and I2
- (d) temperature

- 27. Consider the reactions
  - (i)  $2CO(g) + 2H_2O(g) \implies 2CO_2(g) + 2H_2(g)$ ; Eqm. Constant =  $K_1$
  - (ii)  $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$ ; Eqm. Constant =  $K_2$

(iii)  $CH_4(g) + 2H_2O(g) \rightleftharpoons CO_2(g) + 4H_2(g)$ ; Eqm. Constant =  $K_3$ 

Which of the following relation is correct?

- (a)  $K_3 = \frac{K_1}{K}$

- (b)  $K_3 = \frac{K_1^2}{\kappa^2}$  (c)  $K_3 = K_1 K_2$  (d)  $K_3 = \sqrt{K_1} \cdot K_2$

**28.** For the reaction  $2NO_2(g) + \frac{1}{2}O_2(g) \implies N_2O_5(g)$ , if the equilibrium constant is  $K_p$ , then the equilibrium constant for the reaction  $2N_2O_5(g) \iff 4NO_2(g) + O_2(g)$  would be :

- (a)  $K_{p}^{2}$
- (b)  $\frac{2}{K_n}$
- (c)  $\frac{1}{K_{n}^{2}}$
- (d)  $\frac{1}{\sqrt{K_n}}$

**29.** The equilibrium constant  $(K_c)$  for the reaction

$$2HCl(g) \iff H_2(g) + Cl_2(g)$$

is  $4 \times 10^{-34}$  at 25°C. What is the equilibrium constant for the reaction?

$$\frac{1}{2}$$
H<sub>2</sub>(g) +  $\frac{1}{2}$ Cl<sub>2</sub>(g)  $\implies$  HCl(g)

- (b)  $2.5 \times 10^{33}$
- (c)  $5 \times 10^6$
- (d) None of these

30. At a certain temperature, the following reactions have the equilibrium constants as shown below:

$$S(s) + O_2(g) \iff SO_2(g); \quad K_c = 5 \times 10^{52}$$
  
 $2S(s) + 3O_2(g) \iff 2SO_3(g); \quad K_c = 10^{29}$ 

What is the equilibrium constant  $K_c$  for the reaction at the same temperature?

$$2SO_2(g) + O_2(g) \iff 2SO_3(g)$$

- (a)  $2.5 \times 10^{76}$
- (b)  $4 \times 10^{23}$
- (c)  $4 \times 10^{-77}$
- (d) None of these

31. Consider the following gaseous equilibria given below:

- (I)  $N_2 + 3H_2 \rightleftharpoons 2NH_3$ ; Eqm. Constant =  $K_1$
- (II)  $N_2 + O_2 \implies 2NO$ ; Eqm. Constant =  $K_2$
- (III)  $H_2 + \frac{1}{2}O_2 \iff H_2O$ ; Eqm. Constant =  $K_3$

The equilibrium constant for the reaction,  $2NH_3 + \frac{5}{2}O_2 \iff 2NO + 3H_2O$  in terms of  $K_1$ ,  $K_2$ and  $K_3$  will be:


- (a)  $K_1K_2K_3$

- (b)  $\frac{K_1K_2}{K_2}$  (c)  $\frac{K_1K_3^2}{K_2}$  (d)  $\frac{K_2K_3^3}{K_1}$

**32.** In the reaction  $X(g) + Y(g) \iff 2 Z(g)$ , 2 mole of X, 1 mole of Y and 1 mole of Z are placed in a 10 litre vessel and allowed to reach equilibrium. If final concentration of Z is 0.2 M, then  $K_c$  for the given reaction is:

- (a) 1.60
- (b)  $\frac{80}{2}$
- (c)  $\frac{16}{3}$

(d) None of these



(d) None of these

(a)  $\frac{1}{4}$ 

reaction will be:

	(a) $\frac{1}{4}$	(b) 2	(c) 3	(d) 4	
44.	After equilibrium is	reached, excess of A	1 mole of I <sub>2</sub> and 0.5 molgNO <sub>3</sub> gave 0.25 mol	ole of I <sup>-</sup> in one litre flask. e of yellow precipitate.	
	Equilibrium constant i (a) 1.33	is: (b) 2.66	(c) 2.0	(d) 3.0	
45.	At 87°C, the following	; equilibrium is establis	shed.		
		$H_2(g) + S(s) \rightleftharpoons$	$H_2S(g); K_c = 0.08$		
	If 0.3 mole hydrogen a concentration of H <sub>2</sub> S			L vessel, what will be the	
	(a) 0.011 M	(b) 0.022 M	(c) 0.044 M	(d) 0.08 M	
46.				sure of SO <sub>2</sub> , O <sub>2</sub> and SO <sub>3</sub>	
	are 0.662, 0.10 and 0 that the equilibrium of	oncentrations of SO <sub>2</sub> a	and SO <sub>3</sub> are equal?	tial pressure of oxygen so	
	(a) 0.4 atm	(b) 1.0 atm	(c) 0.8 atm	(d) 0.25 atm	
47.	When heated, ammor			*¢	
•	•	$NH_4COONH_2(s) =$	$\Rightarrow$ 2NH <sub>3</sub> (g) + CO <sub>2</sub> (g)		
	At a certain temperature, the equilibrium pressure of the system is 0.318 atm. $K_p$ for the reaction is:				
	(a) 0.128	(b) 0.426	(c) $4.76 \times 10^{-3}$	(d) None of these	
48.	In a system $A(s) \iff$	2B(g) + 3C(g), if the c	oncentration of C at equ	illibrium is increased by a	
			centration of B to chan		
	(a) two times the ori	ginal value	(b) one half of its or		
	(c) $2\sqrt{2}$ times to the	original value	(d) $\frac{1}{2\sqrt{2}}$ times the or	riginal value	
49.	$A + B \rightleftharpoons C + D$ . If fin concentration of D we reaction?	nally the concentration ill be twice of that of A	of $A$ and $B$ are both $A$ then what will be the	equal but at equilibrium e equilibrium constant of	
	(a) $\frac{4}{9}$	(b) $\frac{9}{4}$	(c) $\frac{1}{9}$	(d) 4	
50.	The equilibrium const	ant $K_c$ for the reaction	$SO_2(g) + NO_2(g) \iff$	$SO_3(g) + NO(g)$ is 16. If	
	1 mole of each of all t	he four gases is taken i	in 1 dm³ vessel, the equ	illibrium concentration of	
٠	NO would be:				
	(a) 0.4 M	(b) 0.6 M	(c) 1.4 M	(d) 1.6 <i>M</i>	
			•	,	

**42.** For the reaction  $2A(g) \iff B(g) + 3C(g)$ , at a given temperature,  $K_c = 16$ . What must be the volume of the flask, if a mixture of 2 mole each of A, B and C exist in equilibrium?

**43.** One mole of pure ethyl alcohol was treated with one mole of pure acetic acid at 25°C. One-third of the acid changes into ester at equilibrium. The equilibrium constant for the

(c) 1

51.	On increasing the temperature, the	rate of a reaction:	•		
	(a) always increases				
,	(b) always decreases				
	(c) first increases and then decreas		,		
	(d) may increase or decrease deper		ne reaction		
52.	A catalyst increases the rate of a re				
	(a) increasing the activation energy				
	(b) increasing the value of rate con	istants $(k_f \text{ and } k_b)$			
	(c) increasing the enthalpy change				
E 2	(d) decreasing the enthalpy change				
55.	At a certain temperature, only 50% I		m in the following reaction:		
	2H	$I(g) \iff H_2(g) + I_2(g)$			
	The equilibrium constant for this re	action is :			
	(a) 0.25 (b) 1.0	(c) 3.0	(d) 0.5		
54.	The equilibrium constant $K_p$ for the	e reaction			
	$H_2(g) + 0$	$CO_2(g) \iff H_2O(g) + CO(g)$	(g)		
	is 4.0 at 1660°C. Initially 0.80 mole	H <sub>2</sub> and 0.80 mole CO <sub>2</sub> are	injected into a 5.0 litre flask.		
	What is the equilibrium concentration	on of CO <sub>2</sub> (g)?	•		
	(a) $0.533 M$ (b) $0.0534 M$	3 5	(d) None of these		
55.	At 273 K and 1 atm, 10 litre of N <sub>2</sub> C	O <sub>4</sub> decomposes to NO <sub>2</sub> acco	ording to equation		
	N	$_{2}O_{4}(g) \rightleftharpoons 2NO_{2}(g)$			
	What is degree of dissociation ( $\alpha$ ) w	hen the original volume is 2	25% less than that of existing		
	volume?	0			
	(a) 0.25 (b) 0.33	(c) 0.66	(d) 0.5		
56.	The equilibrium constant for the re-	action $CO(g) + H_2O(g) \rightleftharpoons$	$CO_2(g) + H_2(g)$ is 5. How		
	many moles of CO <sub>2</sub> must be added to	o 1 litre container already co	ontaining 3 moles each of CO		
	and H <sub>2</sub> O to make 2M equilibrium co				
	(a) 15 (b) 19	(c) 5	(d) 20		
57.	A nitrogen-hydrogen mixture initiall	y in the molar ratio of 1:3	reached equilibrium to form		
	ammonia when 25% of the N <sub>2</sub> and F atm, the partial pressure of ammoni	a at the equilibrium was	ressure of the system was 21		
		(c) 2.0 atm	(d) 1.5 atm		
58.	Ammonia under a pressure of 15 at	m at 27°C is heated to 347	7°C in a closed vessel in the		
	presence of a catalyst. Under the conditions, NH <sub>3</sub> is partially decomposed according to the				
	equation,	•			
	$2NH_3 \rightleftharpoons N_2 + 3H_2$ the vessel is suc	h that the volume remains	effectively constant where as		
	pressure increases to 50 atm. Calcul-	ate the percentage of NH <sub>3</sub>	actually decomposed:		
	(a) 65% (b) 61.3%	(c) 62.5%	(d) 64%		

381731	C. 1. 1. 2. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.			243
59.				eric conditions at 25°C. i $N_2O_4(g) \rightleftharpoons 2NO_2(g)$
	$(K_p = 0.14)$ is reached (a) $1.8 \times 10^2$	1 after some time : (b) $2.8 \times 10^2$	(c) 0.034	(d) $2.8 \times 10^{-2}$
60.				n, it was found that 60% atmosphere, the partial
61.	(a) 0.52 atm $N_2(g) + 3H_2(g) \implies$	(b) 0.21 atm 2NH <sub>3</sub> (g)	(c) 0.41 atm	(d) 0.82 atm
	reacted. If the equilib	orium pressure is P, the	partial pressure of NH <sub>3</sub>	
	(a) P	(L) P	(a) P	(a) P

**62.** 2.0 mole of PCl<sub>5</sub> were introduced in a vessel of 5.0 L capacity of a particular temperature. At equilibrium, PCl<sub>5</sub> was found to be 35% dissociated into PCl<sub>3</sub> and Cl<sub>2</sub>. The value of K<sub>c</sub> for the

reaction 
$$PCl_2(g) + Cl_2(g) \implies PCl_2(g)$$

**63.** At certain temperature compound  $AB_2(g)$  dissociates according to the reaction

$$2AB_2(g) \iff 2AB(g) + B_2(g)$$

With degree of dissociation  $\alpha$ , which is small compared with unity. The expression of  $K_p$ , in terms of  $\alpha$  and initial pressure P is:

(a) 
$$P \frac{\alpha^3}{2}$$

(b) 
$$\frac{P\dot{\alpha}^2}{3}$$

(c) 
$$P \frac{\alpha^3}{3}$$

(d) 
$$\frac{P\alpha^2}{2}$$

**64.** For the reaction

 $H_2(g) + CO_2(g) \rightleftharpoons CO(g) + H_2O(g)$ . If the initial concentration of  $[H_2] = [CO_2]$  and x moles/litre of hydrogen is consumed at equilibrium, the correct expression of  $K_p$  is:

(a) 
$$\frac{x^2}{(1-x)^2}$$

(b) 
$$\frac{(1+x)^2}{(1-x)^2}$$

(c) 
$$\frac{x^2}{(2+x)^2}$$

$$(d) \frac{x^2}{1-x^2}$$

**65.** If  $D_T$  and  $D_O$  are the theoretical and observed vapour densities at a definite temperature and  $\alpha$ be the degree of dissociation of a substance. Then,  $\alpha$  in the terms of  $D_0$ ,  $D_T$  and n (number of moles of products formed from 1 mole reactant) is calculated by the formula:


(a) 
$$\alpha = \frac{D_O - D_T}{(1-n) D_T}$$

(b) 
$$\alpha = \frac{D_T - D_O}{(n-1) D_T}$$

(a) 
$$\alpha = \frac{D_O - D_T}{(1 - n) D_T}$$
 (b)  $\alpha = \frac{D_T - D_O}{(n - 1) D_T}$  (c)  $\alpha = \frac{D_T - D_O}{(n - 1) D_O}$  (d)  $\alpha = \frac{D - D_T}{(n - 1) D_T}$ 

(d) 
$$\alpha = \frac{D - D_T}{(n-1) D_T}$$

**66.** For the dissociation of PCl<sub>5</sub> into PCl<sub>5</sub> and Cl<sub>2</sub> in gaseous phase reaction, if d is the observed vapour density and D the theoretical vapour density with ' $\alpha$ ' as degree of dissociation. Variation of D/d with ' $\alpha$ ' is given by which graph?



- 67. At 27°C and 1 atm pressure, N<sub>2</sub>O<sub>4</sub> is 20% dissociation into NO<sub>2</sub>. What is the density of equilibrium mixture of N<sub>2</sub>O<sub>4</sub> and NO<sub>2</sub> at 27°C and 1 atm?
  - (a) 3.11 g/litre
- (b) 2.11 g/litre
- (c) 4.5 g/litre
- (d) None of these
- **68.**  $COCl_2$  gas dissociates according to the equation,  $COCl_2(g) \rightleftharpoons CO(g) + Cl_2(g)$ . When heated to 700 K the density of the gas mixture at 1.16 atm and at equilibrium is 1.16 g/litre. The degree of dissociation of  $CO_2$  at 700 K is:
  - (a) 0.28
- (b) 0.50
- (c) 0.72
- (d) 0.42
- **69.** The degree of dissociation of I<sub>2</sub> molecule of 1000°C and under atmospheric pressure is 40% by volume. If the dissociation is reduced to 20% at the same temp, total equilibrium pressure on the gas is:
  - (a) 1.57 atm
- (b) 2.57 atm
- (c) 3.57 atm
- (d) 4.57 atm
- **70.** Determine the value of equilibrium constant  $(K_c)$  for the reaction

$$A_2(g) + B_2(g) \iff 2AB(g)$$

If 10 moles of  $A_2$ ; 15 moles of  $B_2$  and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M:

- (a) 4.5
- (b) 1.5
- (c) 0.6
- (d) None of these

71. At 87°C, the following equilibrium is established

$$H_2(g) + S(s) \iff H_2S(g); K_p = 7 \times 10^{-2}$$

If 0.50 mole of hydrogen and 1.0 mole of sulphur are heated to 87°C in 1.0 L vessel, what will be the partial pressure of H<sub>2</sub>S at equilibrium?

- (a) 0.966 atm
- (b) 1.38 atm
- (c) 0.0327 atm
- (d) 1 atm
- 72. Pure PCl<sub>5</sub> is introduced into an evacuated chamber and comes to equilibrium at 247°C and 2.0 atm. The equilibrium gaseous mixture contains 40% chlorine by volume.

Calculate  $K_p$  at 247°C for the reaction

$$PCl_5(g) \Leftrightarrow PCl_3(g) + Cl_2(g)$$

- (a) 0.625 atm
- (b) 4 atm
- (c) 1.6 atm
- (d) None of these

73. For the reaction

$$SnO_2(s) + 2H_2(g) \rightleftharpoons 2H_2O(g) + Sn(l)$$

calculate  $K_p$  at 900 K, where the equilibrium steam-hydrogen mixture was 45%  $H_2$  by volume :


- (a) 1.49
- (b) 1.22
- (c) 0.67
- (d) None of these


CHEM	CAL EQUILIBRIUM			247
74.	For the reaction $XCO_3(s)$	$\Rightarrow XO(s) + CO_{2}$	$(g), K_n = 1.642 \text{ atm a}$	t 727°C. If 4 moles of
	$XCO_3(s)$ was put into a 5			
	What mole percent of the			
		25	(c) 50	(d) None of these
<b>75.</b>	Fe <sub>2</sub> O <sub>3</sub> (s) may be converted	ed to Fe by the reac	tion	
	F	$e_2O_3(s) + 3H_2(g)$	$\Rightarrow$ 2Fe(s) + 3H <sub>2</sub> O(g),	
	for which $K_c = 8$ at temp.	720°C.		;•
	What percentage of the H	${\bf I_2}$ remains unreacte		
	(a) $\approx 22\%$ (b)	<b>≃</b> 34%	(c) $\approx 66\%$	$(d) \approx 78\%$
			1.	. `,-
76.	$AB_3(g)$ is dissociates as A	$B_3(g) \iff AB_2(g)$	$+\frac{1}{2}B_2(g)$ .	
	When the initial pressure	of AB <sub>2</sub> is 800 torr a	nd the total pressure dev	veloped at equilibrium is
	900 torr. What fraction of			
	(a) 10% (b)	20%	(c) 25%	(d) 30%
77.	At 1000 K, a sample of pr	_		
		<del>-</del> -	$2NO(g) + O_2(g)$	
	The equilibrium constant	$K_p$ is 156.25 atm. $I$	Analysis shows that the	partial pressure of $O_2$ is
	0.25 atm at equilibrium.			
=0	(a) 0.01 (b) Pure nitrosyl chloride (NOC	0.02	(c) 0.04	(d) None of these
78.	pressure was 1.0 atm and	the NOCl pressure	was 0.64 atm. What wo	ald be the value of $K_n$ ?
	(a) 1.02 atm		(b) $16.875 \times 10^{-3}$ atm	1
	(c) $16 \times 10^{-2}$ atm		(d) None of these	
79.	At a certain temperature	the equilibrium cor	istant $K_c$ is 0.25 for the	reaction
			$\leftarrow C_2(g) + D_2(g)$	
	If we take 1 mole of each	of the four gases in	a 10 litre container, wh	at would be equilibrium
	concentration of $A_2(g)$ ?	_		
	• •	) 0.033 M	(c) 0.133 M	(d) 1.33 <i>M</i>
80.	At 200°C PCl <sub>5</sub> dissociates		P.O. ( )	
		3 - 3 -	$PCl_3(g) + Cl_2(g)$	·
	It was found that the equ		e 62 times as heavy as l	hydrogen. The degree of
	dissociation of PCl <sub>5</sub> at 20 (a) 10% (b)	) 42%	(c) 50%	(d) 68%
R1	For the dissociation reacti	•	* *	•
<b>U</b> 1.	of $K_p$ and total equilibrium			
	$\sqrt{4P + K_n}$	$K_n$	$\sqrt{K_{\rm p}}$	215 ST 6.1
	(a) $\alpha = \sqrt{\frac{4P + K_p}{K_p}}$ (b)	$\alpha = \sqrt{\frac{\Delta P}{4P + K_{\rm p}}}$	(c) $\alpha = \sqrt{\frac{AP}{4P}}$	(d) None of these
	y P	, F		


82. The graph which represents all the equilibrium concentrations for the reaction


$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Then the select the correct graph for concentrations of NO2 against concentrations of NO4:









**83.** The vapour pressure of mercury is 0.002 mm Hg at 27°C.  $K_c$  for the process Hg(l)  $\iff$  Hg(g) is:

- (a) 0.002°
- (b)  $8.12 \times 10^{-5}$
- (c)  $6.48 \times 10^{-5}$
- (d)  $1.068 \times 10^{-7}$

**84.** Calculate the equilibrium constant  $(K_c)$  for the reaction below if they are present at equilibrium 5.0 mole of  $A_2$ , 3 mole of  $B_2$  and 2 mole of  $AB_2$  at 8.21 atm and 300 K

$$A_2(g) + 2B_2(g) \rightleftharpoons 2AB_2(g) + \text{Heat}$$

- (a) 1.333
- (b) 2.66
- (c) 20

(d) None of these

**85.** For the reaction (1) and (2)

$$A(g) \iff B(g) + C(g) \qquad \dots (1)$$

$$X(g) \rightleftharpoons 2Y(g)$$
 ...(2)

Given,  $K_{P_1}: K_{P_2} = 9:1$ 

If the degree of dissociation of A(g) and X(g) be same then the total pressure at equilibrium (1) and (2) are in the ratio:

- (a) 3:1
- (b) 36:1
- (c) 1:1
- (d) 0.5:1

**86.** Given the following reaction at equilibrium,  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ . Some inert gas at constant pressure is added to the system. Predict which of the following facts will be affected?

- (a) More NH<sub>3</sub>(g) is produced
- (b) Less NH<sub>3</sub>(g) is produced
- (c) No affect on the equilibrium
- (d)  $K_p$  of the reaction is decreased

**87.** Change in volume of the system does not alter the number of moles in which of the following equilibrium:

- (a)  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$
- (b)  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$
- (c)  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
- (d)  $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$

88. For the reaction

$$N_2(g) + 3H_2(g) \implies 2NH_3(g); \quad \Delta H = -93.6 \text{ kJ mol}^{-1}$$

the number of moles of H2 at equilibrium will increase if :

(a) volume is increased

- (b) volume is decreased
- (c) argon gas is added at constant volume
- (d) NH<sub>3</sub> is removed

89. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction

$$SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$$

When equilibrium is re-established:

- (a) the amount of  $Cl_2(g)$  remains unchanged
- (b) the amount of  $Cl_2(g)$  increases
- (c) the amount of SO<sub>2</sub>Cl<sub>2</sub>(g) increases
- (d) the amount of SO<sub>2</sub>(g) decreases
- 90. Some inert gas is added at constant volume to the following reaction at equilibrium

$$NH_4HS(s) \iff NH_3(g) + H_2S(g)$$


Predict the effect of adding the inert gas:

- (a) the equilibrium shifts in the forward direction
- (b) the equilibrium shifts in the backward direction
- (c) the equilibrium remains unaffected
- (d) the value of  $K_p$  is increased
- **91** Consider the reaction where  $K_p = 0.497$  at 500 K


$$PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$$


If the three gases are mixed in a rigid container so that the partial pressure of each gas in initially 1 atm. Which is true?


- (a) More PCl<sub>5</sub> will be produced
- (b) More PCl<sub>3</sub> will be produced
- (c) Equilibrium will be established when 50% reaction is complete
- (d) None of the above
- $SO_3(g)$ **92.** The bv reaction preparation  $SO_2(g) + \frac{1}{2}O_2(g) \implies SO_3(g)$  is an exothermic reaction. If the following follows the temperature-pressure preparation relationship for its % yield, then for temperatures  $T_1$ ,  $T_2$  and  $T_3$ .





- (a)  $T_3 > T_2 > T_1$
- (b)  $T_1 > T_2 > T_3$
- (c)  $T_1 = T_2 = T_3$
- (d) Nothing could be predicted about temperature through given information














- 94. In a vessel containing N2, H2 and NH3 at equilibrium, some helium gas is introduced so that total pressure increase while temperature and volume remain constant. According to Le Chatelier's principle, the dissociation of NH<sub>3</sub>:
  - (a) increases

(b) decreases

(c) remains unaltered

- (d) changes unpredictably
- 95. Le-Chatelier principle is not applicable to:
  - (a)  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$

- (b)  $Fe(s) + S(s) \Longrightarrow FeS(s)$
- (c)  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
- (d)  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$
- 96. Consider the following reactions. In which cases is the product formation favoured by decreased pressure?
  - $(1) CO_2(g) + C(s) \iff 2CO(g);$

$$\Delta H^{\circ} = +172.5 \text{ kJ}$$

(2) 
$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g); \Delta H^\circ = -91.8 \text{ kJ}$$

$$\Delta H^{\circ} = -91.8 \text{ k}$$

(3) 
$$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$$
;

$$\Delta H^{\circ} = 181 \text{ kJ}$$

(4) 
$$2H_2O(g) \rightleftharpoons 2H_2(g) + O_2(g); \Delta H^\circ = 484.6 \text{ kJ}$$

$$\Delta H^{\circ} = 484.6 \text{ kJ}$$

(b) 3, 4

(c) 2, 4

(d) 1, 4

- 97. Consider the following reactions. In which cases is product formation favoured by decreased temperature?
  - (1)  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ ;

$$\Delta H^{\circ} = 181 \text{ kJ}$$

(2) 
$$2CO_2(g) \iff 2CO(g) + O_2(g)$$
;

$$\Delta H^{\circ} = 566 \text{ kJ}$$

(3) 
$$H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$$
;

$$\Delta H^{\circ} = -9.4 \text{ kJ}$$

(4) 
$$H_2(g) + F_2(g) \rightleftharpoons 2HF(g)$$
;

$$\Delta H^{\circ} = -541 \text{ kJ}$$

(d) 3, 4

- **98.** For which of the following reactions is product formation favoured by low pressure and high temperature?
  - (a)  $H_2(g) + I_2(g) \implies 2HI(g)$ ;  $\Delta H^\circ = -9.4 \text{ kJ}$
  - (b)  $CO_2(g) + C(s) \iff 2CO(g); \qquad \Delta H^\circ = 172.5 \text{ kJ}$
  - (c)  $CO(g) + 2H_2(g) \rightleftharpoons CH_3OH$ ;  $\Delta H^{\circ} = -21.7 \text{ kJ}$
  - (d)  $3O_2(g) \iff 2O_3(g); \qquad \Delta H^\circ = 285 \text{ kJ}$
- **99.** For which of the following reaction is product formation favoured by low pressure and low temperature?
  - (a)  $CO_2(g) + C(s) \implies 2CO(g)$ ;  $\Delta H^\circ = 172.5 \text{ kJ}$
  - (b)  $CO(g) + 2H_2(g) \rightleftharpoons CH_3OH; \Delta H^\circ = -21.7 \text{ kJ}$
  - (c)  $2O_3(g) \implies 3O_2(g)$ ;  $\Delta H^\circ = -285 \text{ kJ}$
  - (d)  $H_2(g) + F_2(g) \rightleftharpoons 2HF(g);$   $\Delta H^{\circ} = -541 \text{ kJ}$
- **100.** Consider the following reactions at equilibrium and determine which of the indicated changes will cause the reaction to proceed to the right.
  - (1)  $CO(g) + 3H_2(g) \iff CH_4(g) + H_2O(g)$  (add  $CH_4$ )
  - (2)  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$  (remove  $NH_3$ )
  - $(3) H<sub>2</sub>(g) + F<sub>2</sub>(g) \implies 2HF(g) \qquad (add F<sub>2</sub>)$
  - (4)  $BaO(s) + SO_3(g) \implies BaSO_4(s)$  (add BaO)
  - (a) 2, 3 (b) 1, 4 (c) 2, 4 (d) 2, 3, 4
- 101. If the pressure in a reaction vessel for the following reaction is increased by decreasing the volume, what will happen to the concentrations of CO and CO<sub>2</sub>?

$$H_2O(g) + CO(g) \iff H_2(g) + CO_2(g) + Heat$$

- (a) both the [CO] and [CO2] will decrease
- (b) neither the [CO] nor the [CO2] will change
- (c) the [CO] will decrease and the [CO2] will increase
- (d) both the [CO] and  $[CO_2]$  will increase
- **102.** Consider the following reaction and determine which of the conditions will shift the equilibrium position to the right?

$$4NH_3(g) + 5O_2(g) \implies 4NO(g) + 6H_2O(g) + Heat$$

- (a) increasing the temperature
- (b) increasing the pressure

(c) adding a catalyst

- (d) none of the above is correct
- **103.** The conversion of ozone into oxygen is exothermic. Under what conditions is ozone the most stable?

$$2O_3(g) \rightleftharpoons 3O_2(g)$$

- (a) At low pressure and low temperature
- (b) At high pressure and high temperature
- (c) At high pressure and low temperature
- (d) At low pressure and high temperature
- **104.** A system at equilibrium is described by the equation of fixed temperature T.

$$SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$$

What effect will an increases in the total pressure caused by a decrease in volume have on the equilibrium?


- (a) Concentration of SO<sub>2</sub>Cl<sub>2</sub>(g) increases
- (b) Concentration of SO<sub>2</sub>(g) increases
- (c) Concentration of Cl<sub>2</sub>(g) increases
- (d) Concentration of all gases increases
- **105.** The reaction  $2NO_2(g) \rightleftharpoons N_2O_4(g)$  is an exothermic equilibrium. This means that :
  - (a) equilibration of this gas mixture will be slower at high temperature
  - (b) a mole of N<sub>2</sub>O<sub>4</sub> will occupy twice the volume of a mole of NO<sub>2</sub> at the same
  - (c) the equilibrium will move to the right if an equilibrium mixture is cooled
  - (d) the position of equilibrium will move to the left with increasing gas pressure
- 106. Densities of diamond and graphite are 3.5 and 2.3 gm/mL


C (diamond)  $\rightleftharpoons$  C (graphite);  $\Delta_r H = -1.9 \text{ kJ/mol}$ 


favourable conditions for formation of graphite are:


- (a) high pressure and low temperature(c) high pressure and high temperature
- (b) low pressure and high temperature
- (c) high pressure and high temperature (d) low pressure and low temperature **107.** For an equilibrium  $H_2O(s) \rightleftharpoons H_2O(l)$ , which of the following statements is true?
  - (a) The pressure changes do not affect the equilibrium
  - (b) More of ice melts if pressure on the system is increased
  - (c) More of liquid freezes if pressure on the system is increased
  - (d) The pressure changes may increase or decrease the degree of advancement of the process
- 108. A pressure cooker reduces cooking time for food because:
  - (a) the higher pressure inside the cooker crushes the food material
  - (b) cooking involves chemical changes helped by a rise in temperature
  - (c) heat is more evenly distributed in the cooking space
  - (d) boiling point of water involved in cooking is increased
- 109. The vapour pressure of a liquid in a closed container depends on :
  - (1) temperature of liquid
  - (2) quantity of liquid
  - (3) surface area of the liquid
  - (a) 1 only
- (b) 2 only
- (c) 1 and 3 only
- (d) 1, 2 and 3
- **110.** The pressure on a sample of water at its triple point is reduced while the temperature is held constant. Which phases changes are favoured?
  - (I) melting of ice
  - (II) sublimation of ice
  - (III) vaporization of liquid water
  - (a) I only
- (b) III only
- (c) II only
- (d) II and III

111. An exothermic reaction is represented by the graph:











112. An endothermic reaction is represented by the graph:



113. A schematic plot of  $\ln K_{eq}$  versus inverse of temperature for a reaction is shown below



the reaction must be:

- (a) Exothermic
- (b) Endothermic
- (c) One with negligible enthalpy change
- (d) Highly spontaneous at ordinary temperature

**114.** What is the correct relationship between free energy change and equilibrium constant of a reaction:


(a) 
$$\Delta G^{\circ} = RT \ln K$$


(b) 
$$\Delta G^{\circ} = -RT \ln K$$


(c) 
$$\Delta G = RT \ln K$$


(d) 
$$\Delta G = -RT \ln K$$

**115.** For the chemical equilibrium  $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ ,  $\Delta H_r^0$  can be determined from which one of the following plots ?









116.  $K_p$  has the value of  $10^{-6}$  atm<sup>3</sup> and  $10^{-4}$  atm<sup>3</sup> at 298 K and 323 K respectively for the reaction

. 
$$CuSO_4 \cdot 3H_2O(s) \iff CuSO_4(s) + 3H_2O(g)$$

 $\Delta_r H^{\circ}$  for the reaction is :

(a). 7.7 kJ/mol

(b) -147.41 kJ/mol

(c) 147.41 kJ/mol

(d). None of these

**117.** van't Hoffs equation shows the effect of temperature on equilibrium constants  $K_c$  and  $K_p$ . The  $K_p$  varies with temperature according to the relation :

(a) 
$$\log \frac{K_{p_2}}{K_{p_1}} = \frac{\Delta H^{\circ}}{2.303 R} \left( \frac{T_1 - T_2}{T_1 T_2} \right)$$

(b) 
$$\log \frac{K_{p_2}}{K_{p_1}} = \frac{\Delta H^{\circ}}{2.303 R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$$

(c) 
$$\log \frac{K_{p_2}}{K_{p_1}} = \frac{\Delta E^{\circ}}{2.303 R} \left( \frac{T_1 - T_2}{T_1 T_2} \right)$$

(d) None of these

- **118.** For a reaction, the value of  $K_p$  increases with increase in temperature. The  $\Delta H$  for the reaction would be:
  - (a) positive

(b) negative

(c) zero

- (d) cannot be predicted
- 119. The most stable oxides of nitrogen will be:

(a)  $2NO_2(g) \rightleftharpoons N_2(g) + 2O_2(g)$ ;  $K = 6.7 \times 10^{16} \text{ mol L}^{-1}$ 

(b)  $2N_2O_5(g) \rightleftharpoons 2N_2(g) + 5O_2(g)$ ;  $K = 1.2 \times 10^{24} \text{ mol}^5 \text{ L}^{-5}$ 

(c)  $2NO(g) \rightleftharpoons N_2(g) + O_2(g)$ ;

 $K = 2.2 \times 10^{30}$ 

(d)  $2N_2O(g) \iff 2N_2(g) + O_2(g); K = 3.5 \times 10^{33} \text{ mol } L^{-1}$ 

120. When 1 mole of pure ethyl alcohol (C<sub>2</sub>H<sub>5</sub>OH) is mixed with 1 mole of acetic acid at 25°C, the equilibrium mixture contains 2/3 mole each of ester and water

$$C_2H_5OH(l) + CH_3COOH(l) \Leftrightarrow CH_3COOC_2H_5(l) + H_2O$$

The  $\Delta G^{\circ}$  for the reaction at 298 K is:

(a) 3435 J

(b) 4 J

(c) -3435 J

**121.** The value of  $\Delta G^{\circ}$  for a reaction in aqueous phase having  $K_c = 1$ , would be:

(a) -RT

(b) -1

(d) +RT

- 122. A plot of Gibbs energy of a reaction mixture against the extent of the reaction is :
  - (a) minimum at equilibrium

(b) zero at equilibrium

(c) maximum at equilibrium

(d) None of these

123. For the reaction at 300 K

$$A(g) \iff V(g) + S(g)$$

$$\Delta_r H^{\circ} = -30 \text{ kJ/mol}, \ \Delta_r S^{\circ} = -0.1 \text{ kJ. K}^{-1}. \text{ mol}^{-1}$$

What is the value of equilibrium constant?

(a) 0

(c) 10

(d) None of these

124. Solid Ca(HCO<sub>3</sub>)<sub>2</sub> decomposes as

$$Ca(HCO_3)_2(s) \rightleftharpoons CaCO_3(s) + CO_2(g) + H_2O(g)$$

If the total pressure is 0.2 bar at 420 K, what is the standard free energy change for the given reaction  $(\Delta, G^{\circ})$ ?

(a) 840 kJ/mol

(b) 3.86 kJ/mol

(c) 6.98 kJ/mol

(d) 16.083 kJ/mol

- 125. The standard free energy change of a reaction is  $\Delta G^{\circ} = -115$  kJ at 298 K. Calculate the value of  $\log_{10} K_p$  (R = 8.314 JK<sup>-1</sup> mol<sup>-1</sup>)
  - (a) 20.16
- (b) 2.303
- (c) 2.016
- (d) 13.83

1. The following equilibrium constants were determined at 1120 K:

$$2CO(g) \rightleftharpoons C(s) + CO_2(g);$$

$$K_{p_1} = 10^{-14} \text{ atm}^{-1}$$

$$CO(g) + Cl_2(g) \iff COCl_2(g);$$

$$K_{p_2} = 6 \times 10^{-3} \text{ atm}^{-1}$$

What is the equilibrium constant  $K_c$  for the following reaction at 1120 K:

$$C(s) + CO_2(g) + 2Cl_2(g) \rightleftharpoons 2COCl_2(g)$$

(a) 
$$3.31 \times 10^{11} M^{-1}$$

(b) 
$$5.5 \times 10^{10} M^{-1}$$
 (c)  $5.51 \times 10^6 M^{-1}$ 

(c) 
$$5.51 \times 10^6 M^{-1}$$

(d) None of these

2. One mole of  $N_2(g)$  is mixed with 2 moles of  $H_2(g)$  in a 4 litre vessel. If 50% of  $N_2(g)$  is converted to  $NH_3(g)$  by the following reaction:


$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

What will be the value of  $K_c$  for the following equilibrium?

$$NH_3(g) \iff \frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$$

(c) 
$$\frac{1}{16}$$

(d) None of these



The gas  $A_2$  in the left flask allowed to react with gas  $B_2$  present in right flask as  $A_2(g) + B_2(g) \rightleftharpoons 2AB(g)$ ;  $K_c = 4$  at 27°C. What is the concentration of AB when equilibrium is established?

Assume that the decomposition of HNO<sub>3</sub> can be represented by the following equation

$$4HNO_3(g) \iff 4NO_2(g) + 2H_2O(g) + O_2(g)$$

and the reaction approaches equilibrium at 400 K temperature and 30 atm pressure. At equilibrium partial pressure of HNO3 is 2 atm.

Calculate  $K_c$  in  $(\text{mol/L})^3$  at 400 K:

(Use: R = 0.08 atm-L/mol-K)

5. For the equilibrium

$$LiCl \cdot 3NH_3(s) \iff LiCl \cdot NH_3(s) + 2NH_3(g);$$

 $K_p = 9$  atm<sup>2</sup> at 37°C. A 5 litre vessel contains 0.1 mole of LiCl·NH<sub>3</sub>. How many moles of NH<sub>3</sub> should be added to the flask at this temperature to derive the backward reaction for completion?

Use: R = 0.082 atm-L/mol K

6.	Ammonium	carbamate	dissociates	as

$$NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$$

In a closed vessel containing ammonium carbamate in equilibrium, ammonia is added such that partial pressure of  $\mathrm{NH}_3$  now equals to the original total pressure. Calculate the ratio of partial pressure of  $\mathrm{CO}_2$  now to the original partial pressure of  $\mathrm{CO}_2$ :

(a) 4

(b) 9

(c)  $\frac{4}{9}$ 

(d)  $\frac{2}{9}$ 

7. For the reaction  $C_2H_6(g) \rightleftharpoons C_2H_4(g) + H_2(g)$ 

 $K_p$  is  $5 \times 10^{-2}$  atm. Calculate the mole per cent of  $C_2H_6(g)$  at equilibrium if pure  $C_2H_6$  at 1 atm is passed over a suitable catalyst at 900 K:

(a) 20

- (b) 33.33
- (c) 66.66
- (d) None of these
- **8.**  $2NOBr(g) \rightleftharpoons 2NO(g) + Br_2(g)$ . If nitrosyl bromide (NOBr) is 40% dissociated at certain temp. and a total pressure of 0.30 atm.  $K_p$  for the reaction  $2NO(g) + Br_2(g) \rightleftharpoons 2NOBr(g)$  is:
  - (a) 45
- (b) 25

- (c) 0.022
- (d) 0.25

**9.** Consider the partial decomposition of A as

$$2A(g) \iff 2B(g) + C(g)$$

At equilibrium 700 mL gaseous mixture contains 100 mL of gas C at 10 atm and 300 K. What is the value of  $K_p$  for the reaction?

(a)  $\frac{40}{7}$ 

- (b)  $\frac{1}{28}$
- (c)  $\frac{10}{28}$
- (d)  $\frac{28}{10}$
- **10.** At a certain temperature and 2 atm pressure equilibrium constant  $(K_p)$  is 25 for the reaction  $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$

Initially if we take 2 moles of each of the four gases and 2 moles of inert gas, what would be the equilibrium partial pressure of NO<sub>2</sub>?

- (a) 1.33 atm
- (b) 0.1665 atm
- (c) 0.133 atm
- (d) None of these
- 11. 0.020 g of selenium vapour at equilibrium occupying a volume of 2.463 mL at 1 atm and 27°C. The selenium is in a state of equilibrium according to reaction

$$3Se_2(g) \iff Se_6(g)$$

What is the degree of association of selenium?

(At. wt. of Se = 79)

- (a) 0.205
- (b) 0.315
- (c) 0.14
- (d) None of these
- 12. Determine the degree of association (polymerization) for the reaction in aqueous solution  $\Theta \leftarrow C_6 H_{12} O_6$

If observed (mean) molar mass of HCHO and  $C_6H_{12}O_6$  is 150 :

- (a) 0.50
- (b) 0.833
- (c) 0.90
- (d) 0.96
- 13. A reaction system in equilibrium according to reaction  $2SO_2(g) + O_2(g) \implies 2SO_3(g)$  in one litre vessel at a given temperature was found to be 0.12 mole each of  $SO_2$  and  $SO_3$  and 5 mole of  $O_2$ . In another vessel of one litre contains 32 g of  $SO_2$  at the same temperature. What mass of  $O_2$  must be added to this vessel in order that at equilibrium 20% of  $SO_2$  is oxidized to  $SO_3$ ?
  - (a) -0.4125 g
- (b) 11.6 g
- (c) 1.6 g
- (d) None of these

**14.** The equilibrium constant  $K_p$  for the reaction

$$N_2O_4(g) \iff 2NO_2(g) \text{ is } 4.5$$

What would be the average molar mass (in g/mol) of an equilibrium mixture of  $N_2O_4$  and  $NO_2$  formed by the dissociation of pure  $N_2O_4$  at a total pressure of 2 atm?

(a) 69

- (b) 57.5
- (c) 80.5 ·
- (d) 85.5
- **15.** A flask containing 0.5 atm pressure of  $A_2(g)$ , some solid AB added into flask which undergoes dissociation according to  $2AB(s) \iff A_2(g) + B_2(g)$   $K_p = 0.06$  atm<sup>2</sup>

The total pressure (in atm) at equilibrium is:

- (a) 0.70
- (b) 0.6
- (c) 0.10
- (d) None of these
- **16.** A vessel of 250 litre was filled with 0.01 mole of Sb<sub>2</sub>S<sub>3</sub> and 0.01 mole of H<sub>2</sub> to attain the equilibrium at 440°C as

$$Sb_2S_3(s) + 3H_2(g) \rightleftharpoons 2Sb(s) + 3H_2S(g)$$

After equilibrium, the  $H_2S$  formed was analysed by dissolved it in water and treating with excess of  $Pb^{2+}$  to give 1.19 g of PbS as precipitate. What is the value of  $K_c$  at 440°C?

(a) :

(b) 2

(c) 4

- (d) 8
- 17. For the reaction  $2A(g) + B(g) \iff C(g) + D(g); K_c = 10^{12}$ . If the initial moles of A, B, C and D are 2, 1, 7 and 3 moles respectively in a one litre vessel. What is the equilibrium concentration of A?
  - (a)  $4 \times 10^{-4}$
- (b)  $2 \times 10^{-4}$
- (c)  $10^{-4}$
- (d)  $8 \times 10^{-4}$
- 18. The equilibrium constant for the reaction in aqueous solution

$$H_3BO_3 + glycerin \Leftrightarrow (H_3BO_3 - glycerin)$$
 is 0.90.

How many moles of glycerin should be added per litre of  $0.10 M H_3 BO_3$  so that 80% of the  $H_3 BO_3$  is converted to the boric-acid-glycerin complex?

- (a) 4.44
- (b) 4.52
- (c) 3.6

- (d) 0.08
- 19. Rate of diffusion of ozonized oxygen is  $0.4\sqrt{5}$  times that of pure oxygen. What is the per cent degree of association of oxygen assuming pure  $O_2$  in the sample initially?
  - (a) 20

(b) 40

(c) 60

- (d) None of these
- 20. One mole of  $SO_3$  was placed in a two litre vessel at a certain temperature. The following equilibrium was established in the vessel

$$2SO_3(g) \rightleftharpoons 2SO_2(g) + O_2(g)$$

the equilibrium mixture reacted with 0.2 mole KMnO<sub>4</sub> in acidic medium. Hence,  $K_c$  is:

(a) 0.50

(b) 0.25

(c) 0.125

- (d) None of these
- 21. At 800°C, the following equilibrium is established as

$$F_2(g) \iff 2F(g)$$

The composition of equilibrium may be determined by measuring the rate of effusion of the mixture through a pin hole. It is found that at 800°C and 1 atm mixture effuses 1.6 times as fast as  $SO_2$  effuses under the similar conditions. (At. wt. of F = 19). What is the value of  $K_p$  (in atm)?

(a) 0.315

(b) 0.685

(c) 0.46

(d) 1.49

22. The equilibrium constant for the ionization of  $RNH_2(g)$  in water as  $RNH_2(g) + H_2O(l) \rightleftharpoons RNH_3^+(aq) + OH^-(aq)$ 

is  $8 \times 10^{-6}$  at 25°C. Find the pH of a solution at equilibrium when pressure of RNH<sub>2</sub>(g) is 0.5 bar :

- (a)  $\approx 12.3$
- (b)  $\approx 11.3$
- (c)  $\approx 11.45$
- (d) None

**23.** Calculate  $\triangle$ , G for the reaction at 27°C

$$H_2(g) + 2Ag^+(aq) \iff 2Ag(s) + 2H^+(aq)$$

Given:  $P_{\text{H}_2} = 0.5 \text{ bar}$ ;  $[\text{Ag}^+] = 10^{-5} M$ ;

 $[H^+] = 10^{-3} M$ ;  $\Delta_r G^{\circ} [Ag^+(aq)] = 77.1 \text{ kJ/mol}$ 

(a) -154.2 kJ/mol

(b) -178.9 kJ/mol

(c) -129.5 kJ/mol

- (d) None of these
- **24.** When  $N_2O_5$  is heated at certain temperature, it dissociates as  $N_2O_5(g) \rightleftharpoons N_2O_3(g) + O_2(g)$ ;  $K_c = 2.5$ . At the same time  $N_2O_3$  also decomposes as :  $N_2O_3(g) \rightleftharpoons N_2O(g) + O_2(g)$ . If initially 4.0 moles of  $N_2O_5$  are taken in 1.0 litre flask and allowed to dissociation, concentration of  $O_2$  at equilibrium is 2.5 M. Equilibrium concentration of  $N_2O_5$  is :
  - (a) 1.0 M
- (b) 1.5 M
- (c) 2.166 M
- (d) 1.846 M

25. Two solid compounds X and Y dissociates at a certain temperature as follows

$$X(s) \iff A(g) + 2B(g); K_{p_1} = 9 \times 10^{-3} \text{ atm}^3$$

$$Y(s) \iff 2B(g) + C(g); K_{p_2} = 4.5 \times 10^{-3} \text{ atm}^3$$

The total pressure of gases over a mixture of X and Y is:

- (a) 4.5 atm
- (b) 0.45 atm
- (c) 0.6 atm
- (d) None of these

For a gaseous reaction

$$aA(g) + bB(g) \iff cC(g) + dD(g)$$

equilibrium constant 
$$K_c$$
,  $K_p$  and  $K_x$  are represented by the following relations 
$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}; \quad K_p = \frac{p_C^c \cdot p_D^d}{p_A^a \cdot p_B^b} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where [A] represents molar concentration of A,  $p_A$  represents partial pressure of A and Prepresents total pressure,  $x_A$  represents mole fraction of A ٠:٠

1. On the basis of above work-up. Select write option

(a) 
$$K_p = K_c (RT)^{\Delta n_g}$$
;  $K_x = K_p (RT)^{\Delta n_g}$ 

(b) 
$$K_c = K_p (RT)^{-\Delta n_g}$$
;  $K_p = K_x P^{\Delta n_g}$ 

(c) 
$$K_c = K_x P^{\Delta n_g}$$
;  $K_p = K_x P^{\Delta n_g}$ 

(d) 
$$K_c = K_p (RT)^{-\Delta n_g}$$
;  $K_x = K_p (RT)^{\Delta n_g}$ 

2. For the reaction  $SO_2Cl_2(g) \iff SO_2(g) + Cl_2(g), K_p > K_x$  is obtained at :

- (a) 0.5 atm
- (b) 0.8 atm
- (c) 1 atm
- (d) 2 atm

3. For the following equilibrium relation between  $K_c$  and  $K_x$  (in terms of mole fraction) is  $PCl_3(g) + Cl_2(g) \iff PCl_5(g)$ 

(a) 
$$K = K (RT)^{-1}$$
 (b)  $K$ 

(b) 
$$K_c = K_x(RT)$$

(c) 
$$K_c = K_x \left(\frac{RT}{P}\right)$$

(a) 
$$K_c = K_x (RT)^{-1}$$
 (b)  $K_c = K_x (RT)$  (c)  $K_c = K_x \left(\frac{RT}{P}\right)$  (d)  $K_p = K_x \left(\frac{P}{RT}\right)$ 

# PASSAGE

Variation of equilibrium constant K with temperature is given by van't Hoff equation  $\ln K = \frac{\Delta S_r^{\circ}}{R} - \frac{\Delta H_r^{\circ}}{RT}$ 

$$\ln K = \frac{\Delta \bar{S}_r^{\circ}}{R} - \frac{\Delta H_r^{\circ}}{RT}$$

from this equation,  $\Delta H_r^{\circ}$  can be evaluated if equilibrium constants  $K_1$  and  $K_2$  at two temperature  $T_1$  and  $T_2$  are known.

$$\log\left(\frac{K_2}{K_1}\right) = \frac{\Delta H_r^{\circ}}{2.303 R} \left[\frac{1}{T_1} - \frac{1}{T_2}\right]$$

1. For an isomerization  $X(g) \rightleftharpoons Y(g)$ , the temperature dependency of equilibrium constant is given by:

$$\ln K = 2 - \frac{1000}{T}$$

The value of  $\Delta_r S^\circ$  at 300 K is:

(a) 2R

- (c) 1000 R
- (d) None of these

2. Select the correct statement:

(a) Value of Kee always increases with increasing temperature


(b) For exothermic reaction value of  $K_{eq}$  increases with decreasing in temperature

(c) For endothermic reaction value of  $K_{eq}$  increases with decreasing in temperature

(d) For exothermic reaction slope is (log  $K \vee s \cdot 1/T$ ) negative

3. Variation of  $\log_{10} K$  with  $\frac{1}{r}$  is shown by the following graph in which straight line is at 45°C,

hence  $\Delta H^{\circ}$  is:



(a) -4.606 kJ/mol

(b) -19.147 kJ/mol (c) 8.314 kJ/mol (d) 10 kJ/mol

The equilibrium constant  $K_p$  for the reaction

$$A(g) \iff B(g) + C(g)$$

is 1 at 27°C and 4 at 47°C. For the reaction calculate enthalpy change for the  $B(g) + C(g) \rightleftharpoons A(g)$  (Given : R = 2 cal/mol-K)

(a) - 13.31 kcal/mol (b) 13.31 kcal/mol

(c) - 19.2 kcal/mol

(d) 55.63 kcal/mol

 $N_2O_3$  is an unstable oxide of nitrogen and it docomposes into NO(g) and  $NO_2(g)$  where NO2(g) is further dimerise into N2O4 as

(i)  $N_2O_3(g) \rightleftharpoons NO_2(g) + NO(g)$ 

 $K_{p_1} = 2.5 \text{ bar}$ 

(ii)  $2NO_2(g) \rightleftharpoons N_2O_4(g)$ 

 $K_{o_2}$ 

A flask is initially filled with pure N<sub>2</sub>O<sub>3</sub>(g) having pressure 2 bar and equilibria were established.

At equilibrium partial pressure of NO(g) was found to be 1.5 bar.

**1.** The equilibrium partial pressure of  $N_2O_3(g)$  is:

(a) 0.5 bar

(b) 1.0 bar

(c) 1.5 bar

(d) 0.1 bar

2. The equilibrium partial pressure of NO<sub>2</sub>(g) is:

(a) 0.066 bar

(b) 0.133 bar

(c) 0.423 bar

(d) 0.83 bar

**3.** The value of  $K_{p_2}$  is: (a) 0.16 bar<sup>-1</sup>

(b)  $0.32 \text{ bar}^{-1}$ 

(c)  $0.48 \text{ bar}^{-1}$ 

(d) 0.64 bar<sup>-1</sup>

### PASSAGE 4

If a system at equilibrium is subjected to a change of any one of the factors such as concentration, pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

#### Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration become 'Q' (reaction quotient) and depending upon.

Q < K: equilibrium will shift in forward direction

Q > K: equilibrium will shift in backward direction

#### Effect of change in pressure:

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is decrease in number of moles *i.e.*, towards the direction in which there is decrease in volume.

#### Effect of change in pressure on melting point: There are two types of solids:

(a) Solids whose volume decreases on melting, e.g., ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume) 

⇒ Liquid (lower volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

(b) Solid whose volume increase on melting, e.g., Fe, Cu, Ag, Au, etc.

Solid (lower volume) ⇌ Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

(c) **Solubility of substances**: When solid substance are dissolved in water, either heat is evolved.

For endothermic solubility process solubility increase with increase in temperature.

For exothermic solubility decrease with increase in temperature.

(d) **Solubility of gases in liquids :** When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

**Effect of temperature:** For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.

- 1. A 'X'(g) solute when dissolved in water heat is evolved. Then solubility of 'X' will increase:
  - (a) High temperature, low pressure
- (b) Low temperature, high pressure
- (c) High temperature, high pressure
- (d) Low temperature, low pressure

2.  $Fe(l) \rightleftharpoons Fe(s)$ 

Above equilibrium is favoured at:

- (a) High pressure, low temperature
- (b) High pressure, high temperature
- (c) Low pressure, high temperature (d) Low pressure, low temperature

#### 3. For the reaction

$$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$$

If pressure is increased by reducing the volume of the container then:

- (a) Total pressure at equilibrium will remain same
- (b) Concentration of all the component at equilibrium will change
- (c) Concentration of all the component at equilibrium will remain same
- (d) Equilibrium will shift in the backward direction

## ONE OR MORE ANSWERS IS/ARE CORRECT

- 1. A catalyst:
  - (a) increases the average kinetic energy of reacting molecules
  - (b) decreases the activation energy
  - (c) can alters the reaction mechanism
  - (d) can change pre-exponential factor
- 2. Which of the following is correct about the chemical equilibrium?
  - (a)  $(\Delta G)_{T,P} = 0$
  - (b) Equilibrium constant is independent of initial concentration of reactants
  - (c) Catalyst has no effect on equilibrium state
  - (d) Reaction stops at equilibrium
- 3. For the reaction

$$AB_2(g) \iff AB(g) + B(g)$$

If  $\alpha$  is negligiable w.r.t. 1 then degree of dissociation ( $\alpha$ ) of  $AB_2$  is proportional to : (a)  $\frac{1}{R}$  (b)  $\frac{1}{V}$  (c)  $\frac{1}{\sqrt{N}}$  (d)  $\sqrt{V}$ 

(a)  $\frac{1}{D}$ 

- 4. Consider the reactions given below. In which cases will the reaction proceed toward right by increasing the pressure?

(a) 
$$4HCl(g) + O_2(g) \longrightarrow 2Cl_2(g) + 2H_2O(g)$$

(b) 
$$\operatorname{Cl}_2(g) + \operatorname{H}_2\operatorname{O}(g) \longrightarrow 2\operatorname{HCl}(g) + \frac{1}{2}\operatorname{O}_2(g)$$

(c) 
$$CO_2(g) + 4H_2(g) \longrightarrow CH_4(g) + 2H_2O(g)$$

(d) 
$$N_2(g) + O_2(g) \longrightarrow 2NO(g)$$

5. Ammonia is a weak base that reacts with water according to the equation

$$NH_3(aq) + H_2O(l) \rightleftharpoons NH_4^+(aq) + OH^-(aq)$$

Select the correct option(s) that can increase the moles of ammonium ion in water:

(a) Addition of HCl

(b) Addition of NaOH

(c) Addition of NH<sub>4</sub>Cl

- (d) Addition of H<sub>2</sub>O
- **6.** Consider the reaction  $2CO(g) + O_2(g) \implies 2CO_2(g) + \text{Heat}$

Under what conditions shift is undeterminable?

- (a) Addition of O2 and decrease in volume
- (b) Addition of CO and removal of CO2 at constant volume

- (c) Increase in temperature and decrease in volume
- (d) Addition of CO and increase in temperature at constant volume
- 7. What will be the effect of addition of catalyst at constant temperature?
  - (a) The equilibrium constant will remain constant
  - (b)  $\Delta H$  of the reaction will remain constant
  - (c)  $k_t$  and  $k_h$  will increase upto same extent
  - (d) equilibrium composition will change
- **8.** For the reaction  $PCl_5(g) \iff PCl_3(g) + Cl_2(g)$ , the forward reaction at constant temperature is favoured by:
  - (a) introducing an inert gas at constant volume
  - (b) introducing chlorine gas at constant volume
  - (c) introducing an inert gas at constant pressure
  - (d) increasing the volume of the container
- 9. For the reaction :  $Cl_2(g) + 3F_2(g) \rightleftharpoons 2ClF_3(g)$ ;  $\Delta H = -329$  kJ, dissociation of  $ClF_3(g)$  will be favoured by :
  - (a) increasing the temperature
  - (b) increasing the volume of the container
  - (c) adding of F2 gas
  - (d) adding of inert gas at constant pressure
- 10. Increase in the pressure for the following equilibrium:  $H_2O(l) \rightleftharpoons H_2O(g)$ , result in the:
  - (a) formation of more H<sub>2</sub>O(l)
- (b) formation of more H<sub>2</sub>O(g)

(c) increase in b.p. of  $H_2O(l)$ 

- (d) decrease in b.p. of H<sub>2</sub>O(l)
- 11. Heating a II group metal carbonate leads to decomposition as :

$$BaCO_3(s) \rightleftharpoons BaO(s) + CO_2(g)$$

Equilibrium will shift left

(a) by addition of BaO(s)

- (b) by addition of CO<sub>2</sub>(g)
- (c) by decreasing the temperature
- (d) by decreasing the volume of the vessel
- 12.  $N_2(g)$  and  $H_2(g)$  are allowed to react in a closed vessel at given temp. and pressure for the formation of  $NH_3(g)[N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g) + 22.4 \text{ kcal}]$  if He(g) is added at equilibrium at constant pressure then which is/are correct?
  - (a) Concentration of N<sub>2</sub>(g), H<sub>2</sub>(g) and NH<sub>3</sub>(g) decrease.
  - (b) Moles of NH3(g) decreases.
  - (c) The extent of cooling depends on amount of He(g) added.
  - (d) Concentration of N2 and H2 increases and concentration of NH3 decreases.

#### MATCH THE COLUMN

**Column-I** and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

#### Column-1

- (A)  $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$
- (B)  $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$
- (C)  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$
- (D)  $HCl(g) \rightleftharpoons H^+(aq) + Cl^-(aq)$

Column-II

- (P)  $K_p > K_c$  above room temperature
- (Q)  $K_p = K_c$  above room temperature
- (R)  $K_p < K_c$  above room temperature
- (S)  $K_p$  and  $K_c$  not defined

2.

#### Column-)

- (A)  $3O_2(g) \rightleftharpoons 2O_3(g)$
- (B)  $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$
- (C)  $2HF(g) \rightleftharpoons H_2(g) + F_2(g)$
- (D)  $CO(g) + 3H_2(g) \rightleftharpoons CH_4(g) + H_2O(g)$

Column-II

- (P) no unit
- (Q)  $atm^{-1/2}$
- (R) atm<sup>-1</sup>
- (S) atm<sup>-2</sup>

3.

# Column-I (Reaction)

- (A)  $2X(g) \rightleftharpoons Y(g) + Z(g)$
- (B)  $X(g) \rightleftharpoons Y(g) + Z(g)$
- (C)  $3X(g) \rightleftharpoons Y(g) + Z(g)$
- (D)  $2X(g) \rightleftharpoons Y(g) + 2Z(g)$

Column-II (If  $\alpha$  is negligiable w.r.t. 1)

- (P)  $\alpha = 2 \times \sqrt{K_c}$
- (Q)  $\alpha = 3 \times \sqrt{K_c}$
- (R)  $\alpha = (2K_c)^{1/3}$
- (S)  $\alpha = \sqrt{K_c}$

4.

#### Column-I

- (A)  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g); \Delta H = -ve$
- (B)  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ ;  $\Delta H = + ve$
- (C)  $A(g) + B(g) \rightleftharpoons 2C(g) + D(g);$  $\Delta H = + \text{ ve}$
- (D)  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ ;  $\Delta H = + ve$

Column-II

- (P) K increases with increase in temperature
- (Q) K decreases with increase in temperature
- (R) Pressure has no effect
- (S) Product moles, increases due to addition of inert gas at constant pressure

#### 5.

#### Column-I

- (A)  $\frac{K_{10+T^{\circ}C}}{K_{T^{\circ}C}} = 2$
- (B)  $\frac{K_{10+T^{\circ}C}}{K_{T^{\circ}C}} = \frac{1}{2}$
- (C)  $A(g) + B(g) \rightleftharpoons C(g)$
- (D)  $X(s) + Y(g) \rightleftharpoons Z(g)$

#### Column-II

- (P) Endothermic
- (Q) Not affected by pressure
- (R) Exothermic
- (S) Affected by volume

#### 6.

#### Column-I

- (A) Pressure increased in  $2NO(g) \rightleftharpoons N_2(g) + O_2(g)$
- (B) Pressure increased in  $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$
- (C) Temp. increased and pressure increased  $3O_2(g) \rightleftharpoons 2O_3(g)$ ;  $\Delta H = 285 \text{ kJ}$
- (D) Pressure decrease and moles of  $N_2$  increased

$$N_2(g) + 2O_2(g) \rightleftharpoons 2NO_2(g);$$


 $\Delta H = 66.4 \text{ kJ}$ 

#### Column-II

- (P) Equilibrium shifted in forward direction
- (Q) Equilibrium shifted in backward direction
- (R) Equilibrium remains unaffected
- (S) Theoretically we cannot predict


### 7.


# Column-I (Reactions)




(B) 
$$I_2(s) \implies I_2(aq)$$

#### Column-II (Equilibrium States)






(C) 
$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$



(D) 
$$AgI(s) \rightleftharpoons Ag^+(aq) + I^-(aq)$$



#### **ASSERTION-REASON TYPE QUESTIONS**

Each question contains STATEMENT-1 (Assersion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE
- **1. STATEMENT-1:** The endothermic reactions are favoured at lower temperature and the exothermic reactions are favoured at higher temperature.
  - **STATEMENT-2:** When a system in equilibrium is disturbed by changing the temperature, it will tend to adjust itself so as to overcome the effect of change.
- 2. STATEMENT-1: The melting point of ice decreases with increase of pressure.
  - **STATEMENT-2:** Ice contracts on melting.
- **3. STATEMENT-1:** The equilibrium of  $A(g) \iff B(g) + C(g)$  is not affected by changing the volume.
  - **STATEMENT-2:**  $K_c$  for the reaction does not depend on volume of the container.
- **4. STATEMENT-1:** For a chemical reaction at initial stage rate of forward reaction  $(r_f)$  is greater than rate of reversed reaction  $(r_h)$ 
  - **STATEMENT-2:** When  $r_f = r_b$ , chemical reaction is at equilibrium.
- **5. STATEMENT-1:** For the reaction  $A(g) \rightleftharpoons B(g) + C(g)$ ,  $K_p = 1$  atm. If we start with equal moles of all gases at 9 atm of initial pressure, then at equilibrium partial pressure of A increases.
  - **STATEMENT-2:** Reaction quotient  $Q_p > K_p$  hence equilibrium shifts in backward direction.

- **6. STATEMENT-1:** The gas phase reaction  $PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$  shifts to the right on increasing pressure.
  - **STATEMENT-2:** When pressure increases, equilibrium shifts towards more number of moles.
- **7. STATEMENT-1:** For a reaction at equilibrium, the Gibb's free energy of reaction is minimum at constant temp. and pressure.
  - **STATEMENT-2:** The Gibb's free energy of both reactants and products increases and become equal at equilibrium.
- 8. STATEMENT-1: The physical equilibrium is not static but dynamic in nature.
  - **STATEMENT-2:** The physical equilibrium is a state in which two opposing process are proceeding at the same rate.
- **9. STATEMENT-1:** Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.
  - **STATEMENT-2:** Eqilibrium constant depends upon the way in which the reaction is written.
- **10. STATEMENT-1:** If  $Q_p < K_p$  reaction moves in direction of reactants.
  - **STATEMENT-2:** Reaction quotient is defined in the same way as equilibrium constant at any stage of the reaction.
- **11. STATEMENT-1:** For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$  if the volume of vessel is reduced to half of its original volume, equilibrium concentration of all gases will be doubled.
  - **STATEMENT-2:** According to Le-Chatelier's principle, reaction shifts in a direction that tends to minimized the effect of the stress.
- **12. STATEMENT-1:** The equilibrium constant of the exothermic reaction at high temperature decreases.
  - **STATEMENT-2:** Since  $\ln \frac{K_2}{K_1} = \frac{\Delta H^{\circ}}{R} \left[ \frac{1}{T_1} \frac{1}{T_2} \right]$  and for exothermic reaction,
    - $\Delta H^{\circ} = -\text{ ve and thereby; } \frac{K_2}{K_1} < 1$
- **13. STATEMENT-1:** For the reaction at certain temperature

$$A(g) + B(g) \iff C(g)$$

there will be no effect by addition of inert gas at constant volume.

- **STATEMENT-2:** Molar concentration of all gases remains constant.
- **14. STATEMENT-1:** For the physical equilibrium  $H_2O(s) \rightleftharpoons H_2O(l)$  on increasing temperature and increasing pressure more water will form.
  - **STATEMENT-2:** Since forward reaction is endothermic in nature and volume of water is greater than that of the volume of ice.
- **15. STATEMENT-1:** The catalyst does not alter the equilibrium constant.
  - **STATEMENT-2:** Because for the catalysed reaction and uncatalysed reaction  $\Delta H$  remains same and equilibrium constant depends on  $\Delta H$ .



### SUBJECTIVE PROBLEM

1. If 50% of CO<sub>2</sub> converts to CO at the following equilibrium:

$$\frac{1}{2}$$
C(s) +  $\frac{1}{2}$ CO<sub>2</sub>(g)  $\Longrightarrow$  CO(g)

and the equilibrium pressure is 12 atm. Calculate  $K_p$ .

2. Calculate partial pressure of B at equilibrium in the following equilibrium

$$A(s) \rightleftharpoons B(g) + 2C(g); K_p = 32 \text{ atm}^3.$$

- **3.** In a gaseous reaction  $A + 2B \rightleftharpoons 2C + D$  the initial concentration of B was 1.5 times that of A. At equilibrium the concentrations of A and D were equal. Calculate the equilibrium constant  $K_C$ .
- 4. For the reaction

$$A(g) \rightleftharpoons B(g); K_C = 10$$

$$B(g) \rightleftharpoons C(g); K_C = 2$$

$$C(g) \rightleftharpoons D(g)$$
;  $K_C = 0.01$ 

Calculate  $K_C$  for the reaction  $D(g) \rightleftharpoons A(g)$ .

- **5.** 5 litre vessel contains 2 moles of each of gases A and B at equilibrium. If 1 mole each of A and B are removed. Calculate  $K_C$  for the reaction  $A(g) \rightleftharpoons B(g)$ .
- **6.** Calculate  $K_p$  for the reaction  $A(g) \rightleftharpoons B(s) + 2C(g)$ ;  $K_C = 0.2$  at 305 K.
- 7. A mixture of 3 moles of  $SO_2$ , 4 moles of  $NO_2$ , 1 moles of  $SO_3$  and 4 moles of NO is placed in a 2.0L vessel.  $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$ .


At equilibrium, the vessel is found to contain 1 mole of  $SO_2$ . Calculate the value of  $K_C$ .

- **8.** The density of an equilibrium mixture of  $N_2O_4$  and  $NO_2$  at 1 atm and 373.5K is 2.0g/L. Calculate  $K_C$  for the reaction  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$
- **9.** If chemical equilibrium is attained at standard states then what is the value of  $\Delta G^{\circ}$ ?
- **10.** Calculate the equilibrium concentration ratio of C to A if equimolar ratio of A and B were allowed to come to equilibrium at 300K.

$$A(g) + B(g) \rightleftharpoons C(g) + D(g)$$
;  $\Delta G^{\circ} = -830$  cal.

11. A definite amount of solid NH<sub>4</sub>HS is placed in a flask already containing ammonia gas at a certain temperature and 0.1 atm pressure. NH<sub>4</sub>HS decomposes to give NH<sub>3</sub> and H<sub>2</sub>S and at equilibrium total pressure in flask is 1.1 atm. If the equilibrium constant  $K_p$  for the reaction NH<sub>4</sub>HS(s)  $\rightleftharpoons$  NH<sub>3</sub>(g) + H<sub>2</sub>S(g) is represented as  $z \times 10^{-1}$  then find the value of z.

12. The gaseous reaction :  $A(g) + nB(g) \implies mC(g)$  is represented by following curves



What is the value of n + m?

## ANSWERS

## Level

																•			
1.	(b)	· 2. (	b)	3.	(c)	4.	(a)	5.	(b)	6.	(a)	7.	(a)	8.	(a)	9.	(a)	10.	(a)
11.	(b)	12. (	b)	13.	(b)	14.	(d)	15.	(c)	16.	(b)	17.	(d)	18.	(d)	19.	(c)	20.	(d)
21.	(b)	22. (	a)	23.	(c)	24.	(a)	25.	(b)	26.	(d)	27.	(d)	28.	(c)	29.	(d)	30.	(c)
31.	(d)	32. (	c)	33.	(c)	34.	(c)	35.	(a)	36.	(a)	37.	(c)	38.	(d)	39.	(c)	· 40.	(c)
41.	(a)	42. (	b)	43.	(a)	44.	(a)	45.	(a)	46.	(a)	47.	(c)	48.	(d)	49.	(d)	50.	(d)
51.	(a)	<b>52</b> . (	b)	53.	(a)	54.	(b)	55.	(b)	56.	(b)	57.	(b)	58.	(b)	59.	(c)	60.	(c)
61.	(a)	62. (	d)	63.	(a)	64.	(a)	65.	(c)	66.	(a)	67.	(a)	68.	(c)	69.	(d)	70.	(a)
71.	(a)	72. (	c)	73.	(a)	74.	(d)	75.	(b)	76.	(c)	77.	(b)	78.	(b)	79.	(c)	80.	(d)
81.	(b)	82. (	b)	83.	(d)	84.	(b)	85.	(b)	86.	(b)	87.	(a)	88.	(a)	89.	(b)	90.	(c)
91.	(a)	92. (	) :	93.	(a)	94.	(c)	95.	(b)	96.	(d)	97.	(d)	98.	(b)	99.	(c)	100.	(a)
101.	(d)	102. (	j) 10	03.	(b)	104.	(d)	105.	(c)	106.	(d)	107.	(b)	108.	(d)	109.	(a)	110.	(d)
111.	(c)	<b>112.</b> (i	o) 1	13.	(a)	114.	(b)	115.	(a)	116.	(c)	117.	(b)	118.	(a)	119.	(a)	120.	(c)
121.	(c)	122. (a	1) 1:	23.	(b)	124.	(d)	125.	(a)										

## Level 2

1.	(a)	2. (c)	3. (c)	<b>4</b> . (d)	5. (d)	6. (c)	7. (c)	8. (a)	9. (c)	10. (c)
11.	(b)	<b>12</b> . (d)	<b>13.</b> (b)	14. (b)	<b>15</b> . (a)	16. (a)	17. (a)	18. (b)	19. (c)	<b>20</b> . (c)
21.	(d)	<b>22.</b> (b)	<b>23</b> . (c)	<b>24.</b> (d)	25. (b)					

Passage-1:

1. (b)

2. (d)

3. (c)

Passage-2:

1. (a)

2. (b)

3. (b)

4. (a)

Passage-3:

1. (a)

**2.** (d)

3. (c)

Passage-4:

1. (b)

2. (a)

3. (b)

### One or More Answers is/are correct

1. (b,c,d)

2. (a,b,c)

3. (c,d)

4. (a,c)

5. (a,c,d)

6. (c,d)

7. (a,b,c)

8. (c,d)

9. (a,b,d)

10. (a,c)

11. (b,c,d)

12. (a,b,c)

### Match the Column

1.  $A \rightarrow P$ ;

 $B \rightarrow R$ ;

 $C \rightarrow Q$ ;

 $D \rightarrow S$ 

2.  $A \rightarrow R$ ;

 $B \rightarrow Q$ ;

 $C \rightarrow P$ ;

 $\mathsf{D} \to \mathsf{S}$ 

3.  $A \rightarrow P$ ;

 $B \rightarrow S$ ;

 $C \rightarrow Q$ ;

 $D \rightarrow R$ 

4.  $A \rightarrow Q$ ;

 $B \rightarrow P, R;$ 

 $C \rightarrow P, S;$ 

 $D \rightarrow P$ , S

5.  $A \rightarrow P$ ;

 $B \rightarrow R$ ;

 $C \rightarrow S$ ;

 $D \rightarrow Q$ 

**6.**  $A \rightarrow R$ ;

 $B \rightarrow Q$ ;

 $C \rightarrow P$ ;

 $D \rightarrow S$ 

7.  $A \rightarrow Q$ ;

 $B \rightarrow S$ ;

 $C \rightarrow P$ ;

 $D \rightarrow S$ 

### **Assertion-Reason Type Questions**

1. (D)

2. (A)

3. (D)

4. (B)

5. (A)

6. (C)

11. (B)

12. (A)

13. (A)

14. (C)

15. (A)

7. (C)

8. (A)

9. (A)

10. (D)

Subjective Problems

1. 4

2. 2

3. 4

4. 5

5. 1

6. 5

7. 9

8. 2

9. 0

10. 2

11. 3

**12**. 5

# Hints and Solutions

### Level 1

**2.** (b) 
$$Q_C = \frac{[P_2(g)]^4}{[P_2(g)]} = \frac{(1)^2}{(3/2)} = \frac{2}{3}$$

7. (a) 
$$Q_C = \frac{\left(\frac{6}{2}\right)^2}{\left(\frac{2}{2}\right)\left(\frac{4}{2}\right)^3} = \frac{9}{8}$$

 $Q_c < K_c$  so reaction will proceed in forward direction.

**36.** (a) 
$$K_p = \frac{P_{S_2(g)}^4}{P_{S_2(g)}} = \frac{(4 \times 0.3)^4}{0.70} = 2.96$$

**39.** (c) 
$$K_p = \frac{1}{P_{H_2O(g)}^4}$$

$$\Rightarrow P_{\mathsf{H}_2\mathsf{Q}(g)} = \left(\frac{1}{K_p}\right)^{1/4}$$

$$\Rightarrow$$
  $(10^{-12})^{1/4} = 10^{-3}$  atm

$$K_p = 10^{-3} \times 760 = 0.76 \text{ torr}$$

**40.** (c) 
$$P_{\text{H}_2\text{O}} = 2 \times 10^{-2} \text{ atm}$$

R. H. = 
$$\frac{2 \times 10^{-2} \times 760}{38} \times 100$$

**42.** (b) 
$$K_c = \frac{n_B n_C^3}{n_A^2} \times \frac{1}{V^2}$$

$$\Rightarrow 16 = \frac{2 \times 2^3}{2^2 \times V^2}$$

$$\Rightarrow V = \frac{1}{2}$$

**45.** (a) 
$$K_c = \frac{[H_2S(g)]}{[H_2(g)]}$$

$$\Rightarrow$$
 8 × 10<sup>-2</sup> =  $\frac{x}{0.3 - x}$ 

$$\Rightarrow x = 0.022$$

$$[H_2S(g)] = \frac{0.022}{2}$$

$$= 0.011 M$$

**47.** (c) 
$$P_{\text{total}} = 3P$$
  

$$\Rightarrow P = \frac{0.318}{3} = 0.106$$

$$\therefore K_p = 4P^3 = 4.76 \times 10^{-3}$$

54. (b) 
$$H_2(g) + CO_2(g) \rightleftharpoons H_2O(g) + CO(g)$$
  
moles of eqm  $0.8-x$   $0.8-x$   $x$ 

conc. at eqm 
$$\frac{0.8-x}{5}$$
  $\left(\frac{0.8-x}{5}\right)\left(\frac{x}{5}\right)\left(\frac{x}{5}\right)$ 

$$\therefore \quad \Delta n_g = 0$$

$$\therefore \quad K_c = \frac{\left(\frac{x}{5}\right)^2}{\left(\frac{0.8 - x}{5}\right)^2}$$

$$\Rightarrow$$
  $2 = \frac{x}{0.8 - x}$ 

$$\Rightarrow x = 0.533 [CO_2(g)] = \frac{0.8 - 0.533}{5}$$

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Initial moles 
$$a = 0$$
  
at eqm  $a(1-\alpha) = 2a\alpha$ 

$$= \frac{75}{100} \times \text{Volume at eqm}$$

at constant T and P:  $V \propto n$ 

$$a = 0.75 \times a(1 + \alpha)$$

$$\Rightarrow \alpha = 0.33$$

**56.** (b) 
$$CO + H_2O \rightleftharpoons CO_2 + H_2$$

Initial conc. 3 3 x 0
At eqm 2 2 (x+1) 1
$$5 = \frac{(x+1)}{4} = 20 = x+1$$

$$\Rightarrow x = 19$$

**67.** (a) 
$$\alpha = \frac{M_T - M_O}{(n-1) M_O}$$

$$\Rightarrow 0.2 = \frac{92 - M_O}{M_O}$$

$$\Rightarrow M_O = 76.66$$

$$d_{\text{mixture}} = \frac{PM_{\text{mixture}}}{RT}$$

$$= \frac{1 \times 76.66}{0.821 \times 300} = 3.11 \text{ g/litre}$$

**68.** (c) 
$$M_O = \frac{dRT}{P} = 57.47$$
  $\alpha = \frac{99 - 57.47}{57.47} = 0.72$ 

70. (a) 
$$A_2(g) + B_2(g) \implies 2AB(g)$$

Initial conc.  $10/2$   $15/2$   $5/2$ 

at eqm  $5-x$   $7.5-x$   $2.5+2x$ 
 $2.5+2x=7.5$ 
 $x=2.5$ 
 $K_c = \frac{(7.5)^2}{2.5 \times 5} = 4.5$ 

71. (a) 
$$H_2(g) + S(s) \rightleftharpoons H_2S(g)$$

Conc. at eqm 
$$0.5-x$$
 —  $x$ 

$$K_c = \frac{[H_2S]}{[H_2]}$$

$$\Rightarrow 7 \times 10^{-2} = \frac{x}{0.5-x}$$

$$\Rightarrow x = 0.0327$$

$$P_{H_2S} = \left(\frac{n_{H_2S}}{V}\right) RT$$

$$= 0.0327 \times 0.0821 \times 360 = 0.966$$

**72.** (c) The gaseous mixture contains 40% Cl<sub>2</sub> and 40% PCl<sub>3</sub>, since they are produced in 1 : 1 mole ratio. The PCl<sub>5</sub> % is 20.

For ideal gases mole % = volume %

$$P_{\text{Cl}_2} = P_{\text{PCl}_3}$$
⇒ 2 × 0.40 = 0.80 atm
$$P_{\text{PCl}_5} = 2 \times 0.2 = 0.40 \text{ atm}$$
∴ 
$$K_p = \frac{P_{\text{PCl}_3} \cdot P_{\text{Cl}_2}}{P_{\text{PCl}_5}}$$

$$= \frac{0.80 \times 0.80}{0.40}$$
= 1.6 atm

73. (a) For an ideal gas mole % ≡ volume %

$$K_p = \frac{P_{\text{H}_2\text{O}}^2}{P_{\text{H}_2}^2} = \frac{(0.55)^2}{(0.45)^2} = 1.49$$

74. (d) Moles of CO<sub>2</sub> present at equilibrium  $= \frac{1.642 \times 50}{0.0821 \times 1000} = 1$ 

Mole % of  $XCO_3$  decomposed =  $\frac{1}{4} \times 100 = 25\%$ 

Hence, 75% remains undecomposed.

**75.** (b) Let initial moles of  $H_2(g)$  is 1

$$\operatorname{Fe}_{2}O_{3}(s) + 3H_{2}(g) \rightleftharpoons 2\operatorname{Fe}(s) + 3H_{2}O(g)$$

at eqm 
$$K_c = \frac{\left(\frac{3x}{V}\right)^3}{\left(\frac{1-3x}{V}\right)^3}$$

$$\Rightarrow 8 = \left(\frac{3x}{1-3x}\right)^3$$

$$\Rightarrow x = 0.22$$
% of H<sub>2</sub> unreacted
$$= \frac{1-3 \times 0.22}{1} \times 100 = 34$$

76. (c) 
$$AB_3(g) \rightleftharpoons AB_2(g) + \frac{1}{2}B_2(g)$$
At eqm  $800-x$   $x$   $x/2$ 
 $800-x+x+\frac{x}{2}=900 \Rightarrow x=200$ 
% dissociated  $=\frac{200}{200} \times 100=25$ 

77. (b) Let P is initial pressure of NO<sub>2</sub>

$$2NO_{2}(g) \rightleftharpoons 2NO(g) + O_{2}(g)$$
At eqm  $P - 2x$   $2x$   $x$ 
as per given  $x = 0.25$ 

$$K_{p} = \frac{(2x)^{2}(x)}{(P - 2x)^{2}}$$

$$\Rightarrow 156.25 = \frac{(0.5)^{2}(0.25)}{P_{NO_{2}}^{2}}$$

$$\Rightarrow P_{NO_{2}} = 0.02$$

78. (b) 
$$2\text{NOCl}(g) \implies 2\text{NO}(g) + \text{Cl}_2(g)$$
At eqm  $P - 2x$   $2x$   $x$ 

$$P + x = 1; P - 2x = 0.64, x = 0.12$$

$$K_p = \frac{(0.24)^2 (0.12)}{(0.64)^2}$$

$$= 16.875 \times 10^{-3} \text{ atm}$$

**79.** (c) 
$$Q_c = \frac{1 \times 1}{1 \times 1} = 1$$

 $Q_c > K_c$  so reaction will proceed in backward direction

$$A_{2}(g) + B_{2}(g) \rightleftharpoons C_{2}(g) + D_{2}(g)$$
Conc. eqm  $\frac{1+x}{10}$   $\frac{1+x}{10}$   $\frac{1-x}{10}$   $\frac{1-x}{10}$ 

$$0.25 = \frac{\left(\frac{1-x}{10}\right)^{2}}{\left(\frac{1+x}{10}\right)^{2}} \implies x = 0.333$$

$$[A_2(g)] = \frac{1+x}{10} = \frac{1.333}{10} = 0.133$$

**83.** (d) 
$$K_c = [Hg(g)]; P = CRT$$

$$C = \left(\frac{0.002}{760}\right) \times \frac{1}{(0.0821 \times 300)}$$

$$= 1.068 \times 10^{-7} M$$

**84.** (b) 
$$K_c = \frac{n_{AB_2}^2}{n_{A_2} \cdot n_{B_2}^2} \times V = \frac{n_{AB_2}^2}{n_{A_2} \cdot n_{B_2}^2} \times \left(\frac{n_{\text{total}}RT}{P}\right)$$

$$= \frac{2^2}{5 \times 3^2} \times \left(\frac{10 \times 0.0821 \times 300}{8.21}\right)$$

104. (d) Concentration of all gases increases and equilibrium shifts toward less no. of moles but new equilibrium concentration of every gas would be higher than earlier.

116. (c) 
$$\ln \frac{K_{P_2}}{K_{P_1}} = \frac{\Delta H^{\circ}}{R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$
  
=  $\ln \left( \frac{10^{-4}}{10^{-6}} \right) = \frac{\Delta H^{\circ}}{8.314} \left( \frac{25}{298 \times 323} \right)$ 

$$\Delta_r H^\circ = 147.41 \text{ kJ/mol}$$
 $\Delta_r G^\circ = \Delta_r H^\circ - T \cdot \Delta_r S^\circ$ 
 $= -30 - 300 \times 0.1 = 0$ 
 $\Delta_r G^\circ = -2.303 RT \log K$ 

124.(d) 
$$P_{\text{total}} = 0.2 \,\text{bar}$$
  
At equilibrium  $P_{\text{CO}_2} = P_{\text{H}_2\text{O}} = 0.1 \,\text{bar}$   
 $K_p = (0.1)^2 = 0.01$   
 $\Delta_r G^\circ = -RT \,\ln K_p$   
 $= -8.314 \times 420 \,\ln (0.01)$ 

= 16083.6 J/mol or 16.083 kJ/mol

### Level 2

1. (a) 
$$C(s) + CO_2(g) \rightleftharpoons 2CO(g);$$
 ...(1)  
 $K_{p_1} = 10^{14} \text{ atm}$   
 $2CO(g) + 2Cl_2(g) \rightleftharpoons 2COCl_2(g);$  ...(2)  
 $K_{p_2} = (6 \times 10^{-3})^2 \text{ atm}^{-2}$   
Add (1) and (2)  
 $C(s) + CO_2(g) + 2Cl_2(g) \rightleftharpoons 2COCl_2(g);$   
 $K_p = 10^{14} \times 36 \times 10^{-6}$   
 $= 36 \times 10^8$   
For given reaction  $\Delta n_g = -1$   
 $\therefore K_c = K_p(RT)$   
 $= 36 \times 10^8 \times 0.0821 \times 1120$   
 $K_c = 3.31 \times 10^{11} M^{-1}$ 

2. (c) 
$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$
Initial moles 1 2 0
at eqm 1-x 2-3x 2x
where  $x = 0.5$ 

$$K_c = \frac{(1/4)^2}{\left(\frac{0.5}{4}\right)\left(\frac{0.5}{4}\right)^3} = 256$$

equilibrium constant for the reaction

$$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$$

$$K_c^1 = \frac{1}{\sqrt{K_c}} = \frac{1}{16}$$

3. (c) 
$$A_2(g) + B_2(g) \implies 2AB(g)$$

Moles at eqm  $2-x$   $4-x$   $2x$ 
 $K_c = \frac{4x^2}{(2-x)(4-x)}$ 
 $\Rightarrow x = \frac{32}{24} = 1.33 \text{ mole}$ 
 $[AB(g)] = \frac{2 \times 1.33}{4} = 0.66 \text{ M}$ 

**4.** (d) 
$$P_{\text{total}} = P_{\text{HNO}_3} + P_{\text{NO}_2} + P_{\text{H}_2\text{O}} + P_{\text{O}_2}$$
  
 $\therefore P_{\text{NO}_2} = 4P_{\text{O}_2} \text{ and } P_{\text{H}_2\text{O}} = 2P_{\text{O}_2}$   
 $\therefore P_{\text{total}} = P_{\text{HNO}_3} + 7P_{\text{O}_2}$   
 $\Rightarrow 30 - 2 = P_{\text{O}_2} \times 7$ 

$$\Rightarrow P_{O_2} = \frac{28}{7} = 4$$

$$K_p = \frac{P_{NO_2}^4 \cdot P_{H_2O} \cdot P_{O_2}}{P_{HNO_3}^4}$$

$$= \frac{(4 \times 4)^4 \times (2 \times 4)^2 \times 4}{2^4} = 2^{20}$$

$$K_p = K_c (RT)^{\Delta t_g} = K_c (0.08 \times 400)^3$$

$$K_c = \frac{2^{20}}{(32)^3} = 32$$

5. (d) LiCl 
$$3NH_3(s) \rightleftharpoons LiCl \cdot NH_3(s) + 2NH_3(g)$$
;  

$$[K_p = 9 \text{ atm}^2]$$

Therefore,

Let initial moles of  $NH_3$  is a for completion of reaction.

At eqm 
$$K'_p = \frac{1}{(P'_{NH_3})^2}$$
 or  $\frac{1}{9} = \frac{1}{(P'_{NH_3})^2}$   
 $\therefore P'_{NH_3} = 3 \text{ atm}$   
 $\therefore PV = nRT \implies 3 \times 5 = n \times 0.0820 \times 310$   
 $\therefore n = 0.59$  i.e.,  $(a - 0.2) = 0.59$   
 $\therefore \text{ initial moles of NH}_3 = 0.79$ 

**6.** (c)  $NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$ 

Let partial pressure at equilibrium of  $CO_2$  be P, then  $P_{NH_3} = 2P$  and total pressure at equilibrium = 3P

$$K_p = (2P)^2 \times P = 4P^3$$
 ...(1)

If  $NH_3$  is added and the pressure of  $NH_3$  after addition at equilibrium is 3P

$$K_p = 4P^3 = (P'_{NH_3})^2 \times P'_{CO_2}$$

$$\Rightarrow \qquad P'_{CO_2} = \frac{4}{9}P$$

$$\therefore \text{ ratio of } \frac{P'_{CO_2}}{P_{CO_2}} = \frac{4}{9}$$

7. (c) 
$$C_2H_6(g) \rightleftharpoons C_2H_4(g) + H_2(g)$$
At eqm  $1-x$   $x$   $x$ 

$$K_p = \frac{P_{C_2H_4} \cdot P_{H_2}}{P_{C_2H_6}} \Rightarrow \frac{x^2}{1-x} = 5 \times 10^{-2}$$

$$x^2 + 0.05 x - 0.05 = 0$$

$$x = \frac{-0.05 + \sqrt{(0.05)^2 + 4 \times 0.05}}{2} = 0.20 \text{ atm}$$
Partial pressure of C<sub>2</sub>H<sub>6</sub> = mole fraction × total pressure

⇒ 0.80 = mole fraction ×1.2  
∴ mole % of 
$$C_2H_6 = \frac{0.8}{1.2} \times 100 = 66.66$$

8. (a) 
$$2\text{NOBr}(g) \rightleftharpoons 2\text{NO}(g) + \text{Br}_2(g)$$

Initial pressure  $P = 0 = 0$ 
at eqm  $P - 2x = 2x = x$ 

where  $2x = 0.40 P$ 
 $\Rightarrow x = 0.20 P$ ;
 $\Rightarrow 1.20 P = 0.30$ 
 $\Rightarrow P = 0.25 \text{ atm}$ 
 $K_p = \frac{P_{\text{NO}}^2 \cdot P_{\text{Br}_2}}{P_{\text{NOBr}}^2}$ 
 $= \frac{(0.4P)^2 (0.2P)}{(0.6P)^2} = 0.0222$ 

 $K_p \text{ for the reaction}$   $2NO(g) + Br_2(g) \iff 2NOBr(g)$ is  $\frac{1}{0.0222} = 45$ 

**9.** (c) : Initially only *A* is present so at eqm *B* and *C* should be present in 2 : 1

$$2A(g) \rightleftharpoons 2B(g) + C(g)$$

At eqm 400 mL 200 mL 100 mL

for ideal gases volume % = mole %

$$K_p = \frac{\left(\frac{200}{700} \times 10\right)^2 \left(\frac{100}{700} \times 10\right)}{\left(\frac{400}{700} \times 10\right)^2} = \frac{10}{28}$$

10. (c) 
$$SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$$
  
Initial moles 2 2 2 2  
at eqm  $2-x$   $2-x$   $2+x$   $2+x$   $(\because Q_p < K_p)$ 

Total no. of moles of gases at equilibrium = 8 + 2 = 10

$$K_p = \frac{P_{SO_3} \cdot P_{NO}}{P_{SO_2} \cdot P_{NO_2}}$$

$$\Rightarrow \qquad 25 = \frac{\left(\frac{2+x}{10} \times P\right)^2}{\left(\frac{2-x}{10} \times P\right)^2}$$

$$\Rightarrow \qquad 5 = \frac{2+x}{2-x}$$

$$\Rightarrow$$
  $x = 1.33$ 

Partial pressure of NO<sub>2</sub> =  $\frac{2-x}{10} \times P_{\text{total}}$  $=\frac{0.666}{10}\times 2$ = 0.133 atm

**11.** (b) 
$$3Se_2(g) \rightleftharpoons Se_6(g)$$

moles at eqm 
$$a(1-\alpha)$$

Observed no. of moles at equilibrium  $M_T$ Initial no. of moles of reactant

$$\Rightarrow \frac{a(1-\alpha)+\frac{a\alpha}{3}}{a}=\frac{M_T}{M_O}=\alpha=0.315$$

where molar mass of Se,

$$(M_T) = 79 \times 2 = 158$$

and molar mass of mixture

$$(M_O) = \frac{wRT}{PV} = \frac{0.02 \times 24.63}{1 \times 2.463 \times 10^{-3}} = 200$$

**12.** (d) 6HCHO 
$$\rightleftharpoons$$
 C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>

Conc. at eqm 
$$C(1-\alpha)$$

Observed moles concentration  $M_T$ Initial moles concentration

 $M_T$  = Theoretical molar mass of HCHO

$$\frac{C(1-\alpha) + \frac{C\alpha}{6}}{C} = \frac{M_T}{M_O} = \frac{30}{150}$$

$$\alpha = 0.96$$

**13.** (b) 
$$2SO_2(g) + O_2(g) \implies 2SO_3(g)$$

$$K_c = \frac{(0.12)^2}{(0.12)^2 \times 5} = 0.2$$

Another vessel

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

moles at eqm 0.5-2x y-x

as per given  $2x = \frac{20}{100} \times 0.5 = 0.1$ 

$$K_c = \frac{(0.1)^2}{(0.4)^2 (y - 0.05)} = 0.20$$

$$y = 0.3625$$
 mole

$$\dots$$
 mass of O<sub>2</sub> added = 11.6 g

### **14.** (b) Let initial mole of $N_2O_4(g)$ is one

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Initial moles

At equilibrium

Total no. of moles at equilibrium =  $1 + \alpha$ 

$$P_{N_2O_4} = \frac{1-\alpha}{1+\alpha} \times P$$

$$P_{\text{NO}_2} = \frac{2\alpha}{(1+\alpha)} \times P$$

Hence, 
$$K_p = \frac{P_{NO_2}^2}{P_{NO_2}} = \frac{4\alpha^2}{(1 - \alpha^2)} \times P$$

$$4.5 = \frac{4\alpha^2}{(1 - \alpha^2)} \times 2$$

$$\Rightarrow$$
  $\alpha = 0.6$ 

Mole fraction of N2O4:

$$X_{\rm N_2O_4} = \frac{1-\alpha}{1+\alpha} = 0.25$$

$$X_{NO_2} = 0.75$$

Average molar mass of mixture

$$= 0.25 \times 92 + 0.75 \times 46 = 57.5$$

**15.** (a) 
$$2AB(s) \iff A_2(g) + B_2(g)$$

$$0.5 + x$$
  $x$ 

$$K_p = P_{A_2} \cdot P_{B_2}$$

$$\Rightarrow 0.06 = (0.5 + x) x$$

$$\Rightarrow 0.00 = (0.5 + x) x$$
$$= x^2 + 0.5x - 0.06$$

$$\Rightarrow x = 0.1$$

$$P_{\text{total}} = P_{A_2} + P_{B_2} = 0.6 + 0.1 = 0.70 \text{ atm}$$

**16.** (a) 
$$Sb_2S_3(s) + 3H_2(g) \longrightarrow 2Sb(s) + 3H_2S(g)$$
  
 $0.01 - x$   $0.01 - 3x$   $2x$   $3x$ 

where 3x = 0.005

$$H_2S + Pb^{2+} \longrightarrow PbS + 2H^+$$

no. of moles of PbS formed

$$=\frac{1.19}{238}=0.005$$
 mole

At eqm[H<sub>2</sub>] = 
$$\left[\frac{0.005}{250}\right]$$
 =  $\left(\frac{.01 - .005}{250}\right)$ ;

$$K_c = \left(\frac{0.005}{0.005}\right)^3 = 1$$

17. (a) 
$$2A(g) + B(g) \iff C(g) + D(g)$$

Initial moles at eqm

Due to very high value of  $K_c$  we can assume that reactant almost converted into products so

$$1 - x = y; \quad 2 - 2x = 2y \Rightarrow \quad x \approx 1$$

$$10^{12} = \frac{8 \times 4}{(2y)^2 y}$$

$$\Rightarrow \qquad y^3 = 8 \times 10^{-12}$$

: equilibrium concentration of

$$A = 2y = 4 \times 10^{-4}$$

**18.** (b) 
$$K_c = \frac{\text{[complex]}}{\text{[H}_3 B O_3] \text{[glycerin]}} = 0.90$$

⇒  $\frac{\text{[complex]}}{\text{[H}_3 B O_3]} = \frac{80}{20}$ 

∴  $\text{[glycerin]} = \frac{80}{20 \times 0.9} = 4.44 M$ 

 $H_3BO_3 + glycerin \iff (H_3BO_3 - glycerin)$ 

At eq<sup>m</sup> 0.1-x 
$$a-x$$
 x  $x = 0.1 \times \frac{80}{100} = 0.08;$   $a-x = 4.44$   $\Rightarrow a = 4.44 + 0.08 = 4.52 M$ 

19. (c) 
$$\frac{r_{\text{mix}}}{r_{\text{O}_2}} = \sqrt{\frac{32}{M_{\text{mix}}}} = 0.4\sqrt{5}$$

$$\Rightarrow \qquad M_{\text{mix}} = 40\text{g/mol}$$

$$O_2(g) \implies \frac{2}{3}O_3(g)$$
Initial moles 1 0
at eqm 1-\alpha \frac{2}{3}\alpha

$$\frac{.32}{40} = \frac{1-\alpha+\frac{2}{3}\alpha}{1}$$

$$\Rightarrow \qquad \alpha = 0.6$$

$$\therefore \qquad \% \ \alpha = 60$$

moles at eqm 1-2x

**20.** (c) 
$$2SO_3(g) \iff 2SO_2(g) + O_2(g)$$

Only SO<sub>2</sub> (O. No. = 4) will oxidized so equivalent of SO<sub>2</sub> = equivalent of KMnO<sub>4</sub>  $2x \times 2 = 0.2 \times 5$ 

$$\Rightarrow 2x = 0.5$$

$$K_c = \frac{\left(\frac{0.5}{2}\right)^2 \left(\frac{0.25}{2}\right)}{\left(\frac{0.5}{2}\right)^2} = 0.125$$

21. (d) 
$$\frac{r_{\text{mix}}}{r_{\text{SO}_2}} = \sqrt{\frac{M_{\text{SO}_2}}{M_{\text{mix}}}}$$

$$\Rightarrow 2.56 = \frac{64}{M_{\text{mix}}}$$

$$\Rightarrow M_{\text{mixture}} = 25$$
Let mole fraction of  $F_2$  is  $x$ 

$$25 = \frac{38 \times x + (1 - x) \times 19}{1}$$

$$x = 0.315;$$

$$\Rightarrow K_p = \frac{P_F^2}{P_{F_2}} = \frac{(0.685 \, P)^2}{(0.315 \, P)} \approx 1.49 \text{atm}$$
22. (b)  $8 \times 10^{-6} = \frac{x^2}{2}$ 

**22.** (b) 
$$8 \times 10^{-6} = \frac{x^2}{0.5}$$
  
 $\Rightarrow x = 2 \times 10^{-3}$   
 $\therefore \text{ pOH} = 2.7$   
So,  $\text{pH} = 11.3$ 

23. (c) 
$$\triangle G^{\circ} = 0 - 77.1 \times 2 = -154.2 \text{ kJ/mol}$$

$$Q = \frac{[H^{+}]^{2}}{P_{H_{2}} [Ag^{+}]^{2}}$$

$$= \frac{10^{-6}}{0.5 \times (10^{-10})} = 2 \times 10^{4}$$

$$\triangle G = \triangle_{r}G^{\circ} + RT \ln Q$$

$$\triangle_{r}G = -154.2 + \frac{8.314 \times 300 \ln (2 \times 10^{4})}{1000}$$

$$= -129.5 \text{ kJ/mol}$$

$$24. \text{ (d)} \quad N_2O_5 \iff N_2O_3 + O_2$$

$$4-x \qquad x-y \qquad x+y$$

$$N_2O_3 \iff N_2O + O_2$$

$$x-y \qquad y \qquad y+x$$

$$\therefore [O_2] = x+y=2.5$$

$$\text{and} \quad 2.5 = \frac{(x+y)(x-y)}{4-x}$$

$$\therefore \qquad x=2.166$$

$$[N_2O_5] = 4-x=1.846$$

**25.** (b) Let *x* is partial pressure of *A* and *y* is partial pressure of *C* when both equilibrium simultaneously established in a vessel

$$X(s) \iff A(g) + 2B(g)$$

$$x \qquad (2x + 2y);$$

$$K_{P_1} = P_A \cdot P_{B(\text{total})}^2$$

$$Y(s) \iff C(g) + 2B(g)$$

$$y \qquad (2y + 2x);$$

$$K_{P_2} = P_C \cdot P_{B(\text{total})}^2$$

$$\frac{K_{p_1}}{K_{p_2}} = \frac{x}{y} \implies x = 2y$$

$$K_{p_1} = x(2x + 2y)^2$$

$$\implies x = 0.1 \text{ atm};$$

$$\therefore y = 0.05 \text{ atm}$$
Total pressure of gases =  $P_A + P_B + P_C$ 

$$= 3(x + y)$$

$$= 0.45 \text{ atm}.$$

#### Level 3

### Passage-3

$$N_{2}O_{3}(g) \rightleftharpoons NO_{2}(g) + NO(g) ; K_{p_{1}} = 2.5 \text{ bar}$$

$$2NO_{2}(g) \rightleftharpoons N_{2}O_{4}(g) ; K_{p_{2}}$$

$$x - y \qquad y/2$$

$$P_{NO} = x = 1.5 \text{ b}$$

$$K_{p_{1}} = \frac{P_{NO} \cdot P_{NO_{2}}}{P_{N_{2}O_{3}}}$$

$$2.5 = \frac{x(x - y)}{2 - x} = \frac{1.5(1.5 - y)}{2 - 1.5} = 4.5 - 3y$$

$$\Rightarrow y = \frac{2}{3} \text{ Bar}$$

$$(1) P_{N_{2}O_{3}} = 2 - x = 2 - 0.5 = 0.5 \text{ b}$$

$$(2) P_{NO_{2}} = x - y = 1.5 - \frac{2}{3} = \frac{2.5}{3} \text{ b}$$

$$(3) K_{p_{2}} = \frac{y}{2(x - y)^{2}} = \frac{2/3}{2\left(\frac{2.5}{3}\right)^{2}} = 0.48 \text{ b}^{-1}$$

### One or More Answers is/are correct

6. (c, d) Le Chatelier's principle is not quantitative. If both stress would cause the same direction of shift, the shift is determinable. If the two stresses would cause shifts in opposite directions, no deduction is possible.

### **Subjective Problems**

11. 
$$NH_4HS(s) \Longrightarrow NH_3(g) + H_2S(g)$$

At eqm.  $-(01+P) P$ ;

As per given  $01 + 2P = 11$ ;  $P = 0.5$ 
 $K_P = P_{NH_3} \cdot P_{H_2S} = (0.1+P) \times P$ 
 $= 0.6 \times 0.5 = 0.3 = 3 \times 10^{-1} = 3$ 

12.  $A + nB \Longrightarrow mC$ 
 $t = 0$   $8M$   $12M$ 
 $t = t_{eq}$   $8 - x$   $12 - nx$   $mx$ 

Given:  $[A]_{eq} = 6M$ 
 $8 - x = 6$ 
 $x = 2M$ 
 $[B]_{eq} = 8M$ 
 $12 - nx = 8$ 
 $n(2) = 4$ 
 $n = 2$ 
 $[C]_{eq} = 6M$ 
 $mx = 6$ 
 $m(2) = 6$ 
 $m = 3$ 

m + n = 3 + 2 = 5