

Properties and Types of solid

- The three states of matter are solid, liquid and gas. 1. Which of the following statement is/are true about them [AIIMS 1991]
 - (a) Gases and liquids have viscosity as a common propertv
 - (b) The molecules in all the three states possess random translational motion
 - (c) Gases cannot be converted into solids without passing through the liquid phase
 - (d) Solids and liquids have vapour pressure as a common property
- A pure crystalline substance, on being heated gradually, 2. first forms a turbid looking liquid and then the turbidity completely disappears. This behaviour is the characteristic of substances forming [BHU 2000]
 - (a) Isomeric crystals (b) Liquid crystals
 - (c) Isomorphous crystals (d) Allotropic crystals
- Which of the following is ferroelectric compound 3.

[AFMC 1997]

(a) $BaTiO_3$ (b) $K_4 [Fe(CN)_6]$

(d) $PbZrO_3$ (c) Pb_2O_3

- Solid CO_2 is an example of 4.
 - (a) Molecular crystal (b) Ionic crystal
 - (c) Covalent crystal (d) Metallic crystal
- Value of heat of fusion of NaCl is 5.
 - (a) Very low
 - (b) Very high
 - (c) Not very low and not very high
 - (d) None of the above
- Piezoelectric crystals are used in 6.
 - (a) TV
 - (c) Record player (d) Freeze
- Which of the following is true for diamond 7.
 - (a) Diamond is a good conductor of electricity

(b) Radio

- (b) Diamond is soft
- (c) Diamond is a bad conductor of heat
- (d) Diamond is made up of C, H and O
- 8. *NaCl* is an example of
 - (a) Covalent solid (b) Ionic solid
 - (d) Metallic solid (c) Molecular solid
- Amorphous substances show 9.
 - (A) Short and long range order
 - (B) Short range order

- (C) Long range order
- (D) Have no sharp M.P.
- (a) A and C are correct
- (b) B and C are correct
- (c) C and D are correct (d) B and D are correct
- The characteristic features of solids are 10. [AMU 1994]
 - (a) Definite shape
 - (b) Definite size
 - (c) Definite shape and size
 - (d) Definite shape, size and rigidity
- Which one of the following is a good conductor of 11. electricity
 - [MP PMT 1994; AFMC 2002]
 - (b) Graphite
 - (c) Silicon

(a) Diamond

12.

14.

- (d) Amorphous carbon [Kerala CET (Med.) 2003]
- A crystalline solid (a) Changes abruptly from solid to liquid when heated
 - (b) Has no definite melting point
 - (c) Undergoes deformation of its geometry easily
 - (d) Has an irregular 3-dimensional arrangements
 - (e) Softens slowly
- Diamond is an example of 13.

[MP PET/PMT 1998; CET Pune 1998]

- (a) Solid with hydrogen bonding
- (b) Electrovalent solid
- (c) Covalent solid
- (d) Glass
- The solid *NaCl* is a bad conductor of electricity since
 - [AIIMS 1980]
 - (a) In solid *NaCl* there are no ions
 - (b) Solid NaCl is covalent
 - (c) In solid *NaCl* there is no velocity of ions
- (d) In solid *NaCl* there are no electrons
- The existence of a substance in more than one solid 15. modifications is known as or Any compound having more than two crystal structures is called
 - [MP PMT 1993; MP PET 1999] (b) Isomorphism
 - (a) Polymorphism
 - (c) Allotropy (d) Enantiomorphism [MP PET 1995]
- Which is not a property of solids 16.
 - (a) Solids are always crystalline in nature
 - (b) Solids have high density and low compressibility
 - (c) The diffusion of solids is very slow
 - (d) Solids have definite volume
- Which solid will have the weakest intermolecular forces 17.
 - (a) Ice (b) Phosphorus
 - (d) Sodium fluoride (c) Naphthalene
- Dulong and Petit's law is valid only for [KCET 2004] 18.
 - (a) Metals (b) Non-metals (c) Gaseous elements
 - (d) Solid elements
- 19. Which of the following is an example of metallic crystal solid (a) C
 - (b) Si

	(c) W	(d) $AgCl$	
20.	Under which category iodine crystals are placed among		
	the following		
	(a) Ionic crystal	(b) Metallic crystal	
	(c) Molecular crystal	(d) Covalent crystal	
21.	Among solids the highest m	elting point is established by	
		[Kerala CET (Med.) 2002]	
	(a) Covalent solids	(b) Ionic solids	
00	(c) recursion solution T_{0} and T_{0} recursion	(d) Molecular solids	
22,	to silicon should have which	th of the following number of	
	valence electrons	[KCET (Engg.) 2001]	
	(a) 1	(b) 2	
	(c) 3	(d) 5	
23.	Which of the following is no	on-crystalline solid	
	(a) CsCl	(b) NaCl	
	(c) CaF_2	(d) Glass	
24.	The lustre of a metal is due	to [AFMC 1998]	
	(a) Its high density	(b) Its high polishing	
-1t-	(c) Its chemical inertness	(d) Presence of free	
	A crystalline solid have		
23.	(a) Long range order	(b) Short range order	
	(c) Disordered arrangemer	(d) None of these	
26.	Crystalline solids are	[Pb. PMT 1999]	
	(a) Glass	(b) Rubber	
	(c) Plastic	(d) Sugar	
27.	Davy and Faraday proved th	nat [Kerala CET (Med.) 2002]	
	(a) Diamond is a form of ca	arbon	
	(b) The bond lengths of ca	arbon containing compounds	
	are always equal		
	(c) The strength of graphi	ite is minimum compared to	
	(d) Graphite is very hard		
	Which one of the fo		
28		ollowing metal oxides is	
28.	antiferromagnetic in nature	bllowing metal oxides is [MP PET 2002]	
28.	antiferromagnetic in nature (a) MnO_2	bllowing metal oxides is [MP PET 2002] (b) <i>TiO</i> 2	
28.	antiferromagnetic in nature (a) MnO_2 (c) VO_2	bllowing metal oxides is (b) <i>TiO</i> ₂ (c) <i>CrO</i> .	
28.	antiferromagnetic in nature (a) MnO_2 (c) VO_2	billowing metal oxides is (MP PET 2002] (b) TiO_2 (d) CrO_2 regioned together due to	
28. 29.	antiferromagnetic in nature (a) MnO_2 (c) VO_2 In graphite, carbon atoms a	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002]	
28. 29.	antiferromagnetic in nature (a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding	billowing metal oxides is (MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces	
28. 29.	antiferromagnetic in nature (a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding	
28. 29. 30.	(a) Ionic bonding (c) Metallic bonding (c) Metallic bonding (c) Metallic bonding	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals	
28. 29. 30.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding Which of the following is no	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002]	
28. 29. 30.	(a) Ionic bonding (c) Metallic bonding (c) Metallic bonding (c) Metallic bonding (c) They possess high melti	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point	
28. 29. 30.	antiferromagnetic in nature (a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point	
28. 29. 30.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of	ollowing metal oxides is (m) PET 2002] (b) TiO2 (d) CrO2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point	
28. 29. 30.	(a) They possess high melti (b) All are electrolyte (c) Exhibit directional prop	billowing metal oxides is [MP PET 2002] (b) TiO_2 (c) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond	
28. 29. 30.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a reference of the following is a referenc	billowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal	
28. 29. 30. 31.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i>	ollowing metal oxides is (b) TiO2 (d) CrO2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) NaCl	
28. 29. 30. 31.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite	oblowing metal oxides is (b) TiO2 (d) CrO2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) NaCl (d) Ice	
28. 29. 30. 31. 32.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet	Dellowing metal oxides is (D) metal oxides is $(MP PET 2002]$ (B) TiO_2 (D) CrO_2 re joined together due to $[AFMC 2002]$ (D) Vander Waal's forces (d) Covalent bonding (D) Vander Waal's forces (d) Covalent bonding (D) rest of the bond (D) molecular crystal (D) NaCl (d) Icety of $[Pb. PMT 2000]$	
28. 29. 30. 31. 32.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet (a) Silica	Dellowing metal oxides is $(mr) PET 2002$] $(b) TiO_2$ $(d) CrO_2$ re joined together due to $[AFMC 2002]$ $(b) Vander Waal's forces$ $(d) Covalent bonding(d) Ice(d) Icety of(pb. PMT 2000](b) Sodium silicate$	
28. 29. 30. 31. 32.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet (a) Silica (c) Silicon carbide	bllowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) $NaCl$ (d) Ice ty of [Pb. PMT 2000] (b) Sodium silicate (d) Silicon	
28. 29. 30. 31. 32. 33.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet (a) Silica (c) Silicon carbide Which type of solid crys	bllowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) $NaCl$ (d) Ice ty of [Pb. PMT 2000] (b) Sodium silicate (d) Silicon stals will conduct heat and	
28. 29. 30. 31. 32. 33.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Exhibit the property of (d) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet (a) Silica (c) Silicon carbide Which type of solid crys electricity	bllowing metal oxides is [MP PET 2002] (b) TiO_2 (d) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) $NaCl$ (d) Ice ty of [Pb. PMT 2000] (b) Sodium silicate (d) Silicon stals will conduct heat and	
28. 29. 30. 31. 32. 33.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variet (a) Silica (c) Silicon carbide Which type of solid cryst electricity (a) Ionic	ollowing metal oxides is (m) PET 2002] (b) TiO2 (d) CrO2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (d) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) NaCl (d) Ice ty of [Pb. PMT 2000] (b) Sodium silicate (d) Silicon stals will conduct heat and [RPET 2000] (b) Covalent	
28. 29. 30. 31. 32. 33.	(a) MnO_2 (c) VO_2 In graphite, carbon atoms a (a) Ionic bonding (c) Metallic bonding (c) Metallic bonding Which of the following is no (a) They possess high melti (b) All are electrolyte (c) Exhibit the property of (d) Exhibit directional prop Which of the following is a r (a) <i>SiC</i> (c) Graphite Quartz is a crystalline variett (a) Silica (c) Silicon carbide Which type of solid cryster electricity (a) Ionic (c) Metallic	bllowing metal oxides is [MP PET 2002] (b) TiO_2 (c) CrO_2 re joined together due to [AFMC 2002] (b) Vander Waal's forces (c) Covalent bonding ot correct for ionic crystals [Orissa JEE 2002] ing point and boiling point isomorphism perties of the bond molecular crystal (b) $NaCl$ (c) Ice ty of [Pb. PMT 2000] (b) Sodium silicate (c) Silicon stals will conduct heat and [RPET 2000] (b) Covalent (c) Molecular	

34.	Which of the following is an example of covalent crystal
	solid

- (a) Si (b) NaF
- (c) *Al* (d) *Ar*
- **35.** Which of the following is an example of ionic crystal solid
 - (a) Diamond (b) *LiF*
 - (c) *Li* (d) Silicon
- 36. Which one is an example of amorphous solid(a) Glass(b) Salt
 - (c) Cesium chloride (d) Calcium fluoride
- **37.** Silicon is [MHCET 2004] (a) Semiconductor (b) Insulator
 - (c) Conductor (d) None of these
- **38.** Which of the following statements about amorphous solids is incorrect **[KCET 2004]**
 - (a) They melt over a range of temperature
 - (b) They are anisotropic
 - (c) There is no orderly arrangement of particles
 - (d) They are rigid and incompressible

39. The ability of a given substance to assume two or more crystalline structure is called [DCE 2004]
(a) Amorphism (b) Isomorphism

- (c) Polymorphism
- 40. Glass is

2.

(a) Supercooled liquid (b) Crystalline solid

(d) Isomerism

(c) Amorphous solid (d) Liquid crystal

Crystallography and Lattice

- **1.** The correct statement in the following is **[MP PET 1997]**
 - (a) The ionic crystal of AgBr has Schottky defect
 - (b) The unit cell having crystal parameters, $a = b \neq c$,
 - $\alpha = \beta = 90^{\circ}, \gamma = 120^{\circ}$ is hexagonal
 - (c) In ionic compounds having Frenkel defect the ratio $\frac{\gamma_+}{\gamma_-}$ is high

(d) The coordination number of Na^+ ion in NaCl is 4

· · ·				
Which of the following is correct				[DPMT 1997]
	Crystal system	Axial distance	Axial angles	Examples
(a)	Cubic	$a \neq b = c$	$\begin{array}{l} \alpha = \beta \neq \gamma \\ = 90^{\rm o} \end{array}$	Cu, KCl
(b)	Monoclinic	$a \neq b = c$	$ \begin{array}{l} \alpha = \beta = \gamma \\ = 90^{\rm o} \end{array} $	PbCrO2, PbCrO4
(c)	Rhombohe dral	a = b = c	$\begin{array}{l} \alpha = \beta = \gamma \\ \neq 90^{\rm o} \end{array}$	CaCO ₃ , HgS
(d)	Triclinic	<i>a</i> = <i>b</i> = <i>c</i>	$\begin{array}{l} \alpha \neq \beta = \gamma \neq \\ 90^{\circ} \end{array}$	K ₂ Cr ₂ O ₇ , CuSO ₄ . 5H ₂ O

- 3. Tetragonal crystal system has the following unit cell dimensions [MP PMT 1993]
 - (a) a = b = c and $\alpha = \beta = \gamma = 90^{\circ}$
 - (b) $a = b \neq c$ and $\alpha = \beta = \gamma = 90^{\circ}$
 - (c) $a \neq b \neq c$ and $\alpha = \beta = \gamma = 90^{\circ}$
 - (d) $a = b \neq c$ and $\alpha = \beta = 90^{\circ}$, $\gamma = 120^{\circ}$
- 4. Rhombic sulphur has the following structure

(a) Open chain

(b) Tetrahedral

	(c) Puckered 6-membered ring
F	(d) Puckered 8-membered ring Space lattice of $C_{0}E_{-}$ is [MP PMT 1002]
5.	Space lattice of Car_2 is [MP FM1 1993]
	(b) Body centred cubic
	(c) Simple cubic
	(d) Hexagonal closed packing
6.	For cubic coordination the value of radius ratio is
	(a) $0.732 - 1.000$ (b) $0.225 - 0.414$
-	(c) $0.000 - 0.225$ (d) $0.414 - 0.732$
/•	different crystal systems [MP PMT 1996; MP PET/PMT 199
	(a) 7 (b) 14
0	(c) 32 (d) 230
8.	Example of unit cell with crystallographic dimensions $a \neq b \neq a$ $a = x = 00^{\circ}$, $a \neq 00^{\circ}$ is LAENC to set
	$a \neq b \neq c, a = \gamma = 90$, $p \neq 90$ is [AFMC 1998]
	(a) Calcule (b) Graphile (c) Rhombic sulphur (d) Monoclinic sulphur
9.	In a face-centered cubic lattice, a unit cell is shared
	equally by how many unit cells [CBSE PMT 2005]
	(a) 8 (b) 4 (d) 6
10.	The maximum radius of sphere that can be fitted in the
101	octahedral hole of cubical closed packing of sphere of
	radius r is (b) a tria
	(a) $0.732 r$ (b) $0.414 r$ (c) $0.225 r$ (d) $0.155 r$
11.	The unit cell of a <i>NaCl</i> lattice
	(a) Is body centred cube (b) Has $3Na^+$ ions
	(c) Has 4 <i>NaCl</i> units (d) Is electrically charged
12.	For tetrahedral coordination number, the radius ratio
	r
	$\frac{r_{c^+}}{r_{-}}$ is [KCET 2000]
	$r_{c^+} = r_{a^-}$ is [KCET 2000] (a) 0.732 -1.000 (b) 0.414 - 0.732
	$\frac{r_{c^+}}{r_{a^-}}$ is [KCET 2000] (a) 0.732 - 1.000 (b) 0.414 - 0.732 (c) 0.225 - 0.414 (d) 0.155 - 0.225
13.	r_{c^+} is [KCET 2000] (a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride
13.	r_{c^+} is [KCET 2000] (a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal [MP PMT 1996]
13.	r_{c^+} is [KCET 2000] (a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal [MP PMT 1996] (a) Face centred cubic (b) Body centred cubic
13.	r_{c^+} is [KCET 2000] (a) 0.732 - 1.000 (b) 0.414 - 0.732 (c) 0.225 - 0.414 (d) 0.155 - 0.225 What type of lattice is found in potassium chloride crystal [MP PMT 1996] (a) Face centred cubic (b) Body centred cubic (c) Simple cubic (d) Simple tetragonal
13. 14.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called
13. 14.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called (a) Space lattice(a) Space lattice(b) Simple lattice
13. 14.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal lattice
13. 14. 15.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal latticeCrystals can be classified into basic crystal habits
13. 14. 15.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal lattice(c) Unit cell(d) Crystal lattice(d) Crystal scan be classified into basic crystal habits [MP PMT 1994](a) 3(b) 7
13. 14. 15.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)Space lattice(b)(c)Unit cell(d)(c)Unit cell(d)(d)Simple lattice(e)14(d)
13. 14. 15. 16.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(d)Simple tetragonalThe three dimensional graph of lattice points which setsthe pattern for the whole lattice is called(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal latticeCrystals can be classified into basic crystal habits[MP PMT 1994](a)3(b)7(c)14(d)4How many molecules are there in the unit cell of sodium
13. 14. 15. 16.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Unit cell(d)(c)Unit cell(d)(c)Unit cell(d)(c)Unit cell(d)(d)3(b)(e)14(d)(f)14(d)(g)2(b)(h)4
13. 14. 15. 16.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)Body centred cubic(c)Simple cubic(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal latticeCrystals can be classified intom basic crystal habits[MP PMT 1994](a)3(b)(a)3(b)7(c)14(d)4How many molecules are there in the unit cell of sodium chloride[MP PMT 1996; MP PET 1997](a)2(b)4(c)6(d)8
13. 14. 15. 16.	r_{c^+} r_{a^-} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(c)Simple cubic(d)(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)Space lattice(b)(c)Unit cell(d)(d)Crystal lattice(c)Unit cell(d)(a)3(b)(b)7(c)14(d)(d)4How many molecules are there in the unit cell of sodium chloride(mP PMT 1996; MP PET 1997](a)2(b)4(c)6(d)8In a crystal, the atoms are located at the position of
13. 14. 15. 16.	r_{a^-} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)Body centred cubic(c)Simple cubic(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal lattice(c)14(d)4How many molecules are there in the unit cell of sodium chloride[MP PMT 1996]; MP PET 1997](a)2(b)4(c)6(d)8In a crystal, the atoms are located at the position of [AMU 1985](a)Maximum P.E.(b)(a)Maximum P.E.(b)
13. 14. 15. 16. 17.	r_{c^+} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)Body centred cubic(c)Simple cubic(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal lattice(c)Unit cell(d)4How many molecules are there in the unit cell of sodium chloride[MP PMT 1996; MP PET 1997](a)2(b)4(c)6(d)8In a crystal, the atoms are located at the position of [AMU 1985][AMU 1985](a)Maximum P.E.(b)Minimum P.E.(c)Zero P.E.(d)Infinite P.E.
 13. 14. 15. 16. 17. 18. 	r_{a^-} is[KCET 2000](a) 0.732 -1.000(b) 0.414 -0.732(c) 0.225 -0.414(d) 0.155 -0.225What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal latticeCrystals can be classified into basic crystal habits[MP PMT 1994](a) 3(b) 7(c) 14(d) 4How many molecules are there in the unit cell of sodium chloride(mp PMT 1996; MP PET 1997](a) 2(b) 4(c) 6(d) 8In a crystal, the atoms are located at the position of [AMU 1985](a) Maximum P.E.(b) Minimum P.E.(c) Zero P.E.(d) Infinite P.E.The total number of lattice arrangements in different urter terms in
13. 14. 15. 16. 17.	r_{a^-} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)Body centred cubic(c)Simple cubic(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal lattice(c)Unit cell(d)4(d)4(d)4How many molecules are there in the unit cell of sodium chloride[MP PMT 1996; MP PET 1997](a)2(b)4(c)6(d)8In a crystal, the atoms are located at the position of [AMU 1985][AMU 1985](a)Maximum P.E.(b)Minimum P.E.(c)Zero P.E.(d)Infinite P.E.The total number of lattice arrangements in different crystal systems is[KCET (Engg.) 2001](a)3(b)7
 13. 14. 15. 16. 17. 18. 	r_{a^-} is[KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a)Face centred cubic(b)Body centred cubic(c)Simple cubic(d)Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a)(a)Space lattice(b)Simple lattice(c)Unit cell(d)Crystal lattice(c)Unit cell(d)4(d)4(d)4How many molecules are there in the unit cell of sodium chloride[MP PMT 1996; MP PET 1997](a)2(b)4(c)6(d)8In a crystal, the atoms are located at the position of LAMU 1985][AMU 1985](a)Maximum P.E.(b)Minimum P.E.(c)Zero P.E.(d)Infinite P.E.The total number of lattice arrangements in different crystal systems is[KCET (Engg.) 2001](a)3(b)7(c)8(d)14
 13. 14. 15. 16. 17. 18. 19. 	r_{a^-} is[KCET 2000](a) 0.732 - 1.000(b) 0.414 - 0.732(c) 0.225 - 0.414(d) 0.155 - 0.225What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal lattice(c) Unit cell(d) 4(a) 3(b) 7(c) 14(d) 4How many molecules are there in the unit cell of sodium chloride(mP PMT 1996; MP PET 1997](a) 2(b) 4(c) 6(d) 8In a crystal, the atoms are located at the position of [AMU 1985] (a) Maximum P.E.(b) Minimum P.E.(c) Zero P.E.(d) Infinite P.E.The total number of lattice arrangements in different crystal systems is [KCET (Engg.) 2001] (a) 3(b) 7(c) 8(d) 14Monoclinic crystal has dimension [DCE 2000]
 13. 14. 15. 16. 17. 18. 19. 	r_{a^-} [KCET 2000] (a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal [MP PMT 1996] (a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal lattice(c) Unit cell(d) 4(d) 3(b) 7(c) 14(d) 4How many molecules are there in the unit cell of sodium chloride(mP PMT 1996; MP PET 1997](a) 2(b) 4(c) 6(d) 8In a crystal, the atoms are located at the position of [AMU 1985] (a) Maximum P.E.(b) Minimum P.E.(c) Zero P.E.(d) Infinite P.E.The total number of lattice arrangements in different crystal systems is [KCET (Engg.) 2001] (a) 3(b) 7(c) 8(d) 14Monoclinic crystal has dimension [DCE 2000] (a) $a \neq b \neq c, \alpha = \gamma = 90^\circ, \beta \neq 90^\circ$
 13. 14. 15. 16. 17. 18. 19. 	r_{a^-} [KCET 2000](a) $0.732 - 1.000$ (b) $0.414 - 0.732$ (c) $0.225 - 0.414$ (d) $0.155 - 0.225$ What type of lattice is found in potassium chloride crystal[MP PMT 1996](a) Face centred cubic(b) Body centred cubic(c) Simple cubic(d) Simple tetragonalThe three dimensional graph of lattice points which sets the pattern for the whole lattice is called(a) Space lattice(b) Simple lattice(c) Unit cell(d) Crystal lattice(c) Unit cell(d) 4How many molecules are there in the unit cell of sodium chloride(m) PMT 1996; MP PET 1997](a) 2(b) 4(c) 6(d) 8In a crystal, the atoms are located at the position of LAMU 1985](a) Maximum P.E.(b) Minimum P.E.(c) Zero P.E.(d) Infinite P.E.The total number of lattice arrangements in different crystal systems is[KCET (Engg.) 2001](a) 3(b) 7(c) 8(d) 14Monoclinic crystal has dimension[DCE 2000](a) $a \neq b \neq c, \alpha = \gamma = 90^\circ, \beta \neq 90^\circ$ (b) $a = b = c, \alpha = \beta = \gamma = 90^\circ$

	(d) $a \neq b \neq c, \alpha \neq \beta \neq \gamma \neq 0$	90°		
20.	20. The low solubility of $BaSO_4$ in water can be attribute			
	to			
		[CBSE PMT 1991]		
	(a) High lattice energy	(b) Dissociation energy		
01	(c) Low lattice energy Brownic lattices are of	(d) Ionic bond		
21.	(a) 8 types	(b) 12 types		
	(c) 14 types	(d) 9 types		
22.	The structure of <i>TlCl</i> is si	milar to <i>CsCl</i> . What would be		
	the radius ratio in TlCl			
98]	(a) 0.155 – 0.225	(b) 0.225 – 0.414		
	(c) $0.414 - 0.732$	(d) 0.732 – 1.000		
23.	Structure similar to zinc bl	ende is found in		
	(a) AgCl	(b) NaCl		
	(c) CuCl	(d) <i>TlCl</i>		
24.	The structure of Na_2O cry	ystal is		
	(a) CsCl type	(b) <i>NaCl</i> type		
	(c) ZnS type	(d) Antifluorite		
25.	Structure of <i>ZnS</i> is			
	(a) Body centred cubic	(b) Face centred cubic		
	(c) Simple cube	(d) Fluorite structure		
26.	The crystal system of	a compound with unit cell		
	dimensions $a = 0.387$, $b =$	= 0.387 and $c = 0.504 nm$ and		
	$\alpha = \beta = 90^{\circ}$ and $\gamma = 120^{\circ}$	is [AIIMS 2004]		
	(a) Cubic	(b) Hexagonal		
	(c) Orthorhombic	(d) Rhombohedral		
27.	The number of tetrahedra	l voids in the unit cell of a face		
	centered cubic lattice of sin	milar atoms is [Kerala PMT 2004]		
	(a) 4	(b) 6		
- 0	(c) 8	(d) 10		
28.	28. An <i>fcc</i> unit cell of aluminium contains the equivalent			
	now many atoms	[DCE 2003]		
	(a) 1 (a) 0	$(\mathbf{D}) = 2$		
_	(0) 3	(d) 4		
	Crystal p	backing		
1.	If Z is the number of	atoms in the unit cell that		
	ABCABC, the	ne number of tetrahedral voids		
	in the unit cell is equal to			
	in the ant con is equal to	[AIIMS 2005]		
	(a) Z	(b) 2 <i>Z</i>		
	(c) $Z/2$	(d) Z/4		
2.	The close packing represen	nts ABC ABC order of		
	(a) Body centred cubic pa	cking		
	(b) Face centred cubic pac	cking		
	(c) Simple cubic packing	0		
	(d) Hexagonal cubic close	d packing		
3.	The arrangement ABCAB	CABC is referred as		
-	-	[MP PET 2001]		
	(a) Octahedral close pack	ing (b)Hexagonal close packing		
	(c) Tetragonal close packi	ng (d) Cubic close packing		
4.	The number of close neigh	hbour in a body-centred cubic		
	lattice of identical sphere i	S [MP PET 2001]		
	(a) 8	(b) 6		
	(c) 4	(d) 2		
5۰	The number of equidistan	t oppositely charged ions in a		
	sodium chloride crystal is	[MP PET 2001]		
	(a) 8	(D) 6		

(c) 4

(d) 2

6.	Na and Mg crystallize in BC	C and FCC type crystals
	respectively, then the number o	f atoms of Na and Mg
	present in the unit cell of their re	espective crystal is
		[AIEEE 2002] 2
	(a) 4 and 2 (b)	9 and 14
	(c) 14 and 9 (d)	2 and 4
7•	An AB_2 type structure is found	in [AIIMS 2002]
	(a) <i>NaCl</i> (b)	Al_2O_3
	(c) CaF_2 (d)	N ₂ O
8.	Potassium crystallizes with a	- [MP PET/PMT 1008]
0.	(a) Face-centred cubic lattice	
	(b) Body-centred cubic lattice	2
	(c) Simple cubic lattice	
	(d) Orthorhombic lattice	
9.	If the number of atoms per un	nit in a crystal is 2, the
	structure of crystal is	2
	(a) Octahedral	
	(b) Body centred cubic <i>bcc</i>	
	(c) Face centred cubic fcc	
10	(d) simple cubic	2 1. amutallizaz in auhio
10.	lattice in which both lithium and	silver have coordination
	number of eight. The crystal clas	s is
	0 2	[CBSE PMT 1997]
	(a) Simple cube (b)	Body-centred cube
	(c) Face-centred cube (d)	None of these
11.	The number of octahedral sit	es per sphere in a fcc
	structure is	[MP PMT 2000, 01]
	(a) 8 (b)	4
19	(C) 2 (C) Heveronal close packed arrange	1 ment of ions is described
	as	inclife of folio is described
		[MP PMT 1994]
	(a) ABCABA (b)	ABC ABC
	(c) ABABA (d)	ABBAB
13.	An example of a body cube is	[AIIMS 1996] Magnagium
	$(a) \text{Southin} \qquad (b) \\ (c) \text{Zinc} \qquad (d)$	Copper 1
14.	An example of fluorite structure	is
-4.	(a) <i>NaF</i> (b)	SrF ₂
	$(c) AlCl \qquad (d)$	SiF
	$(\mathbf{u}) \mathbf{u} = \mathbf{u} = \mathbf{u} = \mathbf{u} = \mathbf{u}$	ola alternate tetral altern
15.	voids are occupied?	ais alternate tetraneoral 2
	(a) <i>NaCl</i> (b)	ZnS
	(c) CaF_2 (d)	Na ₂ O
16.	Which of the following contains	rock salt structure
	(a) SrF_2 (b)	MgO 3
	(c) Al_2O_3 (d)	All
17.	In the fluorite structure, the	coordination number of
• / •	Ca^{2+} ion is	
	(a) 4 (b)	6 4
	(c) 8 (d)	3
18.	The ratio of close-packed atom	s to tetrahedral holes in
	cubic close packing is	[Pb. PMT 1998]
	(a) 1.1 (D)	1,2

19. A solid is made of two elements *X* and *Z*. The atoms *Z* are in CCP arrangement while the atom *X* occupy all the tetrahedral sites. What is the formula of the compound

(d) 2:1

(c) 1:3

(a) *XZ* (b) XZ_2 (c) X_2Z (d) X_2Z_3 An ionic compound has a unit cell consisting of A ions at D. the corners of a cube and B ions on the centres of the faces of the cube. The empirical formula for this [CBSE PMT 2004; AIEEE 2005] compound would be (a) *AB* (b) A₂B (c) AB_3 (d) A_3B The vacant space in the *bcc* unit cell is (a) 32% (b) 23% (c) 26% (d) None of these The number of octahedral voids in a unit cell of a cubical 2. closest packed structure is (a) 1 (b) 2 (c) 4 (d) 8 In the closest packed structure of a metallic lattice, the 2. number of nearest neighbours of a metallic atom is [JIPMER 2002] (a) Twelve (b) Four (d) Six (c) Eight In the rock salt structure, the number of formula units 4. per unit cell is equal to (a) 1 (b) 2 (d) 4 (c) 3 Hexagonal close packing is found in crystal lattice of 5. [MH CET 2002] (a) Na (b) Mg(c) Al (d) None of these Which ion has the largest radius from the following ions (b) Mg^{2+} (a) Na^+ (c) Al^{3+} (d) Si⁴⁺

[UPSEAT 2004]

Mathematical analysis of cubic system and Bragg's equation

1. The formula for determination of density of unit cell is

(a)
$$\frac{a^3 \times N_o}{N \times M} g \, cm^{-3}$$
 (b) $\frac{N \times M}{a^3 \times N_o} g \, cm^{-3}$
(c) $\frac{a^3 \times M}{N \times N_o} g \, cm^{-3}$ (d) $\frac{M \times N_o}{a^3 \times N} g \, cm^{-3}$

Potassium fluoride has *NaCl* type structure. What is the distance between K^+ and F^- ions if cell edge is *a cm*

- (a) $2a \ cm$ (b) $a/2 \ cm$
- (c) $4a \ cm$ (d) $a/4 \ cm$
- **3.** An element occurring in the *bcc* structure has 12.08×10^{23} unit cells. The total number of atoms of the element in these cells will be [MP PET 1994] (a) 24.16×10^{23} (b) 36.18×10^{23}
 - (c) 6.04×10^{23} (d) 12.08×10^{23}
 - If an atom is present in the centre of the cube, the
- participation of that atom per unit cell is

(a)	$\frac{1}{4}$	(b) 1
(c)	$\frac{1}{2}$	(d) $\frac{1}{8}$

5. For an ionic crystal of the general formula *AX* and coordination number 6, the value of radius ratio will be **[MP PMT 1993]**

- (a) Greater than 0.73
- (b) In between 0.73 and 0.41
- (c) In between 0.41 and 0.22
- (d) Less than 0.22
- 6. The number of spheres contained (i) in one body centred cubic unit cell and (ii) in one face centred cubic unit cell, is
 - (a) In (i) 2 and in (ii) 4 (b) In (i) 3 and in (ii) 2
 - (c) In (i) 4 and in (ii) 2 (d) In (i) 2 and in (ii) 3
- 7. *CsBr* crystal has *bcc* structure. It has an edge length of 4.3 Å. The shortest interionic distance between Cs^+

and Br^- ions is

(a) 1.86 \mathring{A} (b) 3.72 \mathring{A}

- (c) 4.3 Å (d) 7.44 Å
- 8. In octahedral holes (voids)
 - (a) A simple triangular void surrounded by four spheres
 - (b) A bi-triangular void surrounded by four spheres
 - (c) A bi-triangular void surrounded by six spheres
 - (d) A bi-triangular void surrounded by eight spheres
- 9. Bragg's law is given by the equation [MP PMT 1995, 2002]

(a)
$$n\lambda = 2\theta \sin \theta$$
 (b) $n\lambda = 2d \sin \theta$
(c) $2n\lambda = d \sin \theta$ (d) $n\frac{\theta}{2} = \frac{d}{2}\sin \theta$

10. The number of atoms in 100 g of an fcc crystal with

- density $d = 10 g/cm^3$ and cell edge equal to 100 pm, is equal to [CBSE PMT 1994; KCET 2002]
- (a) 4×10^{25} (b) 3×10^{25}
- (c) 2×10^{25} (d) 1×10^{25}
- In the crystals of which of the following ionic compounds would you expect maximum distance between centres of cations and anions [CBSE PMT 1998]

(a)	LiF	(b)	CsF
(c)	CsI	(d)	LiI

12. The number of unit cells in 58.5 g of *NaCl* is nearly

[MP PMT 2000, 01]

[IIT 1995]

(a)	6×10^{20}	(b)	3×10^{22}
(c)	1.5×10^{23}	(d)	0.5×10^{24}

- **13.** How many unit cells are present in a cube-shaped ideal
crystal of NaCl of mass 1.00 g [Atomic masses:
Na = 23, Cl = 35.5][AIEEE 2003]
- (a) 2.57×10^{21} unit cells (b) 5.14×10^{21} unit cells (c) 1.28×10^{21} unit cells (d) 1.71×10^{21} unit cells **14.** In the Bragg's equation for diffraction of *X*-rays, *n* represents for [MP PMT 2000] (a) Quantum number (b) An integer
 - (c) Avogadro's numbers (d) Moles
- **15.** In a face centred cubic cell, an atom at the face contributes to the unit cell

	[Karna	taka (Engg./Med.) 2000; AFMC 2001]
(a)	1/4 part	(b) 1/8 part

- (a) 1/4 part (b) 1/8 part (c) 1 part (d) 1/2 part
- 16. The interionic distance for cesium chloride crystal will be [MP PET 2002]

- (c) $\frac{\sqrt{3}a}{2}$ (d) $\frac{2a}{\sqrt{3}}$
- **17.** Sodium metal crystallizes as a body centred cubic lattice with the cell edge 4.29 Å. What is the radius of sodium atom

[AIIMS 1999]

[AIIMS 1991]

(a)	$1.857 \times 10^{-8} cm$	(b)	2.371×10^{-7} cm
(c)	$3.817 \times 10^{-8} cm$	(d)	$9.312 \times 10^{-7} cm$

- 18. For an ionic crystal of the type *AB*, the value of (limiting) radius ratio is 0.40. The value suggests that the crystal structure should be(a) Octahedral(b) Tetrahedral
 - (c) Square planar (d) Plane triangle
- **19.** Potassium has a bcc structure with nearest neighbour distance 4.52 Å. Its atomic weight is 39. Its density (in
 - $kg m^{-3}$) will be (a) 454 (b) 804
 - (c) 852 (d) 908
- **20.** If the value of ionic radius ratio $\left(\frac{r_c}{r_a}\right)$ is 0.52 in an ionic

compound, the geometrical arrangement of ions in crystal is

- (a) Tetrahedral (b) Planar
- (c) Octahedral (d) Pyramidal
- 21. The number of atoms/molecules contained in one face centred cubic unit cell of a monoatomic substance is [CPMT 1989, 94; CBSE PMT 1989, 96; NCERT 1990;
 - MP PET 1993; KCET 1999]
 - (a) 1 (b) 2
 - (c) 4 (d) 6
- **22.** The number of atoms/molecules contained in one body centered cubic unit cell is
 - (a) 1 (b) 2 (c) 4 (d) 6
- (c) 4 (d) 6 **23.** It the distance between Na^+ and Cl^- ions in sodium
 - chloride crystal is *X* pm, the length of the edge of the unit cell is **[KCET 2004]**
 - (a) 4X pm (b) X/4 pm
- (c) X/2 pm
 (d) 2X pm
 24. The edge of unit cell of FCC Xe crystal is 620 pm. The radius of Xe atom is [MP PET 2004]
 (a) 219.25 Pm
 (b) 235.16 Pm
 - (c) 189.37 Pm (d) 209.87 Pm
- **25.** In orthorhombic, the value of *a*, *b* and *c* are respectively $4.2\mathring{A}, 8.6\mathring{A}$ and $8.3\mathring{A}$. given the molecular mass of the solute is 155 gm mol⁻¹ and that of density is 3.3 gm/cc,
 - the number of formula units per unit cell is
 - [Orrisa JEE 2005]
 - (a) 2 (b) 3 (c) 4 (d) 6
- **26.** A metal has bcc structure and the edge length of its unit cell is 3.04. The volume of the unit cell in cm^3 will be **[Orrisa JEE 2005]**
 - (a) $1.6 \times 10^{21} cm^3$ (b) $2.81 \times 10^{-23} cm^3$
 - (c) $6.02 \times 10^{-23} cm^3$ (d) $6.6 \times 10^{-24} cm^3$
- 27. In face centred cubic unit cell edge length is [DPMT 2005]

(a)
$$\frac{4}{\sqrt{3}}r$$
 (b) $\frac{4}{\sqrt{2}}r$

(b) $\frac{a}{2}$

(a) a

(c)
$$2r$$
 (d) $\frac{\sqrt{3}}{2}r$

Crystal structure and Coordination number

A solid has a structure in which 'W' atoms are located at 1. the corners of a cubic lattice 'O' atoms at the centre of edges and 'Na' atoms at the centre of the cube. The formula for the compound is [KCET 1996] (a) $NaWO_{2}$ (b) $NaWO_{2}$

(c)	Na_2WO_3	(d)	$NaWO_{4}$

Potassium crystallizes in a bcc lattice, hence the 2. coordination number of potassium in potassium metal is [KCEE 1993]

(a) o	(b) 4
(c) 6	(d) 8

Body centered cubic lattice has a coordination number of 3. [AIIMS 1996: MP PMT 2002] \sim

(a)	4	(D) 8
(c)	12	(d) 6

A compound is formed by elements A and B. This 4. crystallizes in the cubic structure when atoms A are the corners of the cube and atoms *B* are at the centre of the body. The simplest formula of the compounds is

[KCET 1993; CBSE PMT 2000; Kerala PMT 2002]

- (a) *AB* (b) AB_2 (c) A_2B (d) AB_4
- Coordination number for *Cu* is [AMU 1982] 5. (a) 1 (b) 6
 - (d) 12 (c) 8
- In the crystal of CsCl, the nearest neighbours of each 6. Cs ion are [MP PET 1993]
- In a cubic structure of compound which is made from X7. and Y, where X atoms are at the corners of the cube and *Y* at the face centres of the cube. The molecular formula of the compound is [AIIMS 2000] (a) $X_{2}Y$ (b) $X_{3}Y$
 - (c) XY_2 (d) XY_3
- Ferrous oxide has a cubic structure and each edge of the 8. unit cell is 5.0 Å. Assuming density of the oxide as $4.0g - cm^{-3}$, then the number of Fe^{2+} and O^{2-} ions present in each unit cell will be [MP PET 2000]
 - (a) Four Fe^{2+} and four O^{2-}
 - (b) Two Fe^{2+} and four O^{2-}
 - (c) Four Fe^{2+} and two O^{2-}
 - (d) Three Fe^{2+} and three O^{2-}
- Which of the following statements is not true about 9. NaCl structure [DCE 2001]
 - (a) Cl^{-} ions are in fcc arrangement
 - (b) Na^+ ions has coordination number 4
 - (c) Cl^{-} ions has coordination number 6
 - (d) Each unit cell contains 4 NaCl molecules
- In *CsCl* structure, the coordination number of Cs^+ is 10. [MP PMT 2001]
 - (a) Equal to that of Cl^{-} , that is 6
 - (b) Equal to that of Cl^{-} , that is 8

(c) Not equal to that of Cl^{-} , that is 6

(d) Not equal to that of Cl^{-} , that is 8

- In a solid 'AB' having the NaCl structure, 'A' atoms 11. occupy the corners of the cubic unit cell. If all the facecentered atoms along one of the axes are removed, then the resultant stoichiometry of the solid is [IIT Screening 2001] (a) AB_{2} (b) A_2B
 - (c) $A_4 B_3$ (d) $A_3 B_4$
- 12. In solid *CsCl* each *Cl* is closely packed with how many Cs [MP PET 2003]
 - (a) 8 (b) 6 (c) 10 (d) 2
- In A^+B^- ionic compound, radii of A^+ and B^- ions are 13. 180 pm and 187 pm respectively. The crystal structure of this compound will be
 - (a) NaCl type (b) CsCl type
 - (d) Similar to diamond (c) ZnS type
- In which of the following substances the carbon atom is 14. arranged in a regular tetrahedral structure [NCERT 1978] (a) Diamond (b) Benzene
 - (c) Graphite (d) Carbon black
- The coordination number of a metal crystallizing in a 15. hexagonal close packed structure is
 - [NCERT 1978; IIT 1999] (a) 4 (b) 12
 - (c) 8 (d) 6
- The structure of M_gO is similar to NaCl. What would 16. be the coordination number of magnesium
 - (a) 2 (b) 4 (c) 6 (d) 8

How many chloride ions are there around sodium ion in 17. sodium chloride crystal [NCERT 1979, 80; CPMT 1988; BHU 1982, 87; MP PET 1995, 99]

- (a) 3 (b) 8
- (d) 6 (c) 4
- 18. Most crystals show good cleavage because their atoms, ions or molecules are [CBSE PMT 1991]
 - (a) Weakly bonded together
 - (b) Strongly bonded together
 - (c) Spherically symmetrical
 - (d) Arranged in planes
- An example of a non-stoichiometric compound is 19.

(a)
$$Al_2O_3$$
 (b) Fe_3O_4

(c)
$$NiO_2$$
 (d) PbO

- If the radius ratio is in the range of 0.731 1, then the 20. coordination number will be
 - (a) 2 (b) 4 (c) 6 (d) 8
- If the radius ratio is in the range of 0.414 0.732, then 21. the coordination number will be
 - (a) 2 (b) 4 (c) 6 (d) 8
- What is the coordination number of sodium in Na_2O 22.

[AIIMS 2003]

- (a) 6 (b) 4 (c) 8 (d) 2
- The ratio of cationic radius to anionic radius in an ionic 23. crystal is greater than 0.732. Its coordination number is

[KCET 2003]

- (a) Six chloride ions (b) Eight chloride ions (d) Eight Cs ions (c) Six Cs ions

	(a) 6	(b) 8	
	(c) 1	(d) 4	
24 .	In <i>CsCl</i> lattice the coordinate	ation number of Cs^+ ion is	F
	(a) 2	(b) 4 (d) 10	1
07	(c) δ	(U) 12	4.
25.	(a) for	(b) <i>bcc</i>	-
	(a) $f(t)$	(d) None	
06	(c) Dotti (a) and (b) In $N_{-}CL$ lattice the coordin	(u) None Cl^{-} ion is	
20.	(a) 2	(b) a	5.
	(a) <u>2</u> (c) 6	(d) 8	
27.	In zinc blende structure t	he coordination number of	
	Zn^{2+} ion is		
	(a) 2	(b) 4	6.
	(c) 6	(d) 8	
28.	Coordination number of Na	⁺ ion in rock salt is	
	(-) 10	[BVP 2004]	
	(a) 12	(b) 4 (d) 6	
20	The number of Cl^- ions	around one Na^+ in $NaCl$	
29.	crystal lattice is	[MP PET 1996; BVP 2004]	7
	(a) 12	(b) 4	/•
	(c) 8	(d) 6	
30.	The number of atoms j	present in unit cell of a	004]
	(a) 6	(b) 3	5041
	(c) 2	(d) 1	
31.	The coordination number	of a metal crystallizing in a	
	hexagonal close packed chep	b structure is [MP PMT 2004]	0
	(a) 12 (c) 4	(d) 6	0.
32.	Which of the following state	ment(s) is(are) correct	
-	() m	[IIT 1998]	
	(a) The coordination num $CsCl$ crystal is 8	ber of each type of ion in	9.
	(b) A metal that crystalliz	zes in <i>bcc</i> structure has a	
	(c) A unit cell of an ionic c	rvstal shares some of its ions	10.
	with other unit cells	- J	
	(d) The length of the unit	t cell in <i>NaCl</i> is 552 pm	
	$(r_{Na^+} = 95 \ pm; \ r_{Cl^-} = 18)$	31 pm)	
33.	The co-ordination number of	of Na^+ in $NaCl$ is	
		[Orrisa JEE 2005]	
	(a) 6	(b) 8 (d) 1	
34.	In the calcium fluoride	structure the co-ordination	11.
01	number of the cation and an	ions are respectively [J & K 200	5]
	(a) 6, 6	(b) 8,4	
	(c) 4, 4	(d) 4,8	
	Defects in	crystal	
1.	Certain crystals produce ele	ctric signals on application of	12.
	(a) Pyroelectricity	(b) Ferroelectricity	
	(c) Peizoelectricity	(d) Ferrielectricity	
2.	Which defect causes decreas	e in the density of crystal	
		[KCET 2000, 05]	
	(a) Frenkel	(b) Schottky	13.
	(c) Interstitial	(d) F – centre	
3.	The correct statement regard	ding F – centre is	

- (a) Electron are held in the voids of crystals
- (b) F centre produces colour to the crystals
- (c) Conductivity of the crystal increases due to F-centre

(d) All

13.

Doping of silicon (Si) with boron (B) leads to

- [UPSEAT 2004] (a) n-type semiconductor (b) p-type semiconductor
- (c) Metal (d) Insulator
- If *NaCl* is doped with $10^{-3}mol \% SrCl_2$, then the 5. concentration of cation vacancies will be
 - (a) $1 \times 10^{-3} mol\%$ (b) $2 \times 10^{-3} mol\%$
 - (c) $3 \times 10^{-3} mol\%$ (d) $4 \times 10^{-3} mol\%$
- In the laboratory, sodium chloride is made by burning 6. the sodium in the atmosphere of chlorine which is yellow in colour. The cause of vellow colour is
 - (a) Presence of Na^+ ions in the crystal lattice
 - (b) Presence of Cl^{-} ions in the crystal lattice
 - (c) Presence of electron in the crystal lattice
 - (d) Presence of face centered cubic crystal lattice
 - Frenkel defect is caused due to [MP PET 1994]
 - (a) An ion missing from the normal lattice site creating a vacancy
 - (b) An extra positive ion occupying an interstitial position in the lattice
 - (c) An extra negative ion occupying an interstitial position in the lattice
 - (d) The shift of a positive ion from its normal lattice site to an interstitial site
- 8. Which one of the following has Frenkel defect

[MP PMT 2000]

- (a) Sodium chloride (b) Graphite
- (c) Silver bromide (d) Diamond
- Schottky defect generally appears in 9. [DCE 2004]
 - (a) NaCl (b) *KCl*
 - (c) CsCl (d) All of these
- Schottky defect in crystals is observed when 10.

[CBSE PMT 1998; KCET 2002]

- (a) Density of crystal is increased
- (b) Unequal number of cations and anions are missing from the lattice
- (c) An ion leaves its normal site and occupies an interstitial site
- (d) Equal number of cations and anions are missing from the lattice
- Ionic solids, with Schottky defects, contain in their structure

[CBSE PMT 1994]

- (a) Equal number of cation and anion vacancies
- (b) Anion vacancies and interstitial anions
- (c) Cation vacancies only
- (d) Cation vacancies and interstitial cations
- The following is not a function of an impurity present in 12. a crystal [MP PET 1995]
 - (a) Establishing thermal equilibrium
 - (b) Having tendency to diffuse
 - (c) Contributing to scattering
 - (d) Introducing new electronic energy levels
 - Due to Frenkel defect, the density of ionic solids

[MP PET 1996; MP PMT 2002]

(b) Decreases

- (a) Increases
- (c) Does not change (d) Changes

Point defects are present in [MP PMT 1997]		(a) Hexagonal (b) Orthorhombic
(a) Ionic solids (b) Molecular solids		(c) Cubic (d) Triclinic
(c) Amorphous solids (d) Liquids	5۰	Which of the following molecules has three-fold axis of
If a non-metal is added to the interstitial sites of a metal		symmetry [UPSEAT 2004]
then the metal becomes [DCE 2001]		(a) NH_3 (b) C_2H_4
(a) Softer (b) Less tensile		(c) CO_2 (d) SO_2
(c) Less malleable (d) More ductile	6.	Which one possess a antifluorite structure
In AgBr crystal, the ion size lies in the order		(a) Na_2O (b) MgO
$Ag^+ \ll Br^-$. The $AgBr$ crystal should have the		(c) Fe_2O_2 (d) Al_2O_2
following characteristics	-	Which one of the following is the biggest ion IMP DET 1000
(a) Defectless (perfect) crystal	/•	(-) A^{1+3} (-) B^{-+2}
(b) Schottky defect only		(a) Al^{-1} (b) Ba^{-1}
(c) Frenkel defect only		(c) Mg^{+2} (d) Na^+
(d) Both Schottky and Frenkel defects	8.	The edge length of face centred unit cubic cell is
Frenkel and Schottky defects are [BHU 2003]		508 pm . If the radius of the cation is 110 pm , the
(a) Nucleus defects (b) Non-crystal defects		radius of the anion is
(c) Crystal defects (d) None of these		[CBSE PMT 1998]
(a) Broom in an interstitical allocated billion stability of the		(a) 285 pm (b) 398 pm
(a) Brass is an interstitual alloy, while steel is a substitutional alloy		(c) 144 <i>pm</i> (d) 618 <i>pm</i>
(b) Brass is a substitutional alloy, while steel is an	9.	An element (atomic mass $100 g/mol$) having bcc
interstitial alloy		structure has unit cell edge 400 pm. Then density of the
(c) Brass and steel are both substitutional alloys		element is
(d) Brass and steel are both interstitial alloys		[CBSE PMT 1996; AIIMS 2002]
The flame colours of metal ions are due to [KCET 2003]		(a) 10.376 g/cm^3 (b) 5.188 g/cm^3
(a) Frenkel defect (b) Schottky defect		(c) 7.289 g/cm^3 (d) 2.144 g/cm^3
(c) Metal deficiency defect (d) Metal excess defect	10.	If the pressure on a $NaCl$ structure is increased, then
Which one of the following crystals does not exhibit	10.	its coordination number will [AFMC 2000]
$\begin{array}{c} \text{Frenkei defect} \\ \text{(b)} AaBr \\ \text{(b)} AaCl \end{array}$		(a) Increase (b) Decrease
(a) AyD (b) AyC (c) KBr (d) ZnS		(c) Remain the same (d) Either (b) or (c)
In a solid lattice the cation has left a lattice site and is	11.	The pyknometric density of sodium chloride crystal is
located at an interstitial position, the lattice defect is		$2.165 \times 10^{3} kg m^{-3}$ while its <i>X</i> -rays density is
[AIIMS 1982, 1991; DCE 2002; J & K 2005]		2 178 $\times 10^{3} kgm^{-3}$ The fraction of unoccupied sites in
(a) Interstitial defect (b) Valency defect		sodium chlorida crystal is
(c) Frenkel defect (d) Schottky defect		Solution control crystaris [CBSE F M1 2003]
When electrons are trapped into the crystal in anion		(a) 5.96×10 (b) 5.96
(a) Schotky defect (b) Frenkel defect		(c) 5.96×10^{-2} (d) 5.96×10^{-1}
(c) Stoichiometric defect (d) F-centres	12.	Which of the following statements is correct for C_sBr_3
Schottky defect defines imperfection in the lattice		[IIT 1996]
structure of a [AIIMS 2002]		(a) It is a covalent compound $(a) = \frac{1}{2} \left(\frac{1}{2} - \frac{1}{2} \right)^2$
(a) Solid (b) Liquid		(b) It contains Cs^{3+} and Br^{-1} ions
(c) Gas (d) Plasma		(c) It contains Cs^+ and Br_3^- ions
		(d) It contains Cs^+ , Br^- and lattice Br_2 molecule
Critical Thinking	12	In which compound 8 : 8 coordination is found
	13.	[EAMCET 1984]
		(a) $CsCl$ (b) MgO
- Objective Questions		(c) Al_2O_3 (d) All of these
Amorphous solids are	14.	If the coordination of Ca^{2+} in CaE_{-} is 8 then the
(a) Solid substance in real sense		
(b) Liquid in real sense		coordination number of F ion would be
(c) Supercooled liquid		(a) 3 (b) 4 (d) 8
(d) Substance with definite melting point	15	(u) o For some crystals, the radius ratio for cation and anion is
Silicon is found in nature in the form of [MH CET 2002]	19.	0.525, its coordination number will be
(a) Body centered cubic structure		(a) 2 (b) 4
(b) Hexagonal close-packed structure		(c) 6 (d) 8
(c) Network solid	16.	The basic building unit of all silicates is [UPSEAT 2002]
(d) Face centered cubic structure		(a) SiO_4 square planar (b) $[SiO_4]^{4-}$ tetrahedron
A match box exhibits [MP PET 1993, 95]		(c) SiQ, octahedron (d) SiQ, linear
(a) Cubic geometry (b) Monoclinic geometry	17	What type of grystal defect is indicated in the diagram
(c) Orthorhombic geometry (d) Tetragonal geometry	1'/•	below
which has no rotation of symmetry [Orrisa JEE 2004]		

14.

15.

16.

17.

18.

19.

20.

21.

22.

23.

1.

2.

3.

4.

8

[MHCET 2003]

(b) Spodumene

(d) None of these

 $Na^{+} Cl^{-} Na^{+} Cl^{-} Na^{+} Cl^{-}$ $Cl^{-} \Box Cl^{-} Na^{+} \Box Na^{+}$ $Na^{+} Cl^{-} \Box Cl^{-} Na^{+} Cl^{-}$ $Cl^{-} Na^{+} Cl^{-} Na^{+} \Box Na^{+}$ (a) Interstitial defect
(b) Schottky defect
(c) Frenkel defect
(d) Frenkel and Schottky defects

- Which of the following is a three dimensional silicate
 - 0
 - (a) Mica(c) Zeolite
 - (c) 2con
 - (e) 12

18.

Assertion & Reason

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.

1.	Assertion	:	Diamond is a precious stone.
	Reason	:	Carbon atoms are tetrahedrally arranged in diamond. [AIIMS 1994]
2.	Assertion	:	In crystal lattice, the size of the cation is larger in a tetrahedral hole than in an octahedral hole.
	Reason	:	The cations occupy more space than anions in crystal packing. [AIIMS 1996]
3.	Assertion	:	Crystalline solids have short range order.
	Reason order.	:	Amorphous solids have long range
			[AIIMS 1999]
4.	Assertion	:	In any ionic solid (MX) with Schottky defects, the number of positive and negative ions are same.
	Reason	:	Equal number of cation and anion vacancies are present.
			[IIT Screening 2001]
5.	Assertion	:	Space or crystal lattice differ in symmetry of the arrangement of points.
	Reason	:	$n\lambda = 2d \sin \theta$, is known as Bragg's equation.
6.	Assertion	:	In close packing of spheres, a tetrahedral void is surrounded by four spheres whereas an octahedral void is surrounded by six spheres.
	Reason	:	A tetrahedral void has a tetrahedral shape whereas an octahedral void has an octahedral shape.
7.	Assertion	:	Cyclic silicates and chain silicates have

the same general molecular formula.

			SiO_4 tetrahedron are shared while in
			chain silicates only two are shared with other tetrahedra.
8.	Assertion	:	The presence of a large number of Schottky defects in <i>NaCl</i> lowers its density.
	Reason	:	In <i>NaCl</i> , there are approximately 10^6 Schottky pairs per cm^3 at room temperature.
9.	Assertion	:	Anion vacancies in alkali halides are produced by heating the alkali halide crystals with alkali metal vapour.
	Reason	:	Electrons trapped in anion vacancies are referred to as <i>F</i> -centres.
10.	Assertion	:	Electrical conductivity of semiconductors increases with increasing temperature.
	Reason	:	With increase in temperature, large number of electrons from the valence band can jump to the conduction band.
11.	Assertion	:	On heating ferromagnetic or ferrimagnetic substances they become

In cyclic silicates, three corners of each

- paramagnetic. Reason : The electrons change their spin on
- heating. **12.** Assertion : Lead zirconate is a piezoelectric crystal. Reason : Lead zirconate crystals have no dipole
 - on : Lead zirconate crystals have no dipole moment.

Answers

Type of solid and Their properties

1	а	2	b	3	а	4	а	5	b
6	С	7	С	8	b	9	d	10	d
11	b	12	а	13	C	14	С	15	a
16	а	17	a	18	d	19	c	20	c
21	b	22	d	23	d	24	d	25	a
26	d	27	а	28	а	29	d	30	d
31	d	32	а	33	C	34	а	35	b
36	а	37	a	38	b	39	C	40	ac

Crystallography and Lattice

1	b	2	C	3	b	4	d	5	a
6	а	7	b	8	d	9	d	10	b
11	C	12	C	13	а	14	C	15	b
16	b	17	b	18	b	19	а	20	a
21	C	22	d	23	C	24	d	25	b
26	b	27	С	28	d				

Crystal packing

1	b	2	b	3	d	4	а	5	b
6	d	7	С	8	b	9	b	10	b

11	d	12	C	13	a	14	b	15	b
16	b	17	C	18	b	19	C	20	C
21	а	22	C	23	а	24	d	25	b
26	а								

Mathematical analysis of cubic system and Bragg's equation

1	b	2	b	3	а	4	b	5	b
6	а	7	b	8	С	9	b	10	а
11	С	12	с	13	а	14	b	15	d
16	С	17	а	18	b	19	d	20	C
21	С	22	b	23	d	24	а	25	C
26	b	27	b						

Crystal structure and Coordination number

1	b	2	d	3	b	4	a	5	d
6	b	7	d	8	а	9	b	10	b
11	d	12	а	13	b	14	a	15	b
16	C	17	d	18	d	19	b	20	d
21	С	22	b	23	b	24	С	25	а
26	С	27	b	28	d	29	d	30	d
31	а	32	acd	33	a	34	b		

Defects in crystal

1	с	2	b	3	d	4	d	5	а
6	С	7	d	8	C	9	d	10	d
11	а	12	a	13	C	14	а	15	b
16	C	17	С	18	c	19	d	20	C
21	C	22	d	23	a				

Critical Thinking Questions

1	c	2	c	3	C	4	d	5	a
6	а	7	b	8	С	9	b	10	а
11	а	12	С	13	а	14	b	15	С
16	b	17	b	18	С				

Assertion & Reason

1	b	2	d	3	d	4	а	5	b
6	с	7	C	8	b	9	b	10	а
11	а	12	C						



Properties and Types of solids

- (a) Both gases and liquids possess fluidity and hence viscosity molecules in the solid state do not have translational motion.
- **2.** (b) It is a characteristic of liquid crystal.
- **3.** (a) $BaTiO_3$ is a ferroelectric compound.
- 5. (b) The value of heat of fusion of *NaCl* is very high due to *fcc* arrangement of its ions.
- **6.** (c) Piezoelectric crystals are used in record player.
- 8. (b) NaCl is a ionic solid in which constituent particles are positive (Na^+) and negative (Cl^-) ions.
- (d) Amorphous solids have short range order but no sharp in melting point.
- 10. (d) Solids have definite shape, size and rigidity.
- 12. (a) In crystalline solid there is perfect arrangement of the constituent particles only at OK. As the temperature increases the chance that a lattice site may be unoccupied by an ion increases. As the number of defects increases with temperature solid change in liquid.
- **13.** (c) Diamond is a covalent solid in which constituent particles are atoms.
- **14.** (c) Solid *NaCl* is a bad conductor of electricity because ions are not free to move.
- **15.** (a) The existence of a substance in more than one crystalline form is known as polymorphism.
- **16.** (a) Solids are also non-crystalline in nature.
- 17. (a) Ice has the lowest melting point out of the given solids, hence it has the weakest intermolecular forces.
- **19.** (c) All metals and some alloys are metallic crystal.
- 20. (c) Iodine crystals are molecular crystals, in which constituent particles are molecules having interparticle forces are Vander Waal's forces.
- **21.** (b) Ionic solids have highest melting point due to strong electrostatic forces of attraction.
- **22.** (d) For *n*-type, impurity added to silicon should have more than 4 valence electrons.
- 23. (d) Glass is an amorphous solid.
- **25.** (a) Crystalline solids have regular arrangement of constituent particles, sharp melting points and are anisotropic.

- **26.** (d) Sugar is a crystalline solid while glass, rubber and plastic are amorphous solids.
- **28.** (a) MnO_2 is antiferromagnetic.
- **29.** (d) Graphite is sp^2 hybridised and a covalent crystal.
- **30.** (d) Ionic crystals exhibit non-directional properties of the bond.
- **31.** (d) Ice is a molecular crystal in which the constituent units are molecules and the interparticle forces are hydrogen bonds.
- **32.** (a) Quartz is a covalent crystal having a framework of silicates or silica, *i.e.* a three dimensional network when all the four oxygen atoms of each of SiO_4 tetrahedron are shared.
- **33.** (c) Metallic crystals are good conductor of heat and current due to free electrons in them.
- **34.** (a) Silicon is a covalent crystal in which constituent particles are atoms.
- 35. (b) *LiF* is an example of ionic crystal solid, in which constituent particles are positive (*Li*⁺) and negative (*F*⁻) ions.
- **36.** (a) Amorphous solids neither have ordered arrangement (*i.e.* no definite shape) nor have sharp melting point like crystals, but when heated, they become pliable until they assume the properties usually related to liquids. It is therefore they are regarded as super-cooled liquids.
- 37. (a) Silicon is a semiconductor because it is a thermal active and its conductivity increased with increasing temperature.
- 38. (b) Amorphous solids are isotropic, because of these substances show same properties in all directions.
- **39.** (c) Polymorphism is a ability of a substances which show two or more crystalline structure
- **40.** (ac) Amorphous solids neither have ordered arrangement (*i.e.* no definite shape) nor have sharp melting point like crystals, but when heated, they become pliable until they assume the properties usually related to liquids. It is therefore they are regarded as super-cooled liquids.

Crystallography and Lattice

- 1. (b) A crystal system is hexagonal if its unit cell having $a = b \neq c$ axial ratio and $\alpha = \beta = 90^{\circ}$, $\gamma = 120^{\circ}$ axial angles.
- 2. (c) Rhombohedral crystal system

$$a=b=c$$
, $\alpha=\beta=\gamma\neq90^{\circ}$

ex - $NaNO_3$, $CaSO_4$, calcite $CaCO_3$, HgS

3. (b) Tetragonal system has the unit cell dimension $a = b \neq c, \alpha = \beta = \gamma = 90^{\circ}$.

- **5.** (a) Space lattice of CaF_2 is face centred cubic.
- 6. (a) For body centred cubic arrangement coordination number is 8 and radius ratio (r_+/r_-) is 0.732 - 1.000.
- 7. (b) There are 14 Bravais lattices (space lattices).
- (d) Monoclinic sulphur is an example of Monoclinic crystal system.
- **10.** (b) r = 0.414 r.
- 11. (c) Each unit cell of *NaCl* contains 4 *NaCl* units.
- 12. (c) For tetrahedral arrangement co-ordination number is 4 and radius ratio (r_+/r_-) is 0.225 0.414.
- **13.** (a) Face-centred cubic lattice found in *KCl* and *NaCl*.
- 14. (c) Definition of unit cell.
- **16.** (b) In *NaCl* (rock salt) : Number of Na^+ ions = 12

(at edge centers) $\times \frac{1}{4} + 1$ (at body centre) $\times 1 = 4$. Number of Cl^- ions = 8 (at corners) $\times \frac{1}{8} + 6$ (at face centre) $\times \frac{1}{2} = 4$. Thus

4 formula units per unit cell.

- **17.** (b) Lowest potential energy level provides stable arrangement.
- 18. (b) The seven basic crystal lattice arrangements are :- Cubic, Tetragonal, Orthorhombic, Monoclinic, Hexagonal, Rhombohedral and Triclinic.
- **19.** (a) The conditions for monoclinic crystal system.
- **20.** (a) High lattice energy of $BaSO_4$ causes low solubility of $BaSO_4$ in water.
- 21. (c) 14 kinds of Bravais lattices (space lattices) are possible in a crystal.
- 22. (d) Radius ratio in *TlCl* is 0.732 1.000 and coordination number is 8 and arrangement is body centred cubic.
- 23. (c) Zinc blende (ZnS) has fcc structure and is an ionic crystal having 4 : 4 co-ordination number.
- **24.** (d) Na_2O has antifluorite (A_2B) type structure.
- **25.** (b) Zinc blende (*ZnS*) has *fcc* structure and is an ionic crystal having 4 : 4 co-ordination number.

28. (d)
$$\frac{1}{8} \times 8$$
 (at corners) = 1
 $\frac{1}{2} \times 6$ (at face centre) = 3
 $Z = 1 + 3 = 4$ (total no. of atoms)

Crystal packing

(b) Number of tetrahedral voids in the unit cell
 = 2 × number of atoms =

- (b) The system ABC ABC..... is also referred to as face-centred cubic or fcc.
- **3.** (d) It represents *ccp* arrangement.
- **4.** (a) *BCC* has a coordination number of 8.
- 5. (b) In rock salt structure the co-ordination number of $Na^+: Cl^-$ is 6:6.
- 6. (d) The *bcc* cell consists of 8 atoms at the corners and one atom at centre.

$$\therefore n = \left(8 \times \frac{1}{8}\right) + 1 = 2$$

The *fcc* cell consists of 8 atoms at the eight corners and one atom at each of the six faces. This atom at the face is shared by two unit cells.

$$\therefore n = 8 \times \frac{1}{8} + \left(6 \times \frac{1}{2}\right) = 4.$$

7. (c) AB_2 type of structure is present in CaF_2

$$\therefore AB_2 \rightleftharpoons A^{2+} + 2B^-; \quad CaF_2 \rightleftharpoons Ca^{2+} + 2F^-$$

- **8.** (b) Potassium (*K*) has *bcc* lattice.
- 9. (b) Number of atoms per unit cell in *bcc* system = 2.
- (b) In body centred cubic, each atom/ion has a coordination number of 8.
- 11. (d) Number of octahedral sites = Number of sphere in the packing.
 Number of octahedral sites per sphere = 1

12. (c) *ABAB* is hexagonal close pac13. (a) Sodium (*Na*) is a body cube.

- **14.** (b) SrF_2 has fluorite (CaF_2) type structure.
- **15.** (b) In *ZnS* structure, sulphide ions occupy all *FCC* lattice points while Zn^{2+} ions are present in alternate tetrahedral voids.
- **16.** (b) *MgO* contains rock salt (*NaCl*) structure.
- **17.** (c) CaF_2 (fluorite) has fcc structure with 8 : 4 coordination number.
- 18. (b) Every constituent has two tetrahedral voids. In *ccp* lattice atoms

$$= 8 \times \frac{1}{8} + 6 \times \frac{1}{2} = 4$$

 \therefore Tetrahedral void = 4 × 2 = 8,

- Thus ratio = 4:8::1:2.
- **19.** (c) Tetrahedral sites one double comparable to octahedral sites then ratio of X and Z respectively 2 : 1 since formula of the compound X_2Z .
- **20.** (c) A atoms are at eight corners of the cube. Therefore, the no. of A atoms in the unit cell = $\frac{8}{8} = 1$. B atoms are at the face centre of six faces. Therefore, its share in the unit cell = $\frac{6}{2} = 3$. The formula is AB_3 .
- 21. (a) In bcc structure 68% of the available volume is occupied by spheres. Thus vacant space is 32%.

(c) Number of atoms in the cubic close packed 22. structure = 8. Number of octahedral voids $=\frac{1}{2} \times 8 = 4$.

(a) Co-ordination number in HCP and CCP arrangement is 12 while in *bcc* arrangement is

8. (d) In *NaCl* (rock salt) : Number of Na^+ ions = 12 24. (at edge centers) $\times \frac{1}{4} + 1$ (at body centre) $\times 1 = 4$.

Number of Cl^{-} ions = 8 (at corners) $\times \frac{1}{2} + 6$ (at

face centre) $\times \frac{1}{2} = 4$. Thus 4 formula units per unit cell.

- (b) Co-ordination number in HCP = 1225. Co-ordination number in Mq is also = 12
- (a) All are the iso-electronic species but Na^+ has 26. low positive charge so have largest radius.

Mathematical analysis of cubic system and **Bragg's equation**

(b) Density of unit cell 1. $= \frac{N \times \text{mol.wt.}(M)}{V(=a^3) \times \text{avogadro no.}(N_o)} g \, cm^{-3}$

23.

- (b) Distance between K^+ and $F^- = \frac{1}{2} \times \text{length of}$ 2. the edge
- 3. (a) There are two atoms in a *bcc* unit cell.

So, number of atoms in 12.08×10^{23} unit cells $= 2 \times 12.08 \times 10^{23} = 24.16 \times 10^{23} atom$.

- (b) bcc structure has one atom shared by 1 unit 4. cell.
- (b) The structural arrangement of co-ordination 5٠ number '6' is octahedral and its radius ratio is 0.414 - 0.732. The example of octahedral is KCl and NaCl.
- (a) The number of spheres in one body centred 6. cubic and in one face centred cubic unit cell is 2 and 4 respectively.
- (b) Closest approach in bcc lattice 7.

$$= \frac{1}{2} \text{ of body diagonal} = \frac{1}{2} \times \sqrt{3}a = \frac{\sqrt{3}}{2} \times 4.3 = 3.72 \text{ Å}$$

10. (a)
$$M = \frac{p \times a \times N_0 \times 10}{z}$$

= $\frac{10 \times (100)^3 \times (6.02 \times 10^{23}) \times 10^{-30}}{4} = 15.05$
No. of atoms in 100

$$=\frac{6.02\times10^{23}}{15.05}\times100 = 4\times10^{25}.$$

- (c) Cs^+ and I^- have largest sizes. 11.
- (c) $58.5 g \text{ NaCl} = 1 \text{ mole} = 6.02 \times 10^{23} \text{ Na}^+ \text{Cl}^- \text{ units}$. 12.

One unit cell contains $4 Na^+Cl^-$ units. Hence number of unit cell present

$$=\frac{6.02\times10^{23}}{4}=1.5\times10^{23}.$$

13. (a)
$$\frac{1}{58.5} \times 6.023 \times 10^{23} = 1.029 \times 10^{22}$$

A unit cell contains $4 Na^+$ ion and $4 Cl^-$ ions

:: Unit cell =
$$\frac{1.029 \times 10^{22}}{4} = 2.57 \times 10^{21}$$
 unit cell.

- (b) Bragg's equation is $n\lambda = 2d\sin\theta$ 14. where *n* is an integer *i.e.* 1, 2, 3, 4 etc.
- (d) Face centred cubic structure contribute of 1/8 15. by each atom present on the corner and 1/2 by each atom present on the face.
- (c) As *CsCl* is body-centred, $d = \sqrt{3}a/2$. 16.
- (a) Radius of *Na* (if *bcc* lattice) = $\frac{\sqrt{3}a}{4} = \frac{\sqrt{3} \times 4.29}{4}$ 17.

$$= 1.8574 \text{ Å} = 1.8574 \times 10^{-8} \text{ cm}$$

(b) The crystals in which radius ratio value is 18. between 0.225 – 0.414 found shows tetrahedral crystal structure.

19. (d) For
$$bcc, d = \frac{\sqrt{3}}{2}a$$
 or $a = \frac{2d}{\sqrt{3}} = \frac{2 \times 4.52}{1.732} = 5.219$ Å = 522 pm

$$\rho = \frac{Z \times M}{a^3 \times N_0 \times 10^{-30}} = \frac{2 \times 39}{(522)^3 \times (6.023 \times 10^{23}) \times 10^{-30}}$$
$$= 0.91 \ g / cm^3 = 910 \ kg \ m^{-3}$$

- (c) The value of ionic radius ratio is 0.52 which is 20. between 0.414 - 0.732, then the geometrical arrangement of ions in crystal is octahedral.
- 21. (c) The number of atoms present in sc, fcc and bcc unit cell are 1, 4, 2 respectively.
- (b) The number of atoms present in *sc*, *fcc* and *bcc* 22. unit cell are 1, 4, 2 respectively.

23. (d)
$$Cl^- Na^+ Cl^-$$

 $\overbrace{x}{a}$ $a = 2x$

24. (a)
$$r = \frac{a}{2\sqrt{2}}$$
; $r = \frac{620}{2\sqrt{2}} = 219.25 Pm$
25. (c) $Z = \frac{V \times N_0 \times d}{2\sqrt{2}}$

$$=\frac{4.2\times8.6\times8.3\times10^{-24}\times6.023\times10^{23}\times3.3}{155}=3.84=4$$

26. (b) Volume of unit cell =
$$a^3$$

= $(3.04 \times 10^{-8} cm)^3 = 2.81 \times 10^{-23} cm^3$

2

g

$$bcc, d = \frac{\sqrt{3}}{2}a$$
 or

bcc,
$$d = \frac{\sqrt{3}}{2}a$$

$$4r = \sqrt{2}a$$
$$a = \frac{4r}{\sqrt{2}}$$

Crystal structure and Coordination number

1. (b) In a unit cell, W atoms at the corner $=\frac{1}{8} \times 8 = 1$

O atoms at the centre of edges $=\frac{1}{4} \times 12 = 3$

Na atoms at the centre of the cube = 1 W: O: Na = 1:3:1, hence formula = *NaWO*₃

- 2. (d) For *bcc* lattice, co-ordination number is 8.
- **3.** (b) Body centered cubic lattice has a co-ordination number 8.
- 4. (a) A atoms are at eight corners of the cube. Therefore, the number of A atoms in the unit $cell = \frac{8}{8} = 1$, atoms B per unit cell = 1. Hence the formula is AB.
- 5. (d) Co-ordination number for *Cu* is 12.
- 6. (b) Each Cs^+ in CsCl is surrounded by eight $Cl^$ and each Cl^- in CsCl is surrounded by eight Cs^+ .
- 7. (d) X atoms are at eight corners of the cube. Therefore, the number of X atoms in the unit $\operatorname{cell} = \frac{8}{8} = 1$.

Y atoms are at the face centre of six faces. Therefore, its share in the unit cell $=\frac{6}{2}=3$.

The formula is XY_3 .

8. (a) Let the units of ferrous oxide in a unit cell = n, molecular weight of ferrous oxide $(FeO) = 56 + 16 = 72 g mol^{-1}$,

weight of *n* units $=\frac{72 \times n}{6.023 \times 10^{23}}$

Volume of one unit = $(\text{length of corner})^3$

$$(5\mathring{A})^3 = 125 \times 10^{-24} \, cm^3$$

Density =
$$\frac{\text{wt.of cell}}{\text{volume}}$$
,
 $4.09 = \frac{72 \times n}{6.023 \times 10^{23} \times 125 \times 10^{-24}}$
 $n = \frac{3079.2 \times 10^{-1}}{72} = 42.7 \times 10^{-1} = 4.27 \approx 4$

- **9.** (b) In *NaCl* crystal *Na*⁺ ions has coordination number 6.
- **10.** (b) Cl^{-} ions in CsCl adopt *BCC* type of packing.
- 11. (d) There were 6 *A* atoms on the face-centres removing face-centred atoms along one of the axes means removal of 2 *A* atoms. Now, number of *A* atoms per unit cell

$$= 8 \times \frac{1}{8} + 4 \times \frac{1}{2} = 3$$
(corners) (face-centred)
Number of *B* atoms per unit cell

$$= 12 \times \frac{1}{4} + 1 = 4$$
(edge centred) (body centred)

Hence the resultant stoichiometry is A_3B_4

- **12.** (a) In Cs^+Cl^- crystal co-ordination number of each ion is 8.
- **13.** (b) $r_+/r_- = \frac{180}{187} = 0.962$ which lies in the range of 0.732 1.000, hence co-ordination number = 8 *i.e.* the structure is C_SCl type.
- 14. (a) In diamond, C-atoms are arranged in a regular tetrahedral structure.
- 15. (b) In *hcp*, co-ordination no. is 12.
- **16.** (c) *Mg* has 6 co-ordination number (*fcc* structure).
- 17. (d) In NaCl crystal every Na⁺ ion is surrounded by 6 Cl⁻ ion and every chloride ion is surrounded by 6 Na⁺ ion.
- **18.** (d) Crystals show good cleavage because their constituent particles are arranged in planes.
- **19.** (b) Fe_3O_4 is a non-stoichiometric compound because in it the ratio of the cations to the anions becomes different from that indicated by the chemical formula.
- 20. (d) The radius ratio for co-ordination number 4, 6 and 8 lies in between the ranges [0.225 0.414], [0.414 0.732] and [0.732 1] respectively.
- (c) The radius ratio for co-ordination number 4, 6 and 8 lies in between the ranges [0.225 0.414], [0.414 0.732] and [0.732 1] respectively.
- **22.** (b) In Na_2O , each oxide ions (O^{2^-}) is co-ordinated to $8 Na^+$ ions and each Na^+ ion to 4 oxide ions. Hence it has 4 : 8 co-ordination.
- 23. (b) When radius ratio between 0.732 -1, then co-ordination number is 8 and structural arrangement is body-centred cubic.
- 24. (c) Each Cs^+ is surrounded by eight Cl^- ions in CsCl crystal lattice because its co-ordination number is 8 : 8.
- **25.** (a) *NaCl* has *fcc* arrangement of ions.
- **26.** (c) Each Na^+ is surrounded by six Cl^- ions in NaCl crystal lattice because its co-ordination number is 6 : 6.
- 27. (b) Zinc blende (ZnS) has fcc structure and is an ionic crystal having 4 : 4 co-ordination number.
- **30.** (d) In a simple cubic structure

 $z = \frac{1}{8} \times 8$ (atoms one at a corners)

z = 1

- **31.** (a) Co-ordination number in hcp structure is 12.
- **32.** (acd) A metal that crystallizes in *bcc* structure has a co-ordination number of 8.
- **33.** (a) In sodium chloride, each Na^+ ion is surrounded by six Cl^- ions and each Cl^- ion is surrounded by six Na^+ ions. Thus, both the ions have coordination number six.
- **34.** (b) The Ca^{2+} ions are arranged in (ccp) arrangement, i.e. Ca^{2+} ions are present at all corners and tat the centre of each face of the cube. the fluoride ions occupy all the tetrahedral sites. This is 8 : 4 arrangement i.e., each Ca^{2+} ion is surrounded by $8F^{-}$ ions and each F^{-} ion by four Ca^{2+} ions.

Defects in crystal

- (c) When polar crystal is subjected to a mechanical stress, electricity is produced – a case of piezoelectricity. Reversely, if electric field is applied, mechanical stress is developed. Piezoelectric crystal acts as a mechanical electrical transductor.
- (b) More is the Schottky defect in crystal more is the decrease in density.
- **3.** (d) All the given statements are correct about *F*-centres.
- 5. (a) As each Sr^{2+} ion introduces one cation vacancy, therefore concentration of cation vacancies = mol % of $SrCl_2$ added.
- (c) Yellow colour on heating *NaCl* in presence of *Na* is due to presence of electrons in anion vacancies (*F*-centres).
- 7. (d) Frenkel's defect is due to shift of an ion from the normal lattice site (Creating a vacancy) and occupy interstitial spaces.
- 8. (c) A_gBr exhibits Frenkel defect due to large difference in the size of A_g^+ and Br^- ions.
- **9.** (d) Schottky defects occurs in highly ionic compounds which have high co-ordination number ex. *NaCl*, *KCl*, *CsCl*.
- **10.** (d) Schottky defect is due to missing of equal number of cations and anions.
- **11.** (a) Schottky defect is due to missing of equal number of cations and anions.
- (a) Impurity present in a crystal does not establish thermal equilibrium.
- 13. (c) Since no ions are missing from the crystal as a whole, there is no effect on density.
- **15.** (b) On adding non-metal in metal the metal becomes less tensile.
- **16.** (c) AgBr exhibits Frenkel defect due to large difference in the size of Ag^+ and Br^- ions.
- 17. (c) Both are stoichiometric crystalline defects.

- 18. (c) Brass, Cu = 80%, Zn = 20% substitutional alloy.
 Steel is an interstitial alloy because it is an alloy of *Fe* with *C*, *C* atoms occupy the interstitial voids of *Fe* crystal.
- 19. (d) In metal excess defect when holes created by missing of anions are occupied by electrons, there sites are called *F*-centres and are responsible for colour in the crystal.
- **20.** (c) *KBr* exhibits Schottky defect and not Frenkel defect.
- **21.** (c) When cation shifts from lattice to interstitial site, the defect is called Frenkel defect.
- **22.** (d) F-centres are the sites where anions are missing and instead electrons are present. they are responsible for colour.

Critical Thinking Questions

- (c) Amorphous solids neither have ordered arrangement (*i.e.* no definite shape) nor have sharp melting point like crystals, but when heated, they become pliable until they assume the properties usually related to liquids. It is therefore they are regarded as super-cooled liquids.
- (c) Silicon due to its catenation property form network solid.
- 3. (c) Orthorhombic geometry has $a \neq b \neq c$ and $\alpha = \beta = \gamma = 90^{\circ}$. The shape of match box obey this geometry.
- (d) In a triclinic crystal has no notation of symmetry.
- (a) In NH₃ molecule, the original appearance is repeated as a result of rotation through 120°. Such as axis is said to be an axis of three-fold symmetry or a triad axis.
- **6.** (a) Na_2O has antifluorite (A_2B) type structure.
- 7. (b) Cationic radius increases down the group and decreases along the period.
- 8. (c) Distance between centres of cation and anion $= \frac{d}{2} = \frac{508}{2} = 254 \ pm$

$$r_{a} + r_{a} = 254 \ pm$$
 or $110 + r_{a} = 254$ or $r_{a} = 144 \ pm$

9. (b)
$$\rho = \frac{n \times M}{a^3 \times N_0 \times 10^{-30}}$$

2 × 100

$$=\frac{2\times100}{(400)^3\times(6.02\times10^{23})\times10^{-30}}=5.188\ g/cm^3$$

10. (a) NaCl structure $\xrightarrow{\text{High}}$ CsCl structure (6:6 co.-ord.) (8:8 co.-ord.)

11. (a) Difference =
$$2.178 \times 10^3 - 2.165 \times 10^3 = 0.013 \times 10^3$$

Fraction unoccupied = $\frac{0.013 \times 10^3}{2.178 \times 10^3} = 5.96 \times 10^{-3}$

12. (c) $CsBr_3$ consist of Cs^+ and Br_3^- ions.

- 13. (a) Each Cs^+ is surrounded by eight Cl^- ions in CsCl crystal lattice because its co-ordination number is 8 : 8.
- 14. (b) In each CaF_2 each calcium cation is surrounded by eight fluoride anions in a body centred cubic arrangement. Each fluoride ion is in contact with four calcium ions. Thus CaF_2 has 8 : 4 co-ordination number.
- **15.** (c) The radius ratio for co-ordination number 4, 6 and 8 lies in between the ranges [0.225 - 0.414], [0.414 - 0.732] and [0.732 - 1]respectively.



- 17. (b) In this diagram, equal number of cations (Na⁺) and anions (Cl⁻) are missing, so it, shows schottky defect.
- 18. (c) Zeolite is a three dimensional silicate because of in the silicates all the four oxygen atoms at $(SiO_4)^{-4}$ tetrahedra are shared with other tetrahedra, vesulting in a three dimensional network.

Assertion & Reason

- (b) It is true that in the dimond structure, carbon atoms are arranged in tetrahedrally (*sp*³ hybridized) but it is not the correct explanation of assertion.
- (d) Tetrahedral holes are smaller in size than octahedral holes. Cations usually occupy less space than anions.
- (d) Crystalline solids have regular arrangement of constituent particles and are anisotropic whereas amorphous solids have no regular arrangement and are isotropic.
- (a) Schottky defect is due to missing of equal number of cations and anions.
- 5. (b) Space or crystal lattice is a regular repeating arrangement of points in space and forms the basis of classification of all structures.
- 6. (c) Tetrahedral void is so called because it is surrounded by four spheres tetrahedrally while octahedral void is so called because it is surrounded by six spheres octahedrally.

- (c) Two corners per tetrahedron one shared in both the cases.
- 8. (b) When an atom or an ion is missing from its normal lattice site, a lattice vacancy or defect is created, which is called schottky defect. Due to missing density of crystal will be lowered.
- 9. (b) On heating, the metal atoms deposit on the surface and finally they deffuse into the crystal and after ionisation the alkali metal ion occupies cationic vacancy where as electron occupies anionic vacancy.
- 10. (a) In case of semiconductors, the gap between valence band and the conduction band is small and there fore some of the electrons may jump from valence band to conduction band and thus on increasing temperature conductivity is also increased.
- (a) All magnetically ordered solids (ferromagnetic and antiferromagnetic solids) transform to the paramagnetic state at high temperature due to the randomisation of spins.
- 12. (c) In piezoelectric crystals, the dipoles may align them selves in an ordered manner such that there is a net dipole moment in the crystal.