

Chapter 1 Real Number

Question-1

Write the following rational numbers in decimal form:

(i) $\frac{42}{100}$

(ii) $\frac{327}{500}$

(iii) $3\frac{3}{8}$

(iv) $\frac{5}{6}$

(v) $\frac{1}{5}$

(vi) $\frac{1}{7}$

(vii) $\frac{2}{13}$

(viii) $\frac{11}{17}$

Solution:

(i) $\frac{42}{100} = 0.42$

(ii) $\frac{327}{500} = 0.654$

$$\begin{array}{r} 0.654 \\ 5 \overline{) 3.27} \\ \underline{30} \\ 27 \\ \underline{25} \\ 20 \\ \underline{20} \\ 0 \end{array}$$

(iii) $3\frac{3}{8} = \frac{27}{8} = 3.375$

$$\begin{array}{r} 3.375 \\ 8 \overline{) 27} \\ \underline{24} \\ 30 \\ \underline{24} \\ 60 \\ \underline{56} \\ 40 \\ \underline{40} \\ 0 \end{array}$$

$$(iv) \frac{5}{6} = 0.8333... = 0.\overline{8333}$$

$$\begin{array}{r} 0.8333 \\ 6 \overline{) 5.0000} \\ \underline{50} \\ 30 \\ \underline{20} \\ 18 \\ \underline{20} \\ 18 \\ \underline{20} \\ 18 \\ \underline{20} \\ 18 \\ \underline{18} \\ 2 \end{array}$$

$$(v) \frac{1}{5} = 0.2$$

$$\begin{array}{r} 0.2 \\ 5 \overline{) 1.0} \\ \underline{10} \\ 10 \\ \underline{10} \\ 0 \end{array}$$

$$(vi) \frac{1}{7} = 0.\overline{142857}$$

$$\begin{array}{r} 0.142857 \\ 7 \overline{) 1.000000} \\ \underline{10} \\ 7 \\ \underline{30} \\ 28 \\ \underline{20} \\ 14 \\ \underline{60} \\ 56 \\ \underline{40} \\ 35 \\ \underline{50} \\ 49 \\ \underline{49} \\ 1 \end{array}$$

$$(vii) \frac{2}{13} = 0.\overline{153846}$$

$$\begin{array}{r} 0.153846 \\ 13 \overline{) 2.000000} \\ \underline{20} \\ 13 \\ \underline{70} \\ 65 \\ \underline{50} \\ 39 \\ \underline{110} \\ 104 \\ \underline{60} \\ 52 \\ \underline{80} \\ 78 \\ \underline{2} \end{array}$$

$$(viii) \frac{11}{17} = \overline{0.6470588235294117}$$

$$\begin{array}{r}
 0.6470588235294117 \\
 17 \overline{) 110} \\
 \underline{102} \\
 80 \\
 \underline{68} \\
 120 \\
 \underline{119} \\
 100 \\
 \underline{85} \\
 150 \\
 \underline{136} \\
 140 \\
 \underline{136} \\
 40 \\
 \underline{34} \\
 60 \\
 \underline{51} \\
 90 \\
 \underline{85} \\
 50 \\
 \underline{34} \\
 160 \\
 \underline{153} \\
 70 \\
 \underline{68} \\
 20 \\
 \underline{17} \\
 30 \\
 \underline{17} \\
 130 \\
 \underline{119} \\
 \underline{\underline{11}}
 \end{array}$$

Question-2

If a is a positive rational number and n is a positive integer greater than 1, prove that a^n is a rational number.

Solution:

We know that product of two rational number is always a rational number.

Hence if a is a rational number then

$a^2 = a \times a$ is a rational number,

$a^3 = a^2 \times a$ is a rational number,

$a^4 = a^3 \times a$ is a rational number,

...

...

$\therefore a^n = a^{n-1} \times a$ is a rational number.

Question-3

Find three rational numbers lying between 0 and 0.1. Find twenty rational numbers between 0 and 0.1. Give a method to determine any number of rational numbers between 0 and 0.1.

Solution:

The three rational numbers lying between 0 and 0.1 are 0.01, 0.05, 0.09.

The twenty rational numbers between 0 and 0.1 are 0.001, 0.002, 0.003, 0.004, ... 0.011, 0.012, ... 0.099.

To determine any number of rational numbers between 0 and 0.1 insert 0 after the decimal.

Question-4

Complete the following:

(i) Every point on the number line corresponds to a _____ number which may be either _____ or _____.

(ii) The decimal form of an irrational number is neither _____ or _____.

(iii) The decimal representation of the rational number $\frac{8}{27}$ is _____.

(iv) 0 is _____ number. [Hint: a rational /an irrational]

Solution:

(i) Every point on the number line corresponds to a real number which may be either rational or irrational.

(ii) The decimal form of an irrational number is neither recurring or terminating.

(iii) The decimal representation of the rational number $\frac{8}{27}$ is 0.296

(iv) 0 is a rational number.

Question-5

Which of the following rational numbers have the terminating decimal representation?

- (i) $3/5$ (ii) $7/20$ (iii) $2/13$
(iv) $27/40$ (v) $133/125$ (vi) $23/7$

[Making use of the result that a rational number p/q where p and q have no common factor(s) will have a terminating representation if and only if the prime factors of q are 2's or 5's or both.]

Solution:

(i) The prime factor of 5 is 5. Hence $3/5$ has a terminating decimal representation.

(ii) $20 = 4 \times 5 = 2^2 \times 5$.

The prime factors of 20 are both 2's and 5's. Hence $7/20$ has a terminating decimal.

(iii) The prime factor of 13 is 13. Hence $2/13$ has non-terminating decimal.

(iv) $40 = 2^3 \times 5$.

The prime factors of 40 are both 2's and 5's. Hence $27/40$ has a terminating decimal.

(v) $125 = 5^3$

The prime factor of 125 is 5's. Hence $13/125$ has a terminating decimal.

(vi) The prime factor of 7 is 7. Hence $23/7$ has a non-terminating decimal representation.

Question-6

You have seen that $\sqrt{2}$ is not a rational number. Show that $2 + \sqrt{2}$ is not a rational number.

Solution:

Let $2 + \sqrt{2}$ be a rational number say r .

Then $2 + \sqrt{2} = r$

$\sqrt{2} = r - 2$

But, $\sqrt{2}$ is an irrational number.

Therefore, $r - 2$ is also an irrational number.

$\Rightarrow r$ is an irrational number.

Hence our assumption r is a rational number is wrong.

Hence, $2 + \sqrt{2}$ is not a rational number.

Question-7

Prove that $3\sqrt{3}$ is not a rational number.

Solution:

Let $3\sqrt{3}$ be a rational number say r .

$$\text{Then } 3\sqrt{3} = r$$

$$\sqrt{3} = (1/3)r$$

$(1/3)r$ is a rational number because product of two rational number is a rational number.

$\Rightarrow \sqrt{3}$ is a rational number but $\sqrt{3}$ is not a rational number.

Therefore our assumption that $3\sqrt{3}$ is a rational number is wrong.

Question-8

Show that $\sqrt[3]{6}$ is not a rational number.

Solution:

Let $\sqrt[3]{6}$ be a rational number, say $\frac{p}{q}$ where $q \neq 0$.

$$\text{Then } \sqrt[3]{6} = \frac{p}{q}$$

Since $1^3 = 1$, and $2^3 = 8$, it follows that $1 < \frac{p}{q} < 2$

Then $q > 1$ because if $q = 1$ then $\frac{p}{q}$ will be an integer, and there is no integer between 1 and 2.

$$\text{Now, } 6 = \left(\frac{p}{q}\right)^3$$

$$6 = \frac{p^3}{q^3}$$

$$6q^3 = \frac{p^3}{q}$$

q being an integer, $6q^3$ is an integer, and since $q > 1$ and q does not have a common factor with p and consequently with p^3 .

So, $\frac{p^3}{q}$ is a fraction different from an integer.

$$\text{Thus } 6q^3 \neq \frac{p^3}{q}.$$

This contradiction proves the result.

Question-9

Identify the following as rational or irrational numbers. Give the decimal representation of rational numbers.

(i) $\sqrt{4}$

(ii) $3\sqrt{18}$

(iii) $\sqrt{1.44}$

(iv) $\sqrt{\frac{9}{27}}$

(v) $-\sqrt{0.64}$

(vi) $\sqrt{100}$

Solution:

(i) $\sqrt{4} = 2$ is rational.

(ii) $3\sqrt{18} = 3\sqrt{9 \times 2} = 3 \times 3\sqrt{2} = 9\sqrt{2}$ is irrational.

(iii) $\sqrt{1.44} = \sqrt{\frac{144}{100}} = \frac{12}{10} = 1.2$ is rational.

(iv) $\sqrt{\frac{9}{27}} = \sqrt{\frac{9}{9 \times 3}} = \sqrt{\frac{1}{3}} = \frac{1}{\sqrt{3}}$ is irrational.

(v) $-\sqrt{0.64} = -0.8$ is rational.

(vi) $\sqrt{100} = 10$ is rational.

Question-10

In the following equations, find which of the variables x, y, z etc. represent rational numbers and which represent irrational numbers:

(i) $x^2 = 5$

(ii) $y^2 = 9$

(iii) $z^2 = 0.04$

(iv) $u^2 = 17/4$

(v) $v^2 = 3$

(vi) $w^3 = 27$

(vii) $t^2 = 0.4$

Solution:

(i) $x^2 = 5$

$\therefore x = \sqrt{5}$ is irrational.

(ii) $y^2 = 9$

$\therefore y = 3$ is rational.

(iii) $z^2 = 0.04$

$\therefore z = 0.2$ is rational.

(iv) $u^2 = \frac{17}{4}$

$$\begin{aligned}\therefore u &= \sqrt{\frac{17}{4}} \\ &= \frac{\sqrt{17}}{\sqrt{4}} = \frac{\sqrt{17}}{2} \text{ is irrational.}\end{aligned}$$

(v) $v^2 = 3$

$\therefore v = \sqrt{3}$ is irrational.

(vi) $w^3 = 27$

$$w = \sqrt[3]{27} = 3 \text{ is rational.}$$

(vii) $t^2 = 0.4 \therefore t = \sqrt{0.4} = \sqrt{\frac{4}{10}} = \frac{2}{\sqrt{10}}$ is irrational.

Question-11

Give an example to show that the product of a rational number and an irrational number may be a rational number.

Solution:

A rational number 0 multiplied by an irrational number gives the rational number 0.

Question-12

State with reason which of the following are surds and which are not.

(i) $\sqrt{5} \times$

(ii) $\sqrt{8} \times$

(iii) $\sqrt{27} \times \sqrt{3}$

(iv) $\sqrt{16} \times \sqrt{4}$

(v) $5\sqrt{8} \times 2\sqrt{6}$

(vi) $\sqrt{125} \times \sqrt{5}$

(vii) $\sqrt{100} \times \sqrt{2}$

(viii) $6\sqrt{2} \times 9\sqrt{3}$

(ix) $\sqrt{120} \times \sqrt{45}$

(x) $\sqrt{15} \times \sqrt{6}$.

Solution:

(i) $\sqrt{5} \times \sqrt{10} = \sqrt{5} \times \sqrt{5 \times 2} = \sqrt{5} \times \sqrt{5} \times \sqrt{2} = 5\sqrt{2}$ is a surd.

(ii) $\sqrt{8} \times \sqrt{6} = \sqrt{4 \times 2} \times \sqrt{3 \times 2} = 2\sqrt{2} \times \sqrt{2} \times \sqrt{3} = 4\sqrt{3}$ is a surd.

(iii) $\sqrt{27} \times \sqrt{3} = \sqrt{9 \times 3} \times \sqrt{3} = 3\sqrt{3} \times \sqrt{3} = 9$ is not a surd.

(iv) $\sqrt{16} \times \sqrt{4} = 4 \times 2 = 8$ is not a surd.

(v) $5\sqrt{8} \times 2\sqrt{6} = 5\sqrt{4 \times 2} \times 2\sqrt{3 \times 2} = 5 \times 2\sqrt{2} \times 2\sqrt{2} \times \sqrt{3} = 5 \times 2 \times 2 \times 2 \times \sqrt{3} = 40\sqrt{3}$ is a surd.

(vi) $\sqrt{125} \times \sqrt{5} = \sqrt{25 \times 5} \times \sqrt{5} = 5\sqrt{5} \times \sqrt{5} = 5 \times 5 = 25$ is not a surd.

(vii) $\sqrt{100} \times \sqrt{2} = 10\sqrt{2}$ is a surd.

(viii) $6\sqrt{2} \times 9\sqrt{3} = 54\sqrt{6}$ is a surd.

(ix) $\sqrt{120} \times \sqrt{45} = \sqrt{4 \times 30} \times \sqrt{9 \times 5} = 2\sqrt{6 \times 5} \times 3\sqrt{5} = 2 \times \sqrt{6} \times \sqrt{5} \times 3 \times \sqrt{5} = 30\sqrt{6}$ is a surd.

(x) $\sqrt{15} \times \sqrt{6} = \sqrt{5 \times 3} \times \sqrt{2 \times 3} = \sqrt{5} \times \sqrt{3} \times \sqrt{2} \times \sqrt{3} = 3 \times \sqrt{10}$ is a surd.

Question-13

Give two examples to show that the product of two irrational numbers may be a rational number.

Solution:

Take $a = (2 + \sqrt{3})$ and $b = (2 - \sqrt{3})$; a and b are irrational numbers, but their product

$$(2 + \sqrt{3})(2 - \sqrt{3}) = 4 - 3 = 1 \text{ is a rational number.}$$

Take $c = \sqrt{3}$ and $d = -\sqrt{3}$; c and d are irrational numbers, but their product = -3, is a rational number.

Question-14

Find the value of $\sqrt{5}$ correct to two places of $\sqrt{5}$ decimal.

Solution:

We know that $2^2 = 4 < 5 < 9 = 3^2$

Taking positive square roots we get

$$2 < \sqrt{5} < 3.$$

Next, $(2.2)^2 = 4.84 < 5 < 5.29 = (2.3)^2$

Taking positive square roots, we have

$$2.2 < \sqrt{5} < 2.3$$

Again, $(2.23)^2 = 4.9729 < 5 < 5.0176 = (2.24)^2$

Taking positive square roots, we obtain

$$2.23 < \sqrt{5} < 2.24$$

Hence the required approximation is 2.24 as $(2.24)^2$ is nearest to 5 than $(2.23)^2$.

Question-15

Prove that $\sqrt{3} - \sqrt{2}$ is irrational.

Solution:

Let $\sqrt{3} - \sqrt{2}$ be a rational number, say r

$$\text{Then } \sqrt{3} - \sqrt{2} = r$$

On squaring both sides we have

$$(\sqrt{3} - \sqrt{2})^2 = r^2$$

$$3 - 2\sqrt{6} + 2 = r^2$$

$$5 - 2\sqrt{6} = r^2$$

$$-2\sqrt{6} = r^2 - 5$$

$$\sqrt{6} = -(r^2 - 5)/2$$

Now $-(r^2 - 5)/2$ is a rational number and $\sqrt{6}$ is an irrational number.

Since a rational number cannot be equal to an irrational number. Our assumption that

$\sqrt{3} - \sqrt{2}$ is rational is wrong.

Question-16

Prove that $\sqrt{3} + \sqrt{5}$ is an irrational number.

Solution:

Let $\sqrt{3} + \sqrt{5}$ be a rational number, say r

Then $\sqrt{3} + \sqrt{5} = r$

On squaring both sides,

$$(\sqrt{3} + \sqrt{5})^2 = r^2$$

$$3 + 2\sqrt{15} + 5 = r^2$$

$$8 + 2\sqrt{15} = r^2$$

$$2\sqrt{15} = r^2 - 8$$

$$\sqrt{15} = (r^2 - 8)/2$$

Now $(r^2 - 8)/2$ is a rational number and $\sqrt{15}$ is an irrational number.

Since a rational number cannot be equal to an irrational number. Our assumption that

$\sqrt{3} + \sqrt{5}$ is rational is wrong.

Question-17

Examine, whether the following numbers are rational or irrational:

(i) $(\sqrt{2} + 2)^2$

(ii) $(2 - \sqrt{2}) \times (2 + \sqrt{2})$

(iii) $(\sqrt{2} + \sqrt{3})^2$

(iv) $\frac{6}{3\sqrt{2}}$

Solution:

(i) $(\sqrt{2} + 2)^2 = (\sqrt{2})^2 + 2\sqrt{2} \times 2 + (2)^2 = 2 + 4\sqrt{2} + 4 = 6 + 4\sqrt{2}$.

\ It is an irrational number.

(ii) $(2 - \sqrt{2}) \times (2 + \sqrt{2}) = (2)^2 - (\sqrt{2})^2 = 4 - 2 = 2$.

\ It is a rational number.

(iii) $(\sqrt{2} + \sqrt{3})^2 = (\sqrt{2})^2 + 2\sqrt{2} \times \sqrt{3} + (\sqrt{3})^2 = 2 + 2\sqrt{6} + 3 = 5 + 2\sqrt{6}$

\ It is an irrational number.

(iv) $\frac{6}{3\sqrt{2}} = \frac{2}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{2\sqrt{2}}{2} = \sqrt{2}$

\ It is an irrational number.

Question-18

Prove that

(a) $2 + \sqrt{3}$ is not a rational number and

(b) $\sqrt[3]{7}$ is not a rational number.

Solution:

(a) If possible, let $2 + \sqrt{3} = a$, where a is rational.

$$\text{Then, } (2 + \sqrt{3})^2 = a^2$$

$$7 + 4\sqrt{3} = a^2$$

$$\sqrt{3} = \frac{a^2 - 7}{4} \text{ -----(i)}$$

Now, a is rational $\Rightarrow \frac{a^2 - 7}{4}$ is rational.

$\sqrt{3}$ is rational [from (i)]

This is a contradiction.

Hence, $2 + \sqrt{3}$ is not a rational number.

(b) If possible, let $\sqrt[3]{7} = \frac{p}{q}$, where p and q are integers, having no common factors and $q \neq 0$.

$$\text{Then, } (\sqrt[3]{7})^3 = \left(\frac{p}{q}\right)^3$$

$$\Rightarrow 7q^3 = p^3 \text{ -----(i)}$$

$\Rightarrow p^3$ is a multiple of 7

$\Rightarrow p$ is multiple of 7.

Let $p = 7m$, where m is an integer.

$$\text{Then, } p^3 = 343 m^3 \text{ -----(ii)}$$

$$\Rightarrow 7q^3 = 343 m^3 \text{ [from (i) and (ii)]}$$

$$\Rightarrow q^3 = 49 m^3 \Rightarrow q^3 \text{ is a multiple of 7.}$$

$\Rightarrow q$ is a multiple of 7.

Thus, p and q are both multiples of 7, or 7 is a factor of p and q .

This contradicts our assumption that p and q have no common factors.

Hence $\sqrt[3]{7}$ is not a rational number.

Question-19

Examine whether the following numbers are rational or irrational:

(i) $(3 + \sqrt{2})^2$

(ii) $(3 - \sqrt{3})(3 + \sqrt{3})$

(iii) $\frac{6}{2\sqrt{3}}$

Solution:

(i) $(3 + \sqrt{2})^2 = 9 + 2 + 6\sqrt{2} = 11 + 6\sqrt{2}$, which is irrational.

(ii) $(3 - \sqrt{3})(3 + \sqrt{3}) = (3)^2 - (\sqrt{3})^2 = 9 - 3 = 6$, which is rational.

(iii) $\frac{6}{2\sqrt{3}} = \frac{6}{2\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{6 \times \sqrt{3}}{6} = \sqrt{3}$, which is irrational.

Question-20

Find two irrational numbers lying between $\sqrt{2}$ and $\sqrt{3}$.

Solution:

Irrational numbers lying between $\sqrt{2}$ and $\sqrt{3}$ is $\sqrt{\sqrt{2} \times \sqrt{3}}$, i.e. $\sqrt{\sqrt{6}} = 6^{(1/4)}$

Irrational numbers lying between $\sqrt{2}$ and $6^{(1/4)}$ is $\sqrt{\sqrt{2} \times 6^{1/4}} = 2^{(1/4)} \times 6^{(1/8)}$

Hence two irrational numbers lying between $\sqrt{2}$ and $\sqrt{3}$ are $6^{(1/4)}$ and $2^{(1/4)} \times 6^{(1/8)}$.

Question-21

Express $\frac{7}{64}$ as a decimal fraction.

Solution:

$$\begin{array}{r}
 0.109375 \\
 64 \overline{) 7.000} \\
 \underline{64} \\
 600 \\
 \underline{576} \\
 240 \\
 \underline{192} \\
 480 \\
 \underline{448} \\
 320 \\
 \underline{320} \\
 0
 \end{array}$$

Therefore $\frac{7}{64} = 0.109375$

Question-22

Express $\frac{12}{125}$ as a decimal fraction.

Solution:

$$\begin{array}{r} 0.096 \\ 125 \overline{)12.000} \\ \underline{1125} \\ 750 \\ \underline{750} \\ 0 \end{array}$$

Therefore $\frac{12}{125} = 0.096$.

Question-23

Express $\frac{451}{13}$ as a decimal fraction.

Solution:

$$\begin{array}{r} 34.692307 \\ 13 \overline{)451.0000000} \\ \underline{39} \\ 61 \\ \underline{52} \\ 90 \\ \underline{78} \\ 120 \\ \underline{117} \\ 30 \\ \underline{26} \\ 40 \\ \underline{39} \\ 100 \\ \underline{91} \\ 9 \end{array}$$

Therefore **34.692307**