

CBSE Test Paper 01

CH-2 Polynomials

1. If $p(x) = x + 3$, then $p(x) + p(-x)$ is equal to
 - a. $2x$
 - b. 3
 - c. 0
 - d. 6
2. If $x+y+z = 0$, then $x^3 + y^3 + z^3$ is
 - a. $3xyz$
 - b. xyz
 - c. $2xyz$
 - d. 0
3. The degree of constant function is
 - a. 0
 - b. 3
 - c. 1
 - d. 2
4. $(x + 1)$ is a factor of the polynomial
 - a. $x^3 + x^2 - x + 1$
 - b. $x^3 + x^2 + x + 1$
 - c. $x^4 + 3x^3 + 3x^2 + x + 1$
 - d. $x^4 + x^3 + x^2 + 1$
5. The coefficient of 'x' in the expansion of $(x + 3)^3$ is
 - a. 1
 - b. 27
 - c. 9
 - d. 18
6. Fill in the blanks: A polynomial containing one non-zero term is called a _____.
7. Fill in the blanks: The highest power of the variable in a polynomial is called the _____ of the polynomial.
8. Write $(-3x + y + z)^2$ in the expanded form:

9. Evaluate the following by using identities: $0.54 \times 0.54 - 0.46 \times 0.46$
10. Evaluate: $185 \times 185 - 15 \times 15$
11. Evaluate 105×108 without multiplying directly.
12. Simplify $(x + y)^3 - (x - y)^3 - 6y(x + y)(x - y)$.
13. Factorize : $(a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3$
14. If both $x - 2$ and $x - \frac{1}{2}$ are factors of $px^2 + 5x + r$, show that $p = r$.
15. If the polynomials $az^3 + 4z^2 + 3z - 4$ and $z^3 - 4z + a$ leave the same remainder when divided by $z - 3$, find the value of a .

CBSE Test Paper 01
CH-2 Polynomials

Solution

1. (d) 6

Explanation:

$$p(x) = x + 3$$

$$\text{And } p(-x) = -x + 3$$

$$\text{Then, } p(x) + p(-x)$$

$$= x + 3 - x + 3$$

$$= 6$$

2. (a) $3xyz$

Explanation:

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx) \Rightarrow$$

$$x^3 + y^3 + z^3 - 3xyz = (0)(x^2 + y^2 + z^2 - xy - yz - zx) \Rightarrow$$

$$x^3 + y^3 + z^3 - 3xyz = 0 \Rightarrow x^3 + y^3 + z^3 = 3xyz \text{ If } x+y+z = 0, \text{ then}$$

$$x^3 + y^3 + z^3 \text{ is } 3xyz$$

3. (a) 0

Explanation: The degree of any constant term 5 (say)

$$\text{We can write 5 as } 5 \times 1 = 5x^0 \text{ [Since } a^0 = 1]$$

Therefore the degree of any constant term is 0

4. (b) $x^3 + x^2 + x + 1$

$$\textbf{Explanation: } x^3 + x^2 + x + 1 = x^3(x + 1) + 1(x + 1) = (x^3 + 1)(x + 1)$$

5. (b) 27

$$\textbf{Explanation: } (x + 3)^3 = x^3 + (3)^3 + 3 \times x \times 3(x + 3) = x^3 + 27 + 9x^2 + 27x = x^3 + 9x^2 + 27x + 27 \text{ Therefore, the coefficient of } x, \text{ in the expansion of } (x + 3)^3 \text{ is } 27.$$

6. monomial

7. degree

8. We have,

$$(-3x + y + z)^2$$

$$\text{Using identity } (a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$$

$$\begin{aligned}
&= (-3x)^2 + y^2 + z^2 + 2 \times (-3x) \times y + 2 \times y \times z + 2 \times z \times (-3x) \\
&= 9x^2 + y^2 + z^2 - 6xy + 2yz - 6zx
\end{aligned}$$

9. We have,

$$\begin{aligned}
&0.54 \times 0.54 - 0.46 \times 0.46 \\
&= (0.54)^2 - (0.46)^2 = (0.54 + 0.46)(0.54 - 0.46) = 1 \times 0.08 = 0.08
\end{aligned}$$

10. $185 \times 185 - 15 \times 15$

$$\begin{aligned}
&(185)^2 - (15)^2 \text{ [Using } a^2 - b^2 = (a - b)(a + b)] \\
&(185 + 15)(185 - 15) \\
&200 \times 170 = 34000
\end{aligned}$$

11. $105 \times 108 = (100 + 5)(100 + 8)$

$$\text{Using identity } (x + a)(x + b) = x^2 + (a + b)x + ab$$

$$\begin{aligned}
&\text{We get, } 105 \times 108 = 100^2 + (5 + 8)100 + 5 \times 8 \\
&= 10000 + 1300 + 40 = 11340
\end{aligned}$$

12. $(x + y)^3 - (x - y)^3 - 6y(x + y)(x - y)$

$$\begin{aligned}
&= (x + y)^3 - (x - y)^3 - 3.2y(x + y)(x - y) \\
&= (x + y)^3 - (x - y)^3 - 3(x + y - x + y)(x + y)(x - y) \\
&= [x + y - x + y]^3 [\because a^3 - b^3 - 3ab(a - b) = (a - b)^3] \\
&= (2y)^3 = 8y^3
\end{aligned}$$

13. Let $x = a^2 - b^2$, $y = b^2 - c^2$ and $z = c^2 - a^2$. Then,

$$x + y + z = a^2 - b^2 + b^2 - c^2 + c^2 - a^2 = 0$$

$$\therefore x^3 + y^3 + z^3 = 3xyz$$

$$\Rightarrow (a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3 = 3(a^2 - b^2)(b^2 - c^2)(c^2 - a^2)$$

$$\Rightarrow (a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3 = 3(a + b)(a - b)(b + c)(b - c)(c + a)(c - a)$$

$$\Rightarrow (a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3 = 3(a + b)(b + c)(c + a)(a - b)(b - c)(c - a)$$

14. Let $f(x) = px^2 + 5x + r$ be the given polynomial. Since $x - 2$ and $x - \frac{1}{2}$ are factors of $f(x)$.

$$\therefore f(2) = 0 \text{ and } f\left(\frac{1}{2}\right) = 0$$

$$\Rightarrow p \times 2^2 + 5 \times 2 + r = 0 \text{ and } p\left(\frac{1}{2}\right)^2 + 5 \times \frac{1}{2} + r = 0$$

$$\Rightarrow 4p + 10 + r = 0 \text{ and } \frac{p}{4} + \frac{5}{2} + r = 0$$

$$\Rightarrow 4p + r = -10 \text{ and } \frac{p+4r+10}{4} = 0$$

$$\Rightarrow 4p + r = -10 \text{ and } p + 4r + 10 = 0$$

$$\Rightarrow 4p + r = -10 \text{ and } p + 4r = -10$$

$$\Rightarrow 4p + r = p + 4r \text{ [RHS of the two equations are equal]}$$

$$\Rightarrow 3p = 3r \Rightarrow p = r$$

15. Let $p(z) = az^3 + 4z^2 + 3z - 4$

And $q(z) = z^3 - 4z + a$

As these two polynomials leave the same remainder, when divided by $z - 3$, then $p(3) = q(3)$.

$$\begin{aligned} \therefore p(3) &= a(3)^3 + 4(3)^2 + 3(3) - 4 \\ &= 27a + 36 + 9 - 4 \end{aligned}$$

$$\text{Or } p(3) = 27a + 41$$

$$\begin{aligned} \text{And } q(3) &= (3)^3 - 4(3) + a \\ &= 27 - 12 + a = 15 + a \end{aligned}$$

$$\text{Now, } p(3) = q(3)$$

$$\Rightarrow 27a + 41 = 15 + a$$

$$\Rightarrow 26a = -26; a = -1$$

Hence, the required value of $a = -1$.