

THERMODYNAMICS



Classical thermodynamics... is the only physical theory of universal physical theory of universal content which I am convinced... will never be overthrown.

Albert Einstein



Rudolf Clausius
(1822-1888)

Learning Objectives:



After studying this unit, students will be able to

- distinguish between the system and surroundings
- define closed, open and isolated systems
- distinguish between state and path functions
- describe the relation between internal energy, work and heat
- state four laws of thermodynamics
- correlate internal energy change (ΔU) and enthalpy change (ΔH) and their measurement.
- calculate enthalpy changes for various types of reactions
- apply Hess's law to calculate lattice energy of crystals
- define spontaneous and non spontaneous processes
- establish the relationship between thermodynamic state functions enthalpy (H), entropy (S) and Gibbs free energy (G)
- list the factors that determines spontaneity of the process
- establish the link between ΔG and spontaneity, and the relationship between ΔG^0 and equilibrium constant

7.1 Introduction

The term 'Thermodynamics' means flow of heat and is derived from the Greek 'Thermos' (heat) and 'dynamics' (flow). In our daily life, we come across many useful reactions such as burning of fuel to produce heat energy, flow of electrons through circuit to produce electrical energy, metabolic reactions to produce the necessary energy for biological functions and so on. Thermodynamics, the study of the transformation of energy, explains all such processes quantitatively and allows us to make useful predictions.

In the 19th century, scientists tried to understand the underlying principles of steam engine which were already in operation, in order to improve their efficiency. The basic problem of the investigation was the transformation of heat into mechanical work. However, over time, the laws of thermodynamics were developed and helped to understand the process of steam engine. These laws have been used to deduce powerful mathematical relationships applicable to a broad range of processes.

Thermodynamics evaluates the macroscopic properties (heat, work) and their inter relationships. It deals with properties of systems in equilibrium and is independent of any theories or properties of the individual molecules which constitute the system.

The principles of thermodynamics are based on three laws of thermodynamics. The first two laws (First and second law) summarise the actual experience of inter

conversion of different forms of energy. The third law deals with the calculation of entropy and the unattainability of absolute zero Kelvin. Thermodynamics carries high practical values but bears certain limitations. It is independent of atomic and molecular structure and reaction mechanism. The laws can be used to predict whether a particular reaction is feasible or not under a given set of conditions, but they cannot give the rate at which the reaction takes place. In other words, thermodynamics deals with equilibrium conditions quantitatively, but does not take into account the kinetic approach to the equilibrium state.

7.2 System and Surrounding

Before studying the laws of thermodynamics and their applications, it is important to understand the meaning of a few terms used frequently in thermodynamics.

System:

The universe is divided into two parts, the system and its surroundings. The system is the part of universe which is under thermodynamic consideration. It is separated from the rest of the universe by real or imaginary boundaries.




Figure 7.1 System, surrounding & boundary Homogeneous and heterogeneous systems

Example:

The system may be water in a beaker, a balloon filled with air, an aqueous solution of glucose etc.

On the basis of physical and chemical properties, systems can be divided into two types.

A system is called homogeneous if the physical state of all its constituents are the same. Example: a mixture of gases, completely miscible mixture of liquids etc.

A system is called heterogeneous, if physical state of all its constituents is not the same.

Example: mixture of oil and water

Surrounding:

Everything in the universe that is not the part of the system is called surroundings.

Boundary:

Anything which separates the system from its surrounding is called boundary.

7.2.1 Types of systems:

There are three types of thermodynamic systems depending on the nature of the boundary.




Figure 7.2 Types of Systems

Isolated system:

A system which can exchange neither matter nor energy with its surroundings is called an isolated system. Here boundary is sealed and insulated. Hot water contained in a thermos flask, is an example for an isolated system. In this isolated system both energy (heat) and matter (water vapour) neither enter nor leave the system.

Closed system:

A system which can exchange only energy but not matter with its surroundings is called a closed system. Here the boundary is sealed but not insulated. Hot water contained in a closed beaker is an example for a closed system. In this system energy (heat) is transferred to the surroundings but no matter (water vapour) can escape from this system. A gas contained in a cylinder fitted with a piston constitutes a closed system.

Open system:

A System which can exchange both matter and energy with its surrounding is called an open system. Hot water contained in an open beaker is an example for open system. In this system both matter (water vapour) and energy (heat) is transferred to the surrounding.

All living things and chemical reactions are open systems because they exchange matter and energy with the surroundings.

7.2.2 Properties of the system:

Intensive and extensive properties

Some of the properties of a system depend on its mass or size whereas other properties do not depend on its mass or size. Based on this, the properties of a system are grouped as **extensive** property and **intensive** property.

Extensive properties:

The property that depends on the mass or the size of the system is called an extensive property.

Examples: Volume, Number of moles, Mass, Internal energy, etc.,

Intensive properties:

The property that is independent of the mass or the size of the system is called an intensive property.

Examples : Refractive index, Surface tension, density, temperature, Boiling point, Freezing point, molar volume, etc.,

Table: 7.1 Typical List of Extensive and Intensive properties

Extensive properties	Intensive properties
volume, mass, amount of substance (mole), energy, enthalpy, entropy, free energy, heat capacity	molar volume, density, molar mass, molarity, mole fraction, molality, specific heat capacity

7.2.3 Thermodynamic processes

The method of operation which can bring about a change in the system is called thermodynamic process. Heating, cooling, expansion, compression, fusion, vaporization etc., are some examples of a thermodynamic process.

Types of processes:

A thermodynamic process can be carried out in different ways and under different conditions. The processes can be classified as follows:

Reversible process:

The process in which the system and surrounding can be restored to the initial state from the final state without producing any changes in the thermodynamic properties of the universe is called a reversible process. There are two important conditions for the reversible process to occur. Firstly, the process should occur infinitesimally slowly and secondly throughout the process, the system and surroundings must be in equilibrium with each other.

Irreversible Process:

The process in which the system and surrounding cannot be restored to the initial state from the final state is called an irreversible process. All the processes occurring in nature are irreversible processes. During the irreversible process the system and surroundings are not in equilibrium with each other.

Adiabatic process:

An adiabatic process is defined as one in which there is no exchange of heat (q) between the system and surrounding during the process. Those processes in which no heat can flow into or out of the system are called **adiabatic processes**. This condition is attained by thermally insulating the system. In an adiabatic process if work is done by the system its temperature decreases, if work is done on the system its temperature increases, because, the system cannot exchange heat with its surroundings.

For an adiabatic process $q = 0$

Isothermal process :

An isothermal process is defined as one in which the temperature of the system remains constant, during the change from its initial to final state. The system exchanges heat with its surrounding and the temperature of the system remains constant. For this purpose the experiment is often performed in a thermostat.

For an isothermal process $dT = 0$

Isobaric process

An isobaric process is defined as one in which the pressure of the system remains constant during its change from the initial to final state.

For an isobaric process $dP = 0$.

Isochoric process

An isochoric process is defined as

the one in which the volume of system remains constant during its change from initial to final state. Combustion of a fuel in a bomb calorimeter is an example of an isochoric process.

For an isochoric process, $dV = 0$.

Cyclic process:

When a system returns to its original state after completing a series of changes, then it is said that a cycle is completed. This process is known as a cyclic process.

For a cyclic process $dU = 0$, $dH = 0$, $dP = 0$, $dV = 0$, $dT = 0$

Table: 7.2 Overview of the process and its condition

Process	Condition
Adiabatic	$q = 0$
Isothermal	$dT = 0$
Isobaric	$dP = 0$
Isochoric	$dV = 0$
Cyclic	$dE = 0$, $dH = 0$, $dP = 0$, $dV = 0$, $dT = 0$

State functions, path functions:

State function

A thermodynamic system can be defined by using the variables P , V , T and n . A state function is a thermodynamic property of a system, which has a specific value for a given state and does not depend on the path (or manner) by which the particular state is reached.

Example : Pressure (P), Volume (V), Temperature(T), Internal energy (U), Enthalpy (H), free energy (G) etc.

Path functions:

A path function is a thermodynamic property of the system whose value depends on the path by which the system changes from its initial to final states.

Example: Work (w), Heat (q).

Work (w) will have different values if the process is carried out reversibly or irreversibly.

Internal Energy (U)

The internal energy is a characteristic property of a system which is denoted by the symbol U. The internal energy of a system is equal to the energy possessed by all its constituents namely atoms, ions and molecules. The total energy of all molecules in a system is equal to the sum of their translational energy (U_t), vibrational energy (U_v), rotational energy (U_r), bond energy (U_b), electronic energy (U_e) and energy due to molecular interactions (U_i).

Thus:

$$U = U_t + U_v + U_r + U_b + U_e + U_i$$

The total energy of all the molecules of the system is called internal energy. In thermodynamics one is concerned only with the change in internal energy (ΔU) rather than the absolute value of energy.

Importance of Internal energy

The internal energy possessed by a substance differentiates its physical structure. For example, the allotropes of carbon, namely, graphite (C_{graphite}) and diamond (C_{diamond}), differ from each other because they possess different internal energies and have different structures.

Characteristics of internal energy (U):

- The internal energy of a system is an extensive property. It depends on the amount of the substances present in the system. If the amount is doubled, the internal energy is also doubled.
- The internal energy of a system is a state function. It depends only upon the state variables (T, P, V, n) of the system. The change in internal energy does not depend on the path by which the final state is reached.
- The change in internal energy of a system is expressed as $\Delta U = U_f - U_i$
- In a cyclic process, there is no internal energy change. $\Delta U_{(\text{cyclic})} = 0$
- If the internal energy of the system in the final state (U_f) is less than the internal energy of the system in its initial state (U_i), then ΔU would be negative.
$$\Delta U = U_f - U_i = -ve \quad (U_f < U_i)$$
- If the internal energy of the system in the final state (U_f) is greater than the internal energy of the system in its initial state (U_i), then ΔU would be positive.
$$\Delta U = U_f - U_i = +ve \quad (U_f > U_i)$$

HEAT(q)



The heat (q) is regarded as an energy in transit across the boundary separating a system from its surrounding. Heat changes lead to temperature differences between system and surrounding. Heat is a path function.

Units of heat:

The SI unit of heat is joule (J). Heat quantities are generally measured in calories (cal). A calorie is defined as the quantity of heat required to raise the temperature of 1 gram of water by 1°C in the vicinity of 15°C .

Sign convention of heat:

The symbol of heat is q .

If heat flows into the system from the surrounding, energy of a system increases. Hence it is taken to be positive ($+q$).

If heat flows out of the system into the surrounding, energy of the system decreases. Hence, it is taken to be negative ($-q$).

Work(w)

Work is defined as the force (F) multiplied by the displacement(x).

$$-w = F \cdot x \quad \text{-----}(7.1)$$

The negative sign ($-$) is introduced to indicate that the work has been done by the system by spending a part of its internal energy.

The work,

- (i) is a path function.
- (ii) appears only at the boundary of the system.
- (iii) appears during the change in the state of the system.
- (iv) In thermodynamics, surroundings is so large that macroscopic changes to surroundings do not happen.

Units of work:

The SI unit of work is joule (J), which is defined as the work done by a force of one Newton through a displacement of one meter ($J = \text{Nm}$). We often use kilojoule (kJ) for large quantities of work. $1 \text{ kJ} = 1000 \text{ J}$.

Sign convention of work:

The symbol of work is ' w '.

If work is done by the system, the energy of the system decreases, hence by convention, work is taken to be negative ($-w$).

If work is done on the system, the energy of the system increases, hence by convention, the work is taken to be positive ($+w$).

Pressure - volume work

In elementary thermodynamics the only type of work generally considered is the work done in expansion (or compression) of a gas. This is known as pressure-volume work, PV work or expansion work.

Work involved in expansion and

compression processes:

In most thermodynamic calculations we are dealing with the evaluation of work involved in the expansion or compression of gases. The essential condition for expansion or compression of a system is that there should be difference between external pressure (P_{ext}) and internal pressure (P_{int}).

For understanding pressure-volume work, let us consider a cylinder which contains 'n' moles of an ideal gas fitted with a frictionless piston of cross sectional area A . The total volume of the gas inside is V_i and pressure of the gas inside is P_{int} .

If the external pressure P_{ext} is greater than P_{int} , the piston moves inward till the pressure inside becomes equal to P_{ext} . Let this change be achieved in a single step and the final volume be V_f .

In this case, the work is done on the system (+w). It can be calculated as follows

$$w = -F \cdot \Delta x \quad \text{----- (7.2)}$$

where dx is the distance moved by the piston during the compression and F is the force acting on the gas.




Figure 7.3 showing work involved in compression processes

$$F = P_{\text{ext}} A \quad \text{---- (7.3)}$$

Substituting 7.3 in 7.2

$$w = -P_{\text{ext}} \cdot A \cdot \Delta x$$

$$A \cdot \Delta x = \text{change in volume} = V_f - V_i$$

$$w = -P_{\text{ext}} \cdot (V_f - V_i) \quad \text{----- (7.4)}$$

$$w = -P_{\text{ext}} \cdot (-\Delta V) \quad \text{----- (7.5)}$$

$$= P_{\text{ext}} \cdot \Delta V$$

Since work is done on the system, it is a positive quantity.

If the pressure is not constant, but changes during the process such that it is always infinitesimally greater than the pressure of the gas, then, at each stage of compression, the volume decreases by an infinitesimal amount, dV . In such a case we can calculate the work done on the gas by the relation

$$w_{rev} = - \int_{V_i}^{V_f} P_{ext} dV$$

In a compression process, P_{ext} the external pressure is always greater than the pressure of the system.

$$\text{i.e. } P_{ext} = (P_{int} + dP).$$

In an expansion process, the external pressure is always less than the pressure of the system

$$\text{i.e. } P_{ext} = (P_{int} - dP).$$




Figure 7.4

When pressure is not constant and changes in infinitesimally small steps (reversible conditions) during compression from V_i to V_f , the P-V plot looks like in fig 7.4 Work done on the gas is represented by the shaded area.

In general case we can write,

$P_{ext} = (P_{int} \pm dP)$. Such processes are called reversible processes. For a compression process work can be related to internal pressure of the system under reversible conditions by writing equation

$$w_{rev} = - \int_{V_i}^{V_f} P_{int} dV$$

For a given system with an ideal gas

$$P_{int} V = nRT$$

$$P_{int} = \frac{nRT}{V}$$

$$w_{rev} = - \int_{V_i}^{V_f} \frac{nRT}{V} dV$$

$$w_{rev} = -nRT \int_{V_i}^{V_f} \left(\frac{dV}{V} \right)$$

$$w_{rev} = -nRT \ln \left(\frac{V_f}{V_i} \right)$$

$$w_{rev} = -2.303 nRT \log \left(\frac{V_f}{V_i} \right) \text{ ----- 7.6}$$

If $V_f > V_i$ (expansion), the sign of work done by the process is negative.

If $V_f < V_i$ (compression) the sign of work done on the process is positive.

Table: 7.3 Summary of sign conventions

1.	If heat is absorbed by the system	:	$+q$
2.	If heat is evolved by the system	:	$-q$
3.	work is done by the system	:	$-w$
4.	work is done on the system	:	$+w$

7.3 Zeroth law of thermodynamics:

The zeroth law of thermodynamics, also known as the law of thermal equilibrium, was put forward much after the establishment of the first and second laws of thermodynamics. It is placed

before the first and second laws as it provides a logical basis for the concept of temperature of the system.




Figure : 7.5 Zeroth law of thermodynamics:

The law states that 'If two systems are separately in thermal equilibrium with a third one, then they tend to be in thermal equilibrium with themselves'.

According to this law, if systems B and C separately are in thermal equilibrium with another system A, then systems B and C will also be in thermal equilibrium with each other. This is also the principle by which thermometers are used.

7.4 First Law of Thermodynamics:

The first law of thermodynamics, known as the law of conservation of energy, states that the total energy of an isolated system remains constant though it may change from one form to another.

When a system moves from state 1 to state 2, its internal energy changes from U_1 to U_2 . Then change in internal energy

$$\Delta U = U_2 - U_1.$$

This internal energy change is brought about by the either absorption or evolution of heat and/or by work being done by/on the system.

Because the total energy of the system must remain constant, we can write the mathematical statement of the First Law as:

$$\Delta U = q + w \text{ -----(7.7)}$$

Where q - the amount of heat supplied to the system; w - work done on the system

Other statements of first law of thermodynamics

- (1) Whenever an energy of a particular type disappears, an equivalent amount of another type must be produced.
- (2) The total energy of a system and surrounding remains constant (or conserved)
- (3) "Energy can neither be created nor destroyed, but may be converted from one form to another".
- (4) "The change in the internal energy of a closed system is equal to the energy that passes through its boundary as heat or work".
- (5) "Heat and work are two ways of changing a system's internal energy".

7.4.1 Mathematical statement of the first law

The mathematical statement of the first law of thermodynamics is

$$\Delta U = q + w \text{ ----- (7.7)}$$

Case 1 : For a cyclic process involving isothermal expansion of an ideal gas,

$$\Delta U = 0.$$

$$\text{eqn (7.7)} \Rightarrow \therefore q = -w$$

In other words, during a cyclic process, the amount of heat absorbed by the system is equal to work done by the system.

Case 2 : For an isochoric process (no change in volume) there is no work of expansion. i.e. $\Delta V = 0$

$$\begin{aligned}\Delta U &= q + w \\ &= q - P\Delta V \\ \Delta V &= 0 \\ \Delta U &= q_v\end{aligned}$$

In other words, during an isochoric process, the amount of heat supplied to the system is converted to its internal energy.

Case 3 : For an adiabatic process there is no change in heat. i.e. $q = 0$. Hence

$$q = 0$$

$$\text{eqn (7.7)} \Rightarrow \Delta U = w$$

In other words, in an adiabatic process, the decrease in internal energy

is exactly equal to the work done by the system on its surroundings.

Case 4 : For an isobaric process. There is no change in the pressure. P remains constant. Hence

$$\begin{aligned}\Delta U &= q + w \\ \Delta U &= q - P\Delta V\end{aligned}$$

In other words, in an isobaric process a part of heat absorbed by the system is used for P - V expansion work and the remaining is added to the internal energy of the system.

Problem: 7.1

A gas contained in a cylinder fitted with a frictionless piston expands against a constant external pressure of 1 atm from a volume of 5 litres to a volume of 10 litres. In doing so it absorbs 400 J of thermal energy from its surroundings. Determine the change in internal energy of system.

Solution:

$$\text{Given data } q = 400 \text{ J } V_1 = 5 \text{ L } V_2 = 10 \text{ L}$$

$\Delta u = q - w$ (heat is given to the system (+q); work is done by the system (-w))

$$\begin{aligned}\Delta u &= q - PdV \\ &= 400 \text{ J} - 1 \text{ atm } (10 - 5) \text{ L} \\ &= 400 \text{ J} - 5 \text{ atm L} \\ &[\therefore 1 \text{ L atm} = 101.33 \text{ J}] \\ &= 400 \text{ J} - 5 \times 101.33 \text{ J} \\ &= 400 \text{ J} - 506.65 \text{ J} \\ &= -106.65 \text{ J}\end{aligned}$$

7.5 Enthalpy (H)

The enthalpy (H), is a thermodynamic property of a system, is defined as the sum of the internal energy (U) of a system and the product of pressure and volume of the system. That is,

$$H = U + PV \text{ -----(7.8)}$$

It reflects the capacity to do mechanical work and the capacity to release heat by the system. When a process occurs at constant pressure, the heat involved (either released or absorbed) is equal to the change in enthalpy.

Enthalpy is a state function which depends entirely on the state functions T, P and U. Enthalpy is usually expressed as the change in enthalpy (ΔH) for a process between initial and final states at constant pressure.

$$\Delta H = \Delta U + P\Delta V \text{ -----(7.9)}$$

The change in enthalpy (ΔH) is equal to the heat supplied at the constant pressure to a system (as long as the system does no additional work).

$$\Delta H = q_p$$

In an endothermic reaction heat is absorbed by the system from the surroundings that is $q > 0$ (positive). Therefore, ΔH is also positive. In an exothermic reaction heat is evolved by the system to the surroundings that is, $q < 0$ (negative). If q is negative, then ΔH will also be negative.

7.5.1 Relation between enthalpy 'H' and internal energy 'U'

When the system at constant pressure undergoes changes from an initial state with H_1 , U_1 and V_1 to a final state with H_2 , U_2 and V_2 the change in enthalpy ΔH , can be calculated as follows:

$$H = U + PV$$

In the initial state

$$H_1 = U_1 + PV_1 \text{ -----(7.10)}$$

In the final state

$$H_2 = U_2 + PV_2 \text{ -----(7.11)}$$

change in enthalpy is (7.11) - (7.10)

$$(H_2 - H_1) = (U_2 - U_1) + P(V_2 - V_1)$$

$$\Delta H = \Delta U + P\Delta V \text{ ----- (7.12)}$$

As per first law of thermodynamics,

$$\Delta U = q + w$$

Equation 7.12 becomes

$$\Delta H = q + w + P\Delta V$$

$$w = -P\Delta V$$

$$\Delta H = q_p - P\Delta V + P\Delta V$$

$$\Delta H = q_p \text{ ----- (7.13)}$$

q_p – is the heat absorbed at constant pressure and is considered as heat content.

Consider a closed system of gases which are chemically reacting to form gaseous products at constant temperature

and pressure with V_i and V_f as the total volumes of the reactant and product gases respectively, and n_i and n_f as the number of moles of gaseous reactants and products, then,

For reactants (initial state) :

$$PV_i = n_i RT \text{ ----- (7.14)}$$

For products (final state) :

$$PV_f = n_f RT \text{ ----- (7.15)}$$

(7.15) - (7.14)

$$P (V_f - V_i) = (n_f - n_i) RT$$

$$P \Delta V = \Delta n_{(g)} RT \text{ ----- (7.16)}$$

Substituting in 7.16 in 7.12


$$\Delta H = \Delta U + \Delta n_{(g)} RT \text{ ----- (7.17)}$$

7.5.2 Enthalpy Changes for Different Types of Reactions and Phase Transitions:


The heat or enthalpy changes accompanying chemical reactions is expressed in different ways depending on the nature of the reaction. These are discussed below.

Standard heat of formation


The standard heat of formation of a compound is defined as "the change in enthalpy that takes place when one mole of a compound is formed from its elements, present in their standard states (298 K and 1 bar pressure)". By convention the standard heat of formation of all elements is assigned a value of zero.



$$\Delta H_f^0 = -100.42 \text{ kJ mol}^{-1}$$



$$\Delta H_f^0 = +222.33 \text{ kJ mol}^{-1}$$



$$\Delta H_f^0 = -92.4 \text{ kJ mol}^{-1}$$

The standard heats of formation of some compounds are given in Table 7.4.

Table: 7.4 standard heat of formation of some compounds

Substance	$\Delta H_f^0 \text{ (kJ mol}^{-1}\text{)}$	Substance	$\Delta H_f^0 \text{ (kJ mol}^{-1}\text{)}$
$\text{H}_2\text{O(l)}$	-242	$\text{CH}_4\text{(g)}$	-74.85
HCl(g)	-92.4	$\text{C}_2\text{H}_6\text{(g)}$	-84.6
HBr(g)	-36.4	$\text{C}_6\text{H}_6\text{(g)}$	+49.6
$\text{NH}_3\text{(g)}$	-46.1	$\text{C}_2\text{H}_2\text{(g)}$	+222.33
$\text{CO}_2\text{(g)}$	-393.5	$\text{CH}_3\text{OH(l)}$	-239.2


7.6 Thermochemical Equations:

A thermochemical equation is a balanced stoichiometric chemical equation that includes the enthalpy change (ΔH). The following conventions are adopted in thermochemical equations:


- The coefficients in a balanced thermochemical equation refer to number of moles of reactants and products involved in the reaction.

- (ii) The enthalpy change of the reaction ΔH_r has to be specified with appropriate sign and unit.
- (iii) When the chemical reaction is reversed, the value of ΔH is reversed in sign with the same magnitude.
- (iv) The physical states (gas, liquid, aqueous, solid in brackets) of all species are important and must be specified in a thermochemical reaction, since ΔH depends on the physical state of reactants and products.
- (v) If the thermochemical equation is multiplied throughout by a number, the enthalpy change is also multiplied by the same number.
- (vi) The negative sign of ΔH_r indicates that the reaction is exothermic and the positive sign of ΔH_r indicates an endothermic reaction.

For example, consider the following reaction,



$$\Delta H_r^0 = -967.4 \text{ kJ}$$



$$\Delta H_r^0 = +967.4 \text{ kJ}$$

Standard enthalpy of reaction (ΔH_r^0) from standard enthalpy of formation (ΔH_f^0)


The standard enthalpy of a reaction is the enthalpy change for a reaction

when all the reactants and products are present in their standard states. Standard conditions are denoted by adding the superscript 0 to the symbol (ΔH^0)

We can calculate the enthalpy of a reaction under standard conditions from the values of standard enthalpies of formation of various reactants and products. The standard enthalpy of reaction is equal to the difference between standard enthalpy of formation of products and the standard enthalpies of formation of reactants.

$$\Delta H_r^0 = \sum \Delta H_f^0 (\text{products}) - \sum \Delta H_f^0 (\text{reactants})$$

For a general reaction




$$\Delta H_r^0 = \sum \Delta H_f^0 (\text{products}) - \sum \Delta H_f^0 (\text{reactants})$$

$$\Delta H_r^0 = \{c \Delta H_f^0 (\text{C}) + d \Delta H_f^0 (\text{D})\} - \{a \Delta H_f^0 (\text{A}) + b \Delta H_f^0 (\text{B})\}$$

Problem : 7.2

The standard enthalpies of formation of $\text{C}_2\text{H}_5\text{OH}(\text{l})$, $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\text{l})$ are -277, -393.5 and -285.5 kJ mol^{-1} respectively. Calculate the standard enthalpy change for the reaction




The enthalpy of formation of $\text{O}_2(\text{g})$ in the standard state is Zero, by definition

Solution:

For example, the standard enthalpy change for the combustion of ethanol can be calculated from the standard enthalpies of formation of $\text{C}_2\text{H}_5\text{OH}(\text{l})$, $\text{CO}_2(\text{g})$

and $\text{H}_2\text{O}(\text{l})$. The enthalpies of formation are -277 , -393.5 and $-285.5 \text{ kJ mol}^{-1}$ respectively.




$$\begin{aligned}\Delta H_r^0 &= \left\{ (\Delta H_f^0)_{\text{products}} - (\Delta H_f^0)_{\text{reactants}} \right\} \\ \Delta H_r^0 &= \left[2(\Delta H_f^0)_{\text{CO}_2} + 3(\Delta H_f^0)_{\text{H}_2\text{O}} \right] \\ &\quad - \left[1(\Delta H_f^0)_{\text{C}_2\text{H}_5\text{OH}} + 3(\Delta H_f^0)_{\text{O}_2} \right] \\ \Delta H_r^0 &= \left[2\text{mol}(-393.5) \text{ kJ mol}^{-1} \right] \\ &\quad + \left[3\text{mol}(-285.5) \text{ kJ mol}^{-1} \right] \\ &\quad - \left[1\text{mol}(-277) \text{ kJ mol}^{-1} \right] \\ &\quad + \left[3\text{mol}(0) \text{ kJ mol}^{-1} \right] \\ &= [-787 - 856.5] - [-277] \\ &= -1643.5 + 277\end{aligned}$$

$$\Delta H_r^0 = -1366.5 \text{ KJ}$$

Evaluate Yourself - 1

Calculate ΔH_f^0 for the reaction




given that ΔH_f^0 for $\text{CO}_2(\text{g})$, $\text{CO}(\text{g})$ and $\text{H}_2\text{O}(\text{g})$ are -393.5 , -111.31 and -242 kJ mol^{-1} respectively.

Heat of combustion


The heat of combustion of a substance is defined as “The change in enthalpy of a system when one mole of the substance is completely burnt in excess of air or oxygen”. It is denoted by ΔH_C . For example, the heat of combustion of

methane is $-87.78 \text{ kJ mol}^{-1}$



$$\Delta H_C = -87.78 \text{ kJ mol}^{-1}$$

For the combustion of carbon,



$$\Delta H_C = -394.55 \text{ kJ mol}^{-1}$$

Combustion reactions are always exothermic. Hence the enthalpy change is always negative.

Molar heat capacities

When heat (q) is supplied to a system, the molecules in the system absorb the heat and hence their kinetic energy increases, which in turn raises the temperature of the system from T_1 to T_2 . This increase ($T_2 - T_1$) in temperature is directly proportional to the amount of heat absorbed and inversely proportional to mass of the substance. In other words,

$$q \propto m\Delta T$$

$$q = c m \Delta T$$

$$c = q/m \Delta T$$

The constant c is called heat capacity.

$$c = \left(\frac{q}{m(T_2 - T_1)} \right) \text{ ----- (7.18)}$$

when $m=1 \text{ kg}$ and $(T_2 - T_1) = 1 \text{ K}$ then the heat capacity is referred as specific heat capacity. The equation 7.18 becomes

$$c = q$$



Thus specific heat capacity of a system is defined as “The heat absorbed by one kilogram of a substance to raise its temperature by one Kelvin at a specified temperature”.

The heat capacity for 1 mole of substance, is called molar heat capacity (c_m). It is defined as “The amount of heat absorbed by one mole of the substance to raise its temperature by 1 kelvin”.

Units of Heat Capacity:

The SI unit of molar heat capacity is $\text{JK}^{-1}\text{mol}^{-1}$

The molar heat capacities can be expressed either at constant volume (C_v) or at constant pressure (C_p).

According to the first law of thermodynamics

$$U = q + w \quad \text{or} \quad U = q - PdV$$

$$q = U + PdV \text{ -----(7.19)}$$

Differentiate (7.19) with respect to temperature at constant volume i.e $dV=0$,

$$\left(\frac{\partial q}{\partial T}\right)_v = \left(\frac{\partial U}{\partial T}\right)_v$$

$$C_v = \left(\frac{\partial U}{\partial T}\right)_v \text{ -----7.20}$$

Thus the heat capacity at constant volume (C_v) is defined as the rate of change of internal energy with respect to temperature at constant volume.

Similarly the molar heat capacity at constant pressure (C_p) can be defined as

the rate of change of enthalpy with respect to temperature at constant pressure.

$$c_p = \left(\frac{\partial H}{\partial T}\right)_p \text{ ----- (7.21)}$$

Relation between C_p and C_v for an ideal gas.

From the definition of enthalpy

$$H = U + PV \text{ -----(7.8)}$$

for 1 mole of an ideal gas

$$PV = nRT \text{ -----(7.22)}$$

By substituting (7.22) in (7.8)

$$H = U + nRT \text{ ---- (7.23)}$$

Differentiating the above equation with respect to T,

$$\frac{\partial H}{\partial T} = \frac{\partial U}{\partial T} + nR \frac{\partial T}{\partial T}$$

$$C_p = C_v + nR \quad (1) \quad \left[\begin{array}{l} \because \left(\frac{\partial H}{\partial T}\right)_P = C_P \\ \text{and } \left(\frac{\partial U}{\partial T}\right)_V = C_V \end{array} \right]$$

$$C_p - C_v = nR \text{(7.24)}$$

At constant pressure processes, a system has to do work against the surroundings. Hence, the system would require more heat to effect a given temperature rise than at constant volume, so C_p is always greater than C_v

Calculation of ΔU and ΔH

For one mole of an ideal gas, we have



$$C_V = \frac{dU}{dT}$$

$$dU = C_V dT$$

For a finite change, we have

$$\Delta U = C_V \Delta T$$

$$\Delta U = C_V (T_2 - T_1)$$

and for n moles of an ideal gas we get

$$\Delta U = n C_V (T_2 - T_1) \text{ ----- (7.25)}$$

Similarly for n moles of an ideal gas we get

$$\Delta H = n C_P (T_2 - T_1) \text{ ----- (7.26)}$$

Problem 7.3

Calculate the value of ΔU and ΔH on heating 128.0 g of oxygen from 0°C to 100°C . C_V and C_P on an average are 21 and 29 $\text{J mol}^{-1} \text{K}^{-1}$. (The difference is $8 \text{J mol}^{-1} \text{K}^{-1}$ which is approximately equal to R)

Solution.

We know

$$\Delta U = n C_V (T_2 - T_1)$$

$$\Delta H = n C_P (T_2 - T_1)$$

$$\text{Here } n = \frac{128}{32} = 4 \text{ moles ;}$$

$$T_2 = 100^\circ \text{C} = 373 \text{K}; T_1 = 0^\circ \text{C} = 273 \text{K}$$

$$\Delta U = n C_V (T_2 - T_1)$$

$$\Delta U = 4 \times 21 \times (373 - 273)$$

$$\Delta U = 8400 \text{ J}$$

$$\Delta U = 8.4 \text{ kJ}$$

$$\Delta H = n C_P (T_2 - T_1)$$

$$\Delta H = 4 \times 29 \times (373 - 273)$$

$$\Delta H = 11600 \text{ J}$$

$$\Delta H = 11.6 \text{ kJ}$$



Evaluate Yourself -2

Calculate the amount of heat necessary to raise 180 g of water from 25°C to 100°C . Molar heat capacity of water is $75.3 \text{ J mol}^{-1} \text{K}^{-1}$

7.7 Measurement of ΔU and ΔH using Calorimetry.

Calorimeter is used for measuring the amount of heat change in a chemical or physical change. In calorimetry, the temperature change in the process is measured which is inversely proportional to the heat change. By using the expression $C = q/m\Delta T$, we can calculate the amount of heat change in the process. Calorimetric measurements are made under two different conditions

i) At constant volume (q_V)

ii) At constant pressure (q_P)

(A) ΔU Measurements

For chemical reactions, heat evolved at constant volume, is measured in a *bomb calorimeter*.

The inner vessel (the bomb) and its cover are made of strong steel. The cover is fitted tightly to the vessel by means of metal lid and screws.




Figure 7.6 Bomb calorimeter

A weighed amount of the substance is taken in a platinum cup connected with

electrical wires for striking an arc instantly to kindle combustion. The bomb is then tightly closed and pressurized with excess oxygen. The bomb is immersed in water, in the inner volume of the calorimeter. A stirrer is placed in the space between the wall of the calorimeter and the bomb, so that water can be stirred, uniformly. The reaction is started by striking the substance through electrical heating.

A known amount of combustible substance is burnt in oxygen in the bomb. Heat evolved during the reaction is absorbed by the calorimeter as well as the water in which the bomb is immersed. The change in temperature is measured using a Beckman thermometer. Since the bomb is sealed its volume does not change and hence the heat measurements is equal to the heat of combustion at a constant volume (ΔU_c^0).

The amount of heat produced in the reaction (ΔU_c^0) is equal to the sum of the heat absorbed by the calorimeter and water.

Heat absorbed by the calorimeter

$$q_1 = k \cdot \Delta T$$

where k is a calorimeter constant equal to $m_c C_c$ (m_c is mass of the calorimeter and C_c is heat capacity of calorimeter)

Heat absorbed by the water

$$q_2 = m_w C_w \Delta T$$

where m_w is molar mass of water

C_w is molar heat capacity of water ($75.29 \text{ J K}^{-1} \text{ mol}^{-1}$)

$$\begin{aligned}
 \text{Therefore } \Delta U_c &= q_1 + q_2 \\
 &= k \cdot \Delta T + m_w C_w \Delta T \\
 &= (k + m_w C_w) \Delta T
 \end{aligned}$$

Calorimeter constant can be determined by burning a known mass of standard sample (benzoic acid) for which the heat of combustion is known ($-3227 \text{ kJ mol}^{-1}$)

The enthalpy of combustion at constant pressure of the substance is calculated from the equation (7.17)

$$\Delta H_{C(\text{pressure})}^0 = \Delta U_{C(\text{Vol})}^0 + \Delta n_g RT$$

Applications of bomb calorimeter:

1. Bomb calorimeter is used to determine the amount of heat released in combustion reaction.
2. It is used to determine the calorific value of food.
3. Bomb calorimeter is used in many industries such as metabolic study, food processing, explosive testing etc.

(b) ΔH Measurements

Heat change at constant pressure (at atmospheric pressure) can be measured using a coffee cup calorimeter. A schematic representation of a coffee cup calorimeter is given in Figure 7.7. Instead of bomb, a styrofoam cup is used in this calorimeter. It acts as good adiabatic wall and doesn't allow transfer of heat produced during the reaction to its surrounding. This entire heat energy is absorbed by the water inside the cup. This method can be used for the reactions where there is no

appreciable change in volume. The change in the temperature of water is measured and used to calculate the amount of heat that has been absorbed or evolved in the reaction using the following expression.

$$q = m_w C_w \Delta T$$

where m_w is the molar mass of water and C_w is the molar heat capacity of water ($75.29 \text{ J K}^{-1} \text{ mol}^{-1}$)





Figure 7.7 Coffee cup Calorimeter

Problem 7. 4

Calculate the enthalpy of combustion of ethylene at 300 K at constant pressure, if its heat of combustion at constant volume (ΔU) is -1406 kJ .

The complete ethylene combustion reaction can be written as,



$$\Delta U = -1406 \text{ kJ}$$

$$\Delta n = n_{\text{p}(\text{g})} - n_{\text{r}(\text{g})}$$

$$\Delta n = 2 - 4 = -2$$



$$\Delta H = \Delta U + RT\Delta n_{\text{g}}$$

$$\Delta H = -1406 + (8.314 \times 10^{-3} \times 300 \times (-2))$$

$$\Delta H = -1410.9 \text{ kJ}$$

Evaluate Yourself - 3

From the following data at constant volume for combustion of benzene, calculate the heat of this reaction at constant pressure condition.



$$\Delta U \text{ at } 25^\circ \text{C} = -3268.12 \text{ kJ}$$

Applications of the heat of combustion:

(1) Calculation of heat of formation: Since the heat of combustion of organic compounds can be determined with considerable ease, they are employed to calculate the heat of formation of other compounds.


For example let us calculate the standard enthalpy of formation ΔH_f^0 of CH_4 from the values of enthalpy of combustion for H_2 , $\text{C}(\text{graphite})$ and CH_4 which are -285.8, -393.5, and -890.4 kJ mol^{-1} respectively.

Let us interpret the information about enthalpy of formation by writing out the equations. It is important to note that the standard enthalpy of formation of pure elemental gases and elements is assumed to be zero under standard conditions. Thermochemical equation for the formation of methane from its constituent elements is,




$$\Delta H_f^0 = X \text{ kJ mol}^{-1} \text{--- (i)}$$


Thermo chemical equations for the combustion of given substances are,



$$\Delta H^0 = -285.8 \text{ kJ mol}^{-1} \text{--- (ii)}$$




$$\Delta H^0 = -393.5 \text{ kJ mol}^{-1} \text{--- (iii)}$$



$$\Delta H^0 = -890.4 \text{ kJ mol}^{-1} \text{--- (iv)}$$

Since methane is in the product side of the required equation (i), we have to reverse the equation (iv)



$$\Delta H^0 = +890.4 \text{ kJ mol}^{-1} \text{--- (v)}$$

In order to get equation (i) from the remaining,

$$(i) = [(ii) \times 2] + (iii) + (v)$$

$$X = [(-285.8) \times 2] + [-393.5] + [+890.4]$$

$$= -74.7 \text{ kJ}$$

Hence, the amount of energy required for the formation of 1 mole of methane is -74.7 kJ

The heat of formation methane = -74.7 kJ mol⁻¹


(2) Calculation of calorific value of food and fuels: The calorific value is defined as the amount of heat produced in calories (or joules) when one gram of the substance is completely burnt. The SI unit of calorific value is J kg⁻¹. However, it is usually expressed in cal g⁻¹.

Heat of solution:


Heat changes are usually observed when a substance is dissolved in a solvent. The heat of solution is defined as the change in enthalpy when one mole of a substance is dissolved in a specified quantity of solvent at a given temperature.

Heat of neutralisation:

The heat of neutralisation is defined as “The change in enthalpy when one gram equivalent of an acid is completely neutralised by one gram equivalent of a base or vice versa in dilute solution”.




$$\Delta H = -57.32 \text{ kJ}$$




$$\Delta H = -57.32 \text{ kJ}$$

The heat of neutralisation of a strong acid and strong base is around -57.32 kJ, irrespective of nature of acid or base used which is evident from the below



mentioned examples.



$$\Delta H = -57.32 \text{ kJ}$$




$$\Delta H = -57.32 \text{ kJ}$$



$$\Delta H = -57.32 \text{ kJ} \times 2$$

The reason for this can be explained on the basis of Arrhenius theory of acids and bases which states that strong acids and strong bases completely ionise in aqueous solution to produce H⁺ and OH⁻ ions respectively. Therefore in all the above mentioned reactions the neutralisation can be expressed as follows.




$$\Delta H = -57.32 \text{ kJ}$$

Molar heat of fusion

The molar heat of fusion is defined as “the change in enthalpy when one mole of a solid substance is converted into the liquid state at its melting point”.

For example, the heat of fusion of ice can be represented as




Molar heat of vapourisation

The molar heat of vaporisation is defined as “the change in enthalpy when

one mole of liquid is converted into vapour state at its boiling point”.

For example, heat of vaporisation of water can be represented as



Molar heat of sublimation


Sublimation is a process when a solid changes directly into its vapour state without changing into liquid state. Molar heat of sublimation is defined as “the change in enthalpy when one mole of a solid is directly converted into the vapour state at its sublimation temperature”. For example, the heat of sublimation of iodine is represented as



Another example of sublimation process is solid CO_2 to gas at atmospheric pressure at very low temperatures.

Heat of transition


The heat of transition is defined as “The change in enthalpy when one mole of an element changes from one of its allotropic form to another. For example, the transition of diamond into graphite may be represented as



Similarly the allotropic transitions in sulphur and phosphorous can be represented as follows,



$$\Delta H_{\text{trans}} = - 0.067 \text{ kJ}$$




7.8 Hess's law of constant heat summation

We have already seen that the heat changes in chemical reactions are equal to the difference in internal energy (ΔU) or heat content (ΔH) of the products and reactants, depending upon whether the reaction is studied at constant volume or constant pressure. Since ΔU and ΔH are functions of the state of the system, the heat evolved or absorbed in a given reaction depends only on the initial state and final state of the system and not on the path or the steps by which the change takes place.

This generalisation is known as **Hess's law** and stated as:

The enthalpy change of a reaction either at constant volume or constant pressure is the same whether it takes place in a single or multiple steps provided the initial and final states are same.




Application of Hess's Law: Hess's law can be applied to calculate enthalpies of reactions that are difficult to measure. For example, it is very difficult to measure the heat of combustion of graphite to give pure CO .

However, enthalpy for the oxidation of graphite to CO_2 and CO to CO_2 can easily be measured. For these conversions, the heat of combustion values are -393.5 kJ and -283.5 kJ respectively.

From these data the enthalpy of combustion of graphite to CO can be calculated by applying Hess's law.

The reactions involved in this process can be expressed as follows




According to Hess law,

$$\begin{aligned}
 \Delta H_1 &= \Delta H_2 + \Delta H_3 \\
 -393.5 \text{ kJ} &= X - 283.5 \text{ kJ} \\
 X &= -110.5 \text{ kJ}
 \end{aligned}$$

7.9 Lattice energy ($\Delta H_{\text{lattice}}$)

Lattice energy is defined as the amount of energy required to completely remove the constituent ions from its crystal lattice to an infinite distance from one mole of crystal. It is also referred as lattice enthalpy.




From the above equation it is clear that 788 kJ of energy is required to separate Na^+ and Cl^- ions from 1 mole of NaCl .

Born - Haber cycle

The Born-Haber cycle is an approach to analyse reaction energies. It was named after two German scientists Max Born and Fritz Haber who developed this cycle. The cycle is concerned with the formation of an ionic compound from the reaction of a metal with a halogen or other non-metallic element such as oxygen.

Born-Haber cycle is primarily used in calculating lattice energy, which cannot be measured directly. The Born-Haber cycle applies Hess's law to calculate the lattice enthalpy. For example consider the formation of a simple ionic solid such as an alkali metal halide MX , the following steps are considered.



ΔH_1 - enthalpy change for the sublimation M(s) to M(g)

ΔH_2 - enthalpy change $\frac{1}{2} \text{X}_2(\text{g})$ to X(g) for the dissociation of

ΔH_3 - Ionisation energy for M(g) to $\text{M}^+(\text{g})$

ΔH_4 - electron affinity for the conversion of X(g) to $\text{X}^-(\text{g})$

U - the lattice enthalpy for the formation of solid MX

ΔH_f - enthalpy change for the formation of solid MX directly from elements


According to Hess's law of heat summation

$$\Delta H_f = \Delta H_1 + \Delta H_2 + \Delta H_3 + \Delta H_4 + U$$

Let us use the Born - Haber cycle for determining the lattice enthalpy of NaCl as follows:

Since the reaction is carried out with reactants in elemental forms and products in their standard states, at 1 bar, the overall enthalpy change of the reaction is also the enthalpy of formation for NaCl. Also, the formation of NaCl can be considered in 5 steps. The sum of the enthalpy changes of these steps is equal to the enthalpy change for the overall reaction from which the lattice enthalpy of NaCl is calculated.

Let us calculate the lattice energy of sodium chloride using Born-Haber cycle



ΔH_f = heat of formation of sodium chloride = $-411.3 \text{ kJ mol}^{-1}$

ΔH_1 = heat of sublimation of Na(g) = $108.7 \text{ kJ mol}^{-1}$

ΔH_2 = ionisation energy of Na(g) = $495.0 \text{ kJ mol}^{-1}$

ΔH_3 = dissociation energy of $\text{Cl}_2(\text{g})$ = 244 kJ mol^{-1}

ΔH_4 = Electron affinity of Cl(S) = $-349.0 \text{ kJ mol}^{-1}$

U = lattice energy of NaCl

$$\Delta H_f = \Delta H_1 + \Delta H_2 + \frac{1}{2} \Delta H_3 + \Delta H_4 + U$$

$$\therefore U = (\Delta H_f) - \left(\Delta H_1 + \Delta H_2 + \frac{1}{2} \Delta H_3 + \Delta H_4 \right)$$

$$\Rightarrow U = (-411.3) - (108.7 + 495.0 + 122 - 349)$$

$$U = (-411.3) - (376.7)$$

$$\therefore U = -788 \text{ kJ mol}^{-1}$$

This negative sign in lattice energy indicates that the energy is released when sodium is formed from its constituent gaseous ions Na^+ and Cl^-

Evaluate Yourself - 4



When a mole of magnesium bromide is prepared from 1 mole of magnesium and 1 mole of liquid bromine, 524 kJ of energy is released. The heat of sublimation of Mg metal is 148 kJ mol^{-1} . The heat of dissociation of bromine gas into atoms is 193 kJ mol^{-1} . The heat of vapourisation of liquid bromine is 31 kJ mol^{-1} . The first and second ionisation energies of magnesium are $\text{IE}_1 = 737.5 \text{ kJ mol}^{-1}$ and $\text{IE}_2 = 1450.5 \text{ kJ mol}^{-1}$ and the electron affinity of bromine is $-324.5 \text{ kJ mol}^{-1}$. Calculate the lattice energy of magnesium bromide.

7.10. Second Law of thermodynamics:

Need for the second law of thermodynamics:

We know from the first law of thermodynamics, the energy of the universe is conserved. Let us consider the following processes:



1. A glass of hot water over time loses heat energy to the surrounding and becomes cold.
2. When you mix hydrochloric acid with sodium hydroxide, it forms sodium chloride and water with evolution of heat.

In both these processes, the total energy is conserved and are consistent with the first law of thermodynamics. However, the reverse process i.e. cold water becoming hot water by absorbing heat from surrounding on its own does not occur spontaneously even though the energy change involved in this process is also consistent with the first law. However, if the heat energy is supplied to cold water, then it will become hot. i.e. the change that does not occur spontaneously and can be driven by supplying energy.

Similarly, a solution of sodium chloride does not absorb heat energy on its own, to form hydrochloric acid and sodium hydroxide. But, this process can not be driven even by supplying energy. From these kinds of our natural experiences, we have come to know that certain processes are spontaneous while the others are not, and some processes have a preferred direction. In order to explain the feasibility of a process, we need the second law of thermodynamics.

7.10.1 Various statements of the second law of thermodynamics

Entropy

The second law of thermodynamics introduces another state function called entropy. Entropy is a measure of the

molecular disorder (randomness) of a system. But thermodynamic definition of entropy is concerned with the change in entropy that occurs as a result of a process.

It is defined as, $dS = dq_{\text{rev}} / T$

Entropy statement:

The second law of thermodynamics can be expressed in terms of entropy. i.e. “the entropy of an isolated system increases during a spontaneous process”.

For an irreversible process such as spontaneous expansion of a gas,

$$\Delta S_{\text{total}} > 0$$

$$\Delta S_{\text{total}} > \Delta S_{\text{system}} + \Delta S_{\text{surrounding}}$$

$$\text{i.e. } \Delta S_{\text{total}} > \Delta S_{\text{system}} + \Delta S_{\text{surrounding}}$$

For a reversible process such as melting of ice,

$$\Delta S_{\text{system}} = - \Delta S_{\text{surrounding}}$$

$$\Delta S_{\text{total}} = 0$$

Kelvin-Planck statement:

It is impossible to construct a machine that absorbs heat from a hot source and converts it completely into work by a cyclic process without transferring a part of heat to a cold sink. The second law of thermodynamics explains why even an ideal, frictionless engine cannot convert 100% of its input heat into work. Carnot on his analysis of heat engines, found that the maximum efficiency of a heat engine which operates reversibly, depends only on the two temperatures between which it is operated.

Efficiency = work performed / heat absorbed

$$\eta = \frac{|q_h| - |q_c|}{|q_h|}$$

q_h - heat absorbed from the hot reservoir

q_c - heat transferred to cold reservoir

$$\eta = 1 - \frac{|q_c|}{|q_h|} \quad \text{--- (7.27)}$$

For a reversible cyclic process

$$\Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}} = 0$$

$$\Delta S_{\text{system}} = -\Delta S_{\text{surroundings}}$$

$$\frac{q_h}{T_h} = \frac{-q_c}{T_c}$$

$$\frac{T_c}{T_h} = \frac{-q_c}{q_h}$$

$$\frac{T_c}{T_h} = \frac{|q_c|}{|q_h|} \quad \text{--- (7.28)}$$

Substituting 7.28 in 7.27

$$\Rightarrow \eta = 1 - \frac{T_c}{T_h} \quad \text{--- (7.29)}$$

$$T_h \gg T_c$$

Hence, $\eta < 1$

efficiency in percentage can be expressed as

$$\text{Efficiency in percentage} = \left[1 - \frac{T_c}{T_h} \right] \times 100$$

Clausius statement:

It is impossible to transfer heat from a cold reservoir to a hot reservoir without doing some work.

Problem: 7.10

If an automobile engine burns petrol at a temperature of 816°C and if the surrounding temperature is 21°C , calculate its maximum possible efficiency.

Solution:

$$\% \text{ Efficiency} = \left[\frac{T_h - T_c}{T_h} \right] \times 100$$

Here

$$T_h = 816 + 273 = 1089 \text{ K};$$

$$T_c = 21 + 273 = 294 \text{ K}$$

$$\% \text{ Efficiency} = \left(\frac{1089 - 294}{1089} \right) \times 100$$

$$\% \text{ Efficiency} = 73\%$$

Evaluate Yourself - 5



An engine operating between 127°C and 47°C takes some specified amount of heat from a high temperature reservoir. Assuming that there are no frictional losses, calculate the percentage efficiency of an engine.

Unit of entropy:

The entropy (S) is equal to heat energy exchanged (q) divided by the temperature (T) at which the exchange takes place. Therefore, The SI unit of entropy is JK^{-1} .

Spontaneity and Randomness

Careful examination shows that in each of the processes viz., melting of ice and evaporation of water, there is an increase in randomness or disorder of the system. The water molecules in ice

are arranged in a highly organised crystal pattern which permits little movement. As the ice melts, the water molecules become disorganised and can move more freely. The movement of molecules becomes freer in the liquid phase and even more free in the vapour phase. In other words, we can say that the randomness of the water molecules increases, as ice melts into water or water evaporates. Both are spontaneous processes which result in an increase in randomness (entropy).




Figure 7.8 Illustration showing an increase in disorder.

Standard Entropy Change(ΔS°):

It is possible to calculate the actual entropy of a substance at any temperature above 0 K. The absolute entropy of a substance at 298 K and one bar pressure is called the standard entropy S° . The third law of thermodynamics states, according to Nernst, that the absolute entropy of elements is zero only at 0 K in a perfect crystal, and standard entropies of all substances at any temperature above 0 K always have positive values. Once we know the entropies of different substances, we can calculate the standard entropy change (ΔS_r°) for chemical reactions.


$$\Delta S_r^\circ = \sum S_{\text{products}}^\circ - \sum S_{\text{reactants}}^\circ \quad \text{----- (7.30)}$$

Standard Entropy of Formation:

Standard entropy of formation is defined as “the entropy of formation of 1 mole of a compound from the elements under standard conditions”. It is denoted as ΔS_f° . We can calculate the value of entropy of a given compound from the values of S° of elements.

Problem: 7.6

Calculate the standard entropy change for the following reaction (ΔS_r°), given the standard entropies of $\text{CO}_2(\text{g})$, $\text{C}(\text{s})$, $\text{O}_2(\text{g})$ as 213.6, 5.740 and 205 JK^{-1} respectively.



$$\Delta S_r^\circ = \sum S_{\text{products}}^\circ - \sum S_{\text{reactants}}^\circ$$

$$\Delta S_r^\circ = \{S_{\text{CO}_2}^\circ\} - \{S_{\text{C}}^\circ + S_{\text{O}_2}^\circ\}$$

$$\Delta S_r^\circ = 213.6 - [5.74 + 205]$$

$$\Delta S_r^\circ = 213.6 - [210.74]$$

$$\Delta S_r^\circ = 2.86 \text{ JK}^{-1}$$

Evaluate Yourself- 6



Urea on hydrolysis produces ammonia and carbon dioxide. The standard entropies of urea, H_2O , CO_2 , NH_3 are 173.8, 70, 213.5 and 192.5 $\text{J mole}^{-1}\text{K}^{-1}$ respectively. Calculate the entropy change for this reaction.

Entropy change accompanying change of phase

When there is a change of state from

solid to liquid (melting), liquid to vapour (evaporation) or solid to vapour (sublimation) there is a change in entropy. This change may be carried out at constant temperature reversibly as two phases are in equilibrium during the change.

$$\Delta S = \frac{q_{\text{rev}}}{T} = \frac{\Delta H_{\text{rev}}}{T} \quad \text{----- (7.31)}$$

Entropy of fusion:

The heat absorbed, when one mole of a solid melts at its melting point reversibly, is called molar heat of fusion. The entropy change for this process is given by

$$\Delta S_{\text{rev}} = \frac{dq_{\text{rev}}}{T}$$

$$\Delta S_{\text{fusion}} = \frac{\Delta H_{\text{fusion}}}{T_f} \quad \text{----- (7.32)}$$

where ΔH_{fusion} is Molar heat of fusion. T_f is the melting point.

Entropy of vapourisation:

The heat absorbed, when one mole of liquid is boiled at its boiling point reversibly, is called molar heat of vapourisation. The entropy change is given by

$$\Delta S_v = \frac{\Delta H_v}{T_b} \quad \text{----- (7.33)}$$

where ΔH_v is Molar heat of vaporisation. T_b is the boiling point.

Entropy of transition:

The heat change, when one mole of a solid changes reversibly from one allotropic form to another at its transition temperature is called enthalpy of transition. The entropy change is given

$$\Delta S_t = \frac{\Delta H_t}{T_t} \quad \text{----- (7.34)}$$

where ΔH_t is the molar heat of transition, T_t is the transition temperature.


Problem: 7.7

Calculate the entropy change during the melting of one mole of ice into water at 0°C and 1 atm pressure. Enthalpy of fusion of ice is 6008 J mol^{-1}

Given:

$$\Delta H_{\text{fusion}} = 6008 \text{ J mol}^{-1}$$

$$T_f = 0^\circ \text{C} = 273 \text{ K}$$



$$\Delta S_{\text{fusion}} = \frac{\Delta H_{\text{fusion}}}{T_f}$$

$$\Delta S_{\text{fusion}} = \frac{6008}{273}$$

$$\Delta S_{\text{fusion}} = 22.007 \text{ J K}^{-1} \text{ mole}^{-1}$$

Evaluate Yourself - 7

Calculate the entropy change when 1 mole of ethanol is evaporated at 351 K . The molar heat of vaporisation of ethanol is $39.84 \text{ kJ mol}^{-1}$

7.11 Gibbs free energy (G)

One of the important applications of the second law of thermodynamics is to predict the spontaneity of a reaction under a specific set of conditions. A reaction that occurs under the given set of conditions without any external driving force is called a spontaneous reaction. Otherwise, it is said to be non-spontaneous. In our day today life, we observe many spontaneous physical and chemical processes, which includes the following examples.

1. A waterfall runs downhill, but never uphill, spontaneously.
2. A lump of sugar dissolves spontaneously in a cup of coffee, but never reappears in its original form spontaneously.
3. Heat flows from hotter object to a colder one, but never flows from colder to hotter object spontaneously.
4. The expansion of a gas into an evacuated bulb is a spontaneous process, the reverse process that is gathering of all molecules into one bulb is not spontaneous.






Figure 7.9 Spontaneous process illustration

These examples show that the processes that occur spontaneously in one direction, cannot take place in opposite direction spontaneously.

Similarly a large number of exothermic reactions are spontaneous. An example is combustion of methane.




Another example is acid-base neutralization reaction:



However, some endothermic processes are also spontaneous. For

example ammonium nitrate dissolves in water spontaneously though this dissolution is endothermic.



From the above examples we can come to the conclusion that exothermicity favors the spontaneity but does not guarantee it. We cannot decide whether or not a chemical reaction will occur spontaneously solely on the basis of energy changes in the system. We know from second law of thermodynamics that in a spontaneous process, the entropy increases. But not all the processes which are accompanied by an increase in entropy are spontaneous. In order to predict the spontaneity of a reaction, we need some other thermodynamic function.

The second law of thermodynamics introduces another thermodynamic function called Gibbs free energy which finds useful in predicting the spontaneity of a reaction. The Gibbs free energy (G) was developed in the 1870's by Josiah Willard Gibbs. He originally termed this energy as the "available energy" to do work in a system. This quantity is the energy associated with a chemical reaction that can be used to do work.

Gibbs free energy is defined as below

$$G = H - TS \text{ ----- (7.35)}$$

Gibbs free energy (G) is an extensive property and it is a single valued state function.



Let us consider a system which undergoes a change of state from state (1) to state (2) at constant temperature.

$$G_2 - G_1 = (H_2 - H_1) - T(S_2 - S_1)$$

$$\Delta G = \Delta H - T \Delta S \text{-----(7.36)}$$

Now let us consider how ΔG is related to reaction spontaneity.

We know that

$$\Delta S_{\text{total}} = \Delta S_{\text{sys}} + \Delta S_{\text{surr}}$$

For a reversible process (equilibrium), the change in entropy of universe is zero. $\Delta S_{\text{total}} = 0$ [$\because \Delta S_{\text{sys}} = -\Delta S_{\text{surr}}$]

Similarly, for an equilibrium process $\Delta G = 0$

For spontaneous process, $\Delta S_{\text{total}} > 0$

$$\Delta S_{\text{sys}} + \Delta S_{\text{surr}} > 0$$

$$\Delta S_{\text{sys}} + \frac{dq_{\text{surr}}}{T} > 0$$

$$\Delta S_{\text{sys}} - \frac{\Delta H_{\text{sys}}}{T} > 0$$

$$T\Delta S_{\text{sys}} - \Delta H_{\text{sys}} > 0$$

$$-(\Delta H_{\text{sys}} - T\Delta S_{\text{sys}}) > 0$$

$$-\Delta G > 0$$

hence for a spontaneous process,

$$\Delta G < 0$$

$$\text{i.e. } \Delta H - T \Delta S < 0 \text{-----(7.37)}$$

ΔH_{sys} is the enthalpy change of a reaction, $T\Delta S_{\text{sys}}$ is the energy which is not available to do useful work. So ΔG is the net energy available to do useful work and is thus a measure of the 'free energy'. For this reason, it is also known as the free energy of the reaction. For non spontaneous process, $\Delta G > 0$

Gibbs free energy and the net work done by the system:

For any system at constant pressure and temperature

$$\Delta G = \Delta H - T \Delta S \text{----- (7.36)}$$

We know that,

$$\Delta H = \Delta U + P\Delta V$$

$$\therefore \Delta G = \Delta U + P\Delta V - T\Delta S$$

from first law of thermodynamics

if work is done by the system

$$\Delta U = q - w$$

from second law of thermodynamics

$$\Delta S = \frac{q}{T}$$

$$\Delta G = q - w + P\Delta V - T\left(\frac{q}{T}\right)$$

$$\Delta G = -w + P\Delta V$$

$$-\Delta G = w - P\Delta V \text{.....(7.38)}$$

Here, $-P\Delta V$ represents the work done due to expansion against a constant external pressure. Therefore, it is clear that the decrease in free energy ($-\Delta G$) accompanying a process taking place at constant temperature and pressure is equal to the maximum work obtainable from the system other than the work of expansion.

7.11.1 Criteria for spontaneity of a process

The spontaneity of any process depends on three different factors.

- If the enthalpy change of a process is negative, then the process is exothermic and may be spontaneous. (ΔH is negative)
- If the entropy change of a process is positive, then the process may occur spontaneously. (ΔS is positive)

- The Gibbs free energy which is the combination of the above two ($\Delta H - T\Delta S$) should be negative for a reaction to occur spontaneously, i.e. the necessary condition for a reaction to be spontaneous is $\Delta H - T\Delta S < 0$

Table 7.5 Effect of Temperature on Spontaneity of Reactions:


ΔH_r	ΔS_r	$\Delta G_r = \Delta H_r - T\Delta S_r$	Description	Example
–	+	– (at all T)	Spontaneous at all temperature	$2O_3(g) \rightarrow 3O_2(g)$
–	–	– (at low T)	spontaneous at low temperature	Adsorption of gases
		+ (at high T)	non-spontaneous at high temperature	
+	+	+ (at low T)	non-spontaneous at low temperature	Melting of a solid
		– (at high T)	spontaneous at high temperature	
+	–	+ (at all T)	non spontaneous at all temperatures	$2H_2O(g) + O_2(g) \rightarrow 2H_2O_2(l)$

The Table assumes ΔH and ΔS will remain the way indicated for all temperatures. It may not be necessary that way. The Spontaneity of a chemical reaction is only the potential for the reaction to proceed as written. The rate of such processes is determined by kinetic factors, outside of thermodynamical prediction.

Problem: 7.8

Show that the reaction $CO + \left(\frac{1}{2}\right)O_2 \rightarrow CO_2$

at 300K is spontaneous. The standard Gibbs free energies of formation of CO_2 and CO are -394.4 and $-137.2 \text{ kJ mole}^{-1}$ respectively.



$$\Delta G_{(\text{reaction})}^0 = \sum G_{f(\text{products})}^0 - \sum G_{f(\text{reactants})}^0$$

$$\Delta G_{(\text{reaction})}^0 = [G_{CO_2}^0] - \left[G_{CO}^0 + \frac{1}{2} G_{O_2}^0 \right]$$

$$\Delta G_{(\text{reaction})}^0 = -394.4 + [137.2 + 0]$$

$$\Delta G_{(\text{reaction})}^0 = -257.2 \text{ kJ mol}^{-1}$$

$\Delta G_{(\text{reaction})}$ of a reaction at a given temperature is negative hence the reaction is spontaneous.

Evaluate Yourself - 8




For a chemical reaction the values of ΔH and ΔS at 300K are -10 kJ mole^{-1} and $-20 \text{ J deg}^{-1} \text{ mole}^{-1}$ respectively. What is the value of ΔG of the reaction? Calculate the ΔG of a reaction at 600 K assuming ΔH and ΔS values are constant. Predict the nature of the reaction.

7.11.2 Relationship between standard free energy change (ΔG^0) and equilibrium constant (K_{eq}):

In a reversible process, the system is in perfect equilibrium with its surroundings at all times. A reversible chemical reaction can proceed in either direction simultaneously, so that a dynamic equilibrium is set up. This means that the reactions in both the directions should proceed with decrease in free energy, which is impossible. It is possible only if at equilibrium, the free energy of a system is minimum. Let's consider a general

equilibrium reaction



The free energy change of the above reaction in any state (ΔG) is related to the standard free energy change of the reaction (ΔG^0) according to the following equation.

$$\Delta G = \Delta G^0 + RT \ln Q \text{ ----- (7.39)}$$

where Q is reaction quotient and is defined as the ratio of concentration of the products to the concentrations of the reactants under non equilibrium condition.

When equilibrium is attained, there is no further free energy change i.e. $\Delta G = 0$ and Q becomes equal to equilibrium constant. Hence the above equation becomes.

$$\Delta G^0 = -RT \ln K_{eq}$$

This equation is known as Van't Hoff equation.

$$\Delta G^0 = -2.303 RT \log K_{eq} \text{ ----- (7.40)}$$

We also know that

$$\Delta G^0 = \Delta H^0 - T \Delta S^0 = -RT \ln K_{eq}$$

Problem: 7.9

Calculate ΔG^0 for conversion of oxygen to ozone $3/2 O_2 \rightleftharpoons O_{3(g)}$ at 298 K, if K_p for this conversion is 2.47×10^{-29} in standard pressure units.

Solution:

$$\Delta G^0 = -2.303 RT \log K_p$$

Where

$$R = 8.314 \text{ JK}^{-1}\text{mol}^{-1}$$

$$K_p = 2.47 \times 10^{-29}$$

$$T = 298\text{K}$$

$$\Delta G^0 = -2.303(8.314)(298)\log(2.47 \times 10^{-29})$$

$$\Delta G^0 = 163229 \text{ Jmol}^{-1}$$

$$\Delta G^0 = 163.229 \text{ KJ mol}^{-1}$$

7.12 Third law of Thermodynamics

The entropy of a substance varies directly with temperature. Lower the temperature, lower is the entropy. For example, water above 100°C at one atmosphere exists as a gas and has higher entropy (higher disorder). The water molecules are free to roam about in the entire container. When the system is cooled, the water vapour condenses to form a liquid. The water molecules in liquid phase still can move about somewhat freely. Thus the entropy of the system has decreased. On further cooling, water freezes to form ice crystal. The water molecules in the ice crystal are highly ordered and entropy of the system is very low.

If we cool the solid crystal still further, the vibration of molecules held in the crystal lattice gets slower and they have very little freedom of movement (very little disorder) and hence very small entropy.

At absolute zero $\{0 \text{ K (or)} -273^\circ \text{C}\}$, theoretically all modes of motion stop.

Absolute zero is a temperature that an object can get arbitrarily close to but absolute zero will remain unattainable.

Thus the third law of thermodynamics states that the entropy of pure crystalline substance at absolute zero is zero. Otherwise it can be stated as it is impossible to lower the temperature of an object to absolute zero in a finite number of steps. Mathematically,

$$\lim_{T \rightarrow 0} S = 0 \text{ for a perfectly ordered crystalline state.}$$




Figure 7.10 Third law of Thermodynamics

Crystals with defects (imperfection) at absolute zero, have entropy greater than zero. Absolute entropy of a substance can never be negative.

SUMMARY

The branch of science which deals the relation between energy, heat and work is called Thermodynamics. The main aim of the study of chemical thermodynamics is to learn (i) transformation of energy from one form into another form (ii) Utilization of various forms of energies.

System: A system is defined as any part of universe under consideration. There are three types of thermodynamic systems. They are (i) isolated system (ii) closed system and (iii) open system.

Surrounding: Everything in the universe that is not the part of the system is called surrounding.

Boundary: Anything which separates the system from its surrounding is called boundary.

Thermodynamic Properties: Any quantity that depends only on the state of system is called thermodynamic property of a system. There are two kinds of thermodynamic properties called (1) intensive - independent of the quantity of material and (2) extensive - directly proportional to the quantity of material. There are five basic thermodynamic properties. (U, H, S and G)

Thermodynamic Processes:

Adiabatic process in which no heat transfer takes place ($q = 0$)

Isothermal process in which

temperature remains constant ($dT = 0$).

Isobaric process in which pressure remains constant ($dP = 0$).

Isochoric process in which volume remains constant ($dV = 0$).

Cyclic process in which the system returns to its original state after completing a series of changes.

Internal energy (U): Internal energy of a system is equal to the energy possessed by all its constituents namely atoms, ions and molecules. The energy of a system of molecules is equal to the sum of its translational energy, vibrational energy, rotational energy, bond energy, electronic energy and energy due to molecular interactions.

Heat: Heat is regarded as the energy in transit across the boundary separating a system from its surrounding. Heat is a path function. The SI unit of heat is joule (J)

Work : Work is defined as the force (F) multiplied by the displacement $-w = F \cdot x$, work is measured in Joules, i.e the SI unit of work is Joule. During expansion or compression of a gas the work done is calculated by the relation $w = -P\Delta V$.

The sign conventions for heat and work are as follows:

If heat is absorbed by the system : $+q$

If heat is evolved by the system : -q

If work is done by the system : -w

If work is done on the system : +w

Laws of Thermodynamics:

Zeroth law : When two systems are separately in equilibrium with a third system, they are in equilibrium with each other.

First law: "Energy can neither be created nor destroyed, but may be converted from one form to another".

$$\Delta U = q + w.$$

Enthalpy is a thermodynamic property of a system. Enthalpy H is defined as the sum of the internal energy and pressure volume work.

$$H = U + PV. \text{ Enthalpy change, } \Delta H = \Delta U + \Delta n_g RT.$$

Hess's law: It states that "the enthalpy change of a reaction either at constant volume or constant pressure is the same whether it takes place in a single or multiple steps". Hess's law can be applied to calculate enthalpies of reactions that are difficult to measure.

Heat capacities: [C_p and C_v]

Heat capacity is defined as the amount of energy required to increase the temperature of one unit quantity of material by one degree, under specific conditions. It can be measured under two different conditions, namely,

(a) constant pressure $C_p = (dH/dT)_p$

(b) constant volume $C_v = (dU/dT)_v$

Second law of thermodynamics:

The second law of thermodynamics helps us to predict whether the reaction is feasible or not and also tells the direction of the flow of heat.

To predict spontaneity of a process, a new thermodynamic quantity namely entropy (S) was introduced. Entropy is a measure of the randomness or disorderliness of the system.

Entropy statement: "whenever a spontaneous process takes place, it is accompanied by an increase in the total entropy of the universe".

Kelvin-Planck statement: It is impossible to take heat from a hotter reservoir and convert it completely into work by a cyclic process without transferring a part of that heat to a colder reservoir.

Clausius statement: This statement recognises that heat flows spontaneously from hot objects to cold objects and to get it flow in the opposite direction, we have to spend some work.

Gibbs Free Energy (G):

G is expressed as $G = H - TS$, free energy change of a process is given by the relation $\Delta G = \Delta H - T\Delta S$.

Standard free energy change and equilibrium constants are related by the equation $\Delta G^0 = -RT \ln K_{eq}$

Third law:

The entropy of a perfectly crystalline material at absolute zero is zero.



EVALUATION



I. Choose the best answer

- The amount of heat exchanged with the surrounding at constant pressure is given by the quantity
 - ΔE
 - ΔH
 - ΔS
 - ΔG
- All the naturally occurring processes proceed spontaneously in a direction which leads to
 - decrease in entropy
 - increase in enthalpy
 - increase in free energy
 - decrease in free energy
- In an adiabatic process, which of the following is true ?
 - $q = w$
 - $q = 0$
 - $\Delta E = q$
 - $P \Delta V = 0$
- In a reversible process, the change in entropy of the universe is
 - > 0
 - ≥ 0
 - < 0
 - $= 0$
- In an adiabatic expansion of an ideal gas
 - $w = -\Delta u$
 - $w = \Delta u + \Delta H$
 - $\Delta u = 0$
 - $w = 0$
- The intensive property among the quantities below is
 - mass
 - volume
 - enthalpy
 - $\frac{\text{mass}}{\text{volume}}$
- An ideal gas expands from the volume of $1 \times 10^{-3} \text{ m}^3$ to $1 \times 10^{-2} \text{ m}^3$ at 300 K against a constant pressure at $1 \times 10^5 \text{ Nm}^{-2}$. The work done is
 - 900 J
 - 900 kJ
 - 270 kJ
 - 900 kJ
- Heat of combustion is always
 - positive
 - negative
 - zero
 - either positive or negative
- The heat of formation of CO and CO_2 are - 26.4 kCal and - 94 kCal, respectively. Heat of combustion of carbon monoxide will be
 - + 26.4 kcal
 - 67.6 kcal
 - 120.6 kcal
 - + 52.8 kcal



10. $C(\text{diamond}) \rightarrow C(\text{graphite})$, $\Delta H = -ve$, this indicates that
- graphite is more stable than diamond
 - graphite has more energy than diamond
 - both are equally stable
 - stability cannot be predicted
11. The enthalpies of formation of Al_2O_3 and Cr_2O_3 are -1596 kJ and -1134 kJ , respectively.
- ΔH for the reaction $2Al + Cr_2O_3 \rightarrow 2Cr + Al_2O_3$ is
- -1365 kJ
 - 2730 kJ
 - -2730 kJ
 - -462 kJ
12. Which of the following is not a thermodynamic function ?
- internal energy
 - enthalpy
 - entropy
 - frictional energy
13. If one mole of ammonia and one mole of hydrogen chloride are mixed in a closed container to form ammonium chloride gas, then
- $\Delta H > \Delta U$
 - $\Delta H - \Delta U = 0$
 - $\Delta H + \Delta U = 0$
 - $\Delta H < \Delta U$
14. Change in internal energy, when 4 kJ of work is done on the system and 1 kJ of heat is given out by the system is
- $+1 \text{ kJ}$
 - -5 kJ
 - $+3 \text{ kJ}$
 - -3 kJ
15. The work done by the liberated gas when 55.85 g of iron (molar mass 55.85 g mol^{-1}) reacts with hydrochloric acid in an open beaker at 25°C
- -2.48 kJ
 - -2.22 kJ
 - $+2.22 \text{ kJ}$
 - $+2.48 \text{ kJ}$
16. The value of ΔH for cooling 2 moles of an ideal monatomic gas from 125°C to 25°C at constant pressure will be $\left[\text{given } C_p = \frac{5}{2} R \right]$
- $-250 R$
 - $-500 R$
 - $500 R$
 - $+250 R$
17. Given that $C(g) + O_2(g) \rightarrow CO_2(g)$ $\Delta H^0 = -a \text{ kJ}$; $2CO(g) + O_2(g) \rightarrow 2CO_2(g)$ $\Delta H^0 = -b \text{ kJ}$; Calculate the ΔH^0 for the reaction $C(g) + \frac{1}{2}O_2(g) \rightarrow CO(g)$
- $\frac{b+2a}{2}$
 - $2a-b$
 - $\frac{2a-b}{2}$
 - $\frac{b-2a}{2}$



18. When 15.68 litres of a gas mixture of methane and propane are fully combusted at 0°C and 1 atmosphere, 32 litres of oxygen at the same temperature and pressure are consumed. The amount of heat released from this combustion in kJ is ($\Delta H_{\text{C}}(\text{CH}_4) = -890 \text{ kJ mol}^{-1}$ and $\Delta H_{\text{C}}(\text{C}_3\text{H}_8) = -2220 \text{ kJ mol}^{-1}$)
- a) -889 kJ b) -1390 kJ c) -3180 kJ d) -632.68 kJ
19. The bond dissociation energy of methane and ethane are 360 kJ mol^{-1} and 620 kJ mol^{-1} respectively. Then, the bond dissociation energy of C-C bond is
- a) 170 kJ mol^{-1} b) 50 kJ mol^{-1} c) 80 kJ mol^{-1} d) 220 kJ mol^{-1}
20. The correct thermodynamic conditions for the spontaneous reaction at all temperature is (NEET Phase - I)
- a) $\Delta H < 0$ and $\Delta S > 0$
b) $\Delta H < 0$ and $\Delta S < 0$
c) $\Delta H > 0$ and $\Delta S = 0$
d) $\Delta H > 0$ and $\Delta S > 0$
21. The temperature of the system, decreases in an _____
- a) Isothermal expansion b) Isothermal Compression c) adiabatic expansion
d) adiabatic compression
22. In an isothermal reversible compression of an ideal gas the sign of q , ΔS and w are respectively
- a) $+$, $-$, $-$ b) $-$, $+$, $-$ c) $+$, $-$, $+$ d) $-$, $-$, $+$
23. Molar heat of vapourisation of a liquid is 4.8 kJ mol^{-1} . If the entropy change is $16 \text{ J mol}^{-1} \text{ K}^{-1}$, the boiling point of the liquid is
- a) 323 K b) 27°C c) 164 K d) 0.3 K
24. ΔS is expected to be maximum for the reaction
- a) $\text{Ca}(\text{S}) + \frac{1}{2} \text{O}_2(\text{g}) \rightarrow \text{CaO}(\text{S})$
b) $\text{C}(\text{S}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g})$
c) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}(\text{g})$
d) $\text{CaCO}_3(\text{S}) \rightarrow \text{CaO}(\text{S}) + \text{CO}_2(\text{g})$



25. The values of ΔH and ΔS for a reaction are respectively 30 kJ mol^{-1} and $100 \text{ JK}^{-1} \text{ mol}^{-1}$. Then the temperature above which the reaction will become spontaneous is
- a) 300 K b) 30 K c) 100 K d) 20^0 C

II. Write brief answer to the following questions:

26. State the first law of thermodynamics.
27. Define Hess's law of constant heat summation.
28. Explain intensive properties with two examples
29. Define the following terms:
- a. isothermal process b. adiabatic process
- c. isobaric process d. isochoric process
30. What is the usual definition of entropy? What is the unit of entropy?
31. Predict the feasibility of a reaction when
- i) both ΔH and ΔS positive ii) both ΔH and ΔS negative
- iii) ΔH decreases but ΔS increases
32. Define is Gibbs's free energy.
33. Define enthalpy of combustion.
34. Define molar heat capacity. Give its unit.
35. Define the calorific value of food. What is the unit of calorific value?
36. Define enthalpy of neutralization.
37. What is lattice energy?
38. What are state and path functions? Give two examples.
39. Give Kelvin statement of second law of thermodynamics.
40. The equilibrium constant of a reaction is 10, what will be the sign of ΔG ? Will this reaction be spontaneous?
41. Enthalpy of neutralization is always a constant when a strong acid is neutralized by a strong base: account for the statement.



42. State the third law of thermodynamics.
43. Write down the Born-Haber cycle for the formation of CaCl_2
44. Identify the state and path functions out of the following: a) Enthalpy b) Entropy c) Heat d) Temperature e) Work f) Free energy.
45. State the various statements of second law of thermodynamics.
46. What are spontaneous reactions? What are the conditions for the spontaneity of a process?
47. List the characteristics of internal energy.
48. Explain how heat absorbed at constant volume is measured using bomb calorimeter with a neat diagram.
49. Calculate the work involved in expansion and compression process.
50. Derive the relation between ΔH and ΔU for an ideal gas. Explain each term involved in the equation.
51. Suggest and explain an indirect method to calculate lattice enthalpy of sodium chloride crystal.
52. List the characteristics of Gibbs free energy.
53. Calculate the work done when 2 moles of an ideal gas expands reversibly and isothermally from a volume of 500 ml to a volume of 2 L at 25°C and normal pressure.
54. In a constant volume calorimeter, 3.5 g of a gas with molecular weight 28 was burnt in excess oxygen at 298 K. The temperature of the calorimeter was found to increase from 298 K to 298.45 K due to the combustion process. Given that the calorimeter constant is 2.5 kJ K^{-1} . Calculate the enthalpy of combustion of the gas in kJ mol^{-1} .
55. Calculate the entropy change in the system, and surroundings, and the total entropy change in the universe during a process in which 245 J of heat flow out of the system at 77°C to the surrounding at 33°C .
56. 1 mole of an ideal gas, maintained at 4.1 atm and at a certain temperature, absorbs heat 3710 J and expands to 2 litres. Calculate the entropy change in expansion process.
57. 30.4 kJ is required to melt one mole of sodium chloride. The entropy change during melting is $28.4 \text{ JK}^{-1} \text{ mol}^{-1}$. Calculate the melting point of sodium chloride.
58. Calculate the standard heat of formation of propane, if its heat of combustion is $-2220.2 \text{ kJ mol}^{-1}$. the heats of formation of $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\text{l})$ are -393.5 and $-285.8 \text{ kJ mol}^{-1}$ respectively.

59. You are given normal boiling points and standard enthalpies of vapourisation. Calculate the entropy of vapourisation of liquids listed below.

S. No	Liquid	Boiling points ($^{\circ}\text{C}$)	ΔH (kJ mol $^{-1}$)
1.	Ethanol	78.4	+ 42.4
2.	Toluene	110.6	+ 35.2

60. For the reaction $\text{Ag}_2\text{O}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \frac{1}{2} \text{O}_2(\text{g})$: $\Delta H = 30.56 \text{ kJ mol}^{-1}$ and $\Delta S = 6.66 \text{ J K}^{-1} \text{ mol}^{-1}$ (at 1 atm). Calculate the temperature at which ΔG is equal to zero. Also predict the direction of the reaction (i) at this temperature and (ii) below this temperature.

61. What is the equilibrium constant K_{eq} for the following reaction at 400K.






given that $\Delta H^{\circ} = 77.2 \text{ kJ mol}^{-1}$;

and $\Delta S^{\circ} = 122 \text{ J K}^{-1} \text{ mol}^{-1}$.



62. Cyanamide (NH_2CN) is completely burnt in excess oxygen in a bomb calorimeter, ΔU was found to be $-742.4 \text{ kJ mol}^{-1}$, calculate the enthalpy change of the reaction at 298K. $\text{NH}_2\text{CN}(\text{s}) + \frac{3}{2} \text{O}_2(\text{g}) \rightarrow \text{N}_2(\text{g}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$ $\Delta H = ?$

63. Calculate the enthalpy of hydrogenation of ethylene from the following data. Bond energies of C – H, C – C, C = C and H – H are 414, 347, 618 and 435 kJ mol $^{-1}$.

64. Calculate the lattice energy of CaCl_2 from the given data



65. Calculate the enthalpy change for the reaction $\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2$ from the following data.






66. When 1-pentyne (A) is treated with 4N alcoholic KOH at 175°C, it is converted slowly into an equilibrium mixture of 1.3% 1-pentyne(A) , 95.2% 2-pentyne(B) and 3.5% of 1,2 pentadiene (C) the equilibrium was maintained at 175°C, calculate ΔG^0 for the following equilibria.



67. At 33K, N_2O_4 is fifty percent dissociated. Calculate the standard free energy change at this temperature and at one atmosphere.
68. The standard enthalpies of formation of SO_2 and SO_3 are -297 kJ mol^{-1} and -396 kJ mol^{-1} respectively. Calculate the standard enthalpy of reaction for the reaction: $SO_2 + \frac{1}{2} O_2 \rightarrow SO_3$
69. For the reaction at 298 K : $2A + B \rightarrow C$
- $\Delta H = 400 \text{ KJ mol}^{-1}$; $\Delta S = 0.2 \text{ KJK}^{-1} \text{ mol}^{-1}$ Determine the temperature at which the reaction would be spontaneous.
70. Find out the value of equilibrium constant for the following reaction at 298K, $2 NH_3(g) + CO_2 (g) \rightleftharpoons NH_2CONH_2 (aq) + H_2O (l)$ Standard Gibbs energy change, ΔG_r^0 at the given temperature is $-13.6 \text{ kJ mol}^{-1}$.
71. A gas mixture of 3.67 lit of ethylene and methane on complete combustion at 25°C and at 1 atm pressure produce 6.11 lit of carbondioxide. Find out the amount of heat evolved in kJ, during this combustion. ($\Delta H_C(CH_4) = -890 \text{ kJ mol}^{-1}$ and ($\Delta H_C(C_2H_4) = -1423 \text{ kJ mol}^{-1}$

CONCEPT MAP



Appendix

S.No	Content	Page No.
1.	Fundamental constants	230
2.	Commonly used SI prefixes	230
3.	Inter-conversion of units	231
4.	Critical constants of selected gases	231
5.	Van der waals constants for selected gases	232
6.	Molar heat capacities of selected compounds	233
7.	Standard heat of combustion	233
8.	Thermodynamic data for selected compounds	234
9.	Lattice Enthalpies of alkali metal halides	235
10.	Cp and Cv values for mono, di and triatomic gases	235
	KEY ANSWERS	
11.	Unit - 1 Basic Concepts of Chemistry and Chemical Calculations	236
12.	Unit - 2 Quantum Mechanical Model of Atom	252
13.	Unit -3 Periodic classification of Elements	260
14.	Unit - 4 Hydrogen	263
15.	Unit -5 Alkali and Alkaline Earth Metals	265
16.	Unit - 6 Gaseous State	268
17.	Unit - 7 Thermodynamics	275
18.	List Of Reference Book	289
19.	Glossary	290

Fundamental constants

Avogadro number	N	6.022×10^{23}
Atomic mass unit (or) Unified atomic mass	amu (or) u	1.660×10^{-27} kg
Speed of light	c	2.997×10^8 m s ⁻¹
Elementary charge (or) Charge of an electron	e	1.602×10^{-19} C
Planck's constant	h	6.626×10^{-34} Js
Mass of electron	m _e	9.109×10^{-31} kg
Mass of proton	m _p	1.672×10^{-27} kg
Mass of neutron	m _n	1.674×10^{-27} kg
Bohr's radius	a ₀	0.529 Å
Rydberg constant	R _∞	1.097×10^5 cm ⁻¹
Universal gas constant	R	8.314 J K ⁻¹ mol ⁻¹ 0.082 L atm K ⁻¹ mol ⁻¹ 1.987 cal K ⁻¹ mol ⁻¹ 62.36 L mmHg K ⁻¹ mol ⁻¹ 83.14 L mbar K ⁻¹ mol ⁻¹

Commonly used SI Prefixes

femto	pico	nano	micro	milli	centi	deci	kilo	mega	giga
f	p	n	μ	m	c	d	k	M	G
10 ⁻¹⁵	10 ⁻¹²	10 ⁻⁹	10 ⁻⁶	10 ⁻³	10 ⁻²	10 ⁻¹	10 ³	10 ⁶	10 ⁹

Inter - Conversion of Units

1 Å	=	10^{-10} m	
1 atm	=	101.325 kPa	= $1.01325 \times 10^5 \text{ N m}^{-2}$
1 atm	=	$1.01325 \times 10^5 \text{ J m}^{-3}$	= 760 Torr (mmHg)
1 eV	=	1.60219×10^{-19} J	
1 N	=	1 J m^{-1}	= 10^5 dyne
1 J	=	10^7 erg	= $1 \text{ Kg m}^2 \text{ s}^{-2}$
1 cal	=	4.184 J	

Critical constants of selected gases

Gas	Critical Pressure, P_c (atm)	Critical Volume, V_c ($\text{cm}^3 \text{ mol}^{-1}$)	Critical Temperature, T_c (K)
He	2.26	57.76	5.21
Ne	26.86	41.74	44.44
Ar	48.00	75.25	150.72
Kr	54.27	92.24	209.39
Xe	58.0	118.8	289.75
H ₂	12.8	64.99	33.23
N ₂	33.54	90.10	126.3
O ₂	50.14	78.0	154.8
Cl ₂	76.1	124	417.2
Br ₂	102	135	584
HCl	81.5	81.0	324.7
CO ₂	72.85	94.0	304.2
H ₂ O	218.3	55.3	647.4
NH ₃	111.3	72.5	405.5
CH ₄	45.6	98.7	190.6
C ₂ H ₄	50.50	124	283.1
C ₂ H ₆	48.20	148	305.4
C ₆ H ₆	48.6	260	562.7

Van der waals constants for select gases

Name	a (L ² bar mol ⁻²)	b (L mol ⁻¹)
Ammonia	4.225	0.0371
Benzene	18.24	0.1154
Carbon dioxide	3.64	0.04267
Carbon monoxide	1.505	0.03985
Chlorine	6.579	0.05622
Helium	0.0346	0.0238
Hexane	24.71	0.1735
Hydrogen	0.2476	0.02661
Hydrogen bromide	4.51	0.04431
Hydrogen chloride	3.716	0.04081
Hydrogen sulfide	4.49	0.04287
Mercury	8.2	0.01696
Methane	2.283	0.04278
Nitric oxide	1.358	0.02789
Nitrogen	1.37	0.0387
Nitrogen dioxide	5.354	0.04424
Nitrous oxide	3.832	0.04415
Oxygen	1.382	0.03186
Sulfur dioxide	6.803	0.05636
Water	5.536	0.03049

Molar heat capacities of select compounds:

	$C_{v,m}$ (J K ⁻¹ mol ⁻¹)	$C_{p,m}$ (J K ⁻¹ mol ⁻¹)
Gases		
Inert gases	12.48	20.79
H ₂	20.44	28.82
N ₂	20.74	29.12
O ₂	20.95	29.36
CO ₂	28.46	37.11
Liquids		
H ₂ O	-	75.29
CH ₃ OH	-	81.6
C ₂ H ₅ OH	-	111.5
C ₆ H ₆	-	136.1
Solids		
C (diamond)	-	6.11
Cu	-	244.4
Fe	-	25.1
SiO ₂	-	44.4

Standard heat of combustion (ΔH_f^0 , kJ mol⁻¹)

Methane(g)	=	- 890	Ethanol(l)	=	- 1368
Ethane(g)	=	- 1560	Acetic acid (l)	=	- 875
Propane(g)	=	- 2220	Benzoic acid (s)	=	- 3227
Butane(g)	=	- 2878	Glucose (s)	=	- 2808
Benzene(l)	=	- 3268	Sucrose (s)	=	- 5645
Methanol(l)	=	- 726			

Thermodynamic data for select compounds

	ΔH_f^0 (kJ mol ⁻¹)	ΔG_f^0 (kJ mol ⁻¹)	S^0 (J K ⁻¹ mol ⁻¹)
C (s, graphite)	0	0	5.740
H ₂ (g)	0	0	130.684
N ₂ (g)	0	0	191.61
O ₂ (g)	0	0	205.138
Cl ₂ (g)	0	0	223.07
Ag(s)	0	0	42.55
Cu(s)	0	0	33.150
Au(s)	0	0	47.40
Fe(s)	0	0	27.28
Zinc(s)	0	0	41.63
C(s, diamond)	1.895	2.900	2.377
CO(g)	-110.53	-137.17	197.67
CO ₂ (g)	-393.51	-394.36	213.74
HCl(g)	-92.31	-95.30	186.91
H ₂ O(l)	-285.83	-237.13	69.91
H ₂ O(g)	-241.82	-228.57	188.83
NH ₃ (g)	-46.11	-16.45	192.45
Methane(g)	-74.81	-50.72	186.26
Ethane(g)	-84.68	-32.82	229.60
Propane(g)	-103.85	-23.49	269.91
Butane(g)	-126.15	-17.03	310.23
Benzene(l)	49.0	124.3	173.3
Methanol(l)	-238.66	-166.27	126.8
Ethanol(l)	-277.69	-174.78	160.7

Lattice Enthalpies of alkali metal halides ($-\Delta H^0$ Lattice, kJ mol^{-1})

	F	Cl	Br	I
Li	1022	846	800	744
Na	902	771	733	684
K	801	701	670	629
Rb	767	675	647	609
Cs	716	645	619	585

C_p and C_v for Mono, di and tri atomic gases

	C_v Value	C_p Value ($C_p = R + C_v$)
Monatomic gas	$\frac{3}{2}R$	$\frac{5}{2}R$
Diatomic gas	$\frac{5}{2}R$	$\frac{7}{2}R$
Triatomic	$\frac{6}{2}R$	$\frac{8}{2}R$

Unit - 1

Basic Concepts of Chemistry and Chemical Calculations

Evaluate yourself :

1. (i) **Element** - Copper wire, Silver plate
(ii) **Compound** - Sugar, distilled water, carbon dioxide, Table salt, Naphthalene balls
(iii) **Mixture** - Sea water

2. (i) $\text{C}_2\text{H}_5\text{OH}$: $(2 \times 12) + (5 \times 1) + (1 \times 16) + (1 \times 1) = 46 \text{ g}$
(ii) KMnO_4 : $(1 \times 39) + (1 \times 55) + (4 \times 16) = 158 \text{ g}$
(iii) $\text{K}_2\text{Cr}_2\text{O}_7$: $(2 \times 39) + (2 \times 52) + (7 \times 16) = 294 \text{ g}$
(iv) $\text{C}_{12}\text{H}_{22}\text{O}_{11}$: $(12 \times 12) + (22 \times 1) + (11 \times 16) = 342 \text{ g}$

3. (a) Molar mass of ethane, $\text{C}_2\text{H}_6 = (2 \times 12) + (6 \times 1) = 30 \text{ g mol}^{-1}$

$$n = \text{mass} / \text{molar mass} = 9 \text{ g} / 30 \text{ g mol}^{-1} = 0.3 \text{ mole}$$

- (b) At 273 K and 1 atm pressure 1 mole of a gas occupies a volume of 22.4 L

Therefore,

number of moles of oxygen, that occupies a volume of 224 ml at 273 K and 3 atm pressure

$$= \frac{1 \text{ mole}}{273 \text{ K} \times 1 \text{ atm} \times 22.4 \text{ L}} \times 0.224 \text{ L} \times 273 \text{ K} \times 3 \text{ atm}$$

$$= 0.03 \text{ mole}$$

1 mole of oxygen contains 6.022×10^{23} molecules

0.03 mole of oxygen contains $= 6.022 \times 10^{23} \times 0.03$

$$= 1.807 \times 10^{22} \text{ molecules of oxygen}$$

4. (a) Mass of the metal = 0.456 g

Mass of the metal chloride = 0.606 g

0.456 g of the metal combines with 0.15 g of chlorine.

Mass of the metal that combines with 35.5 g of chlorine is $\frac{0.456}{0.15} \times 35.5$

$$= 107.92 \text{ g eq}^{-1}.$$

$$\begin{aligned}
 \text{(b) Equivalent mass of a oxidising agent} &= \frac{\text{molar mass}}{\text{number of moles of electrons gained by one mole of the reducing agent}} \\
 &= \frac{294.2 \text{ g mol}^{-1}}{6 \text{ eq mol}^{-1}} \\
 &= \mathbf{49.0 \text{ g eq}^{-1}}
 \end{aligned}$$

5.

Element	Percentage Composition	Atomic mass	Relative no. of atoms = $\frac{\text{Percentage}}{\text{Atomic mass}}$	Simple ratio
C	54.55 %	12	$54.55/12 = 4.55$	$4.55 / 2.27 = 2$
H	9.09 %	1	$9.09 / 1 = 9.09$	$9.09 / 2.27 = 4$
O	36.36 %	16	$36.36/16 = 2.27$	$2.27/2.27 = 1$
Empirical formula (C₂H₄O)				

6.

Element	Percentage Composition	Relative no. of atoms = $\frac{\text{Percentage}}{\text{Atomic mass}}$	Atomic mass = $\frac{\text{Percentage}}{\text{Relative no. of atoms}}$	Simple ratio
X	32 %	2	16	4
Y	24 %	1	24	2
Z	44 %	0.5	88	1
Empirical formula (X₄Y₂Z)				

$$\begin{aligned}
 \text{Calculated empirical formula mass} &= (16 \times 4) + (24 \times 2) + 88 \\
 &= 64 + 48 + 88 = 200
 \end{aligned}$$

$$\begin{aligned}
 n &= \frac{\text{molar mass}}{\text{calculated empirical formula mass}} \\
 \therefore n &= \frac{400}{200} \\
 &= 2
 \end{aligned}$$

$$\therefore \text{Molecular formula (X}_4\text{Y}_2\text{Z)}_2 = \mathbf{X_8Y_4Z_2}$$

7.


Content	Reactant		Products	
	x	y	l	m
Stoichiometric coefficient	2	3	4	1
No. of moles allowed to react	8	15	–	–
No. of moles of reactant reacted and product formed	8	12	16	4
No. of moles of un-reacted reactants and the product formed	–	3	16	4

Limiting reagent : x


Product formed : 16 moles of l & 4 moles of m

Amount of excess reactant : 3 moles of y


8.



Equate the total no. of electrons in the reactant side by cross multiplying,



Based on reactant side, balance the products




Product side : 36 hydrogen atoms & 88 oxygen atoms

Reactant side : 28 hydrogen atoms & 84 oxygen atoms

Difference is 8 hydrogen atoms & 4 oxygen atoms

∴ Add 4 H₂O molecule on the reactant side.

Balanced equation is,





EVALUATION




I Choose the best answer

1. (a) 40 ml of CO_2 gas
2. (d) 200 u
3. (c) assertion is true but reason is false
4. (b) oxygen
5. (a) 102 g
6. (c) 6.022×10^{20}
7. (c) 16 %
8. (c) 0.075
9. (d) 1 mole of HCl
10. (c) $\text{BaCl}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2 \text{HCl}$
11. (b) $\text{P}_4 + 3 \text{NaOH} + 3 \text{H}_2\text{O} \rightarrow \text{PH}_3 + 3 \text{NaH}_2\text{PO}_2$
12. (b) 52.7
13. (d) 6.022×10^{24} molecules of water
14. (a) NO
15. (a) 6.022×10^{23}
16. (c) $\text{S}_2\text{O}_4^{2-} < \text{SO}_3^{2-} < \text{S}_2\text{O}_6^{2-} < \text{SO}_4^{2-}$
17. (c) molar mass of ferrous oxalate / 3
18. (d) the mass of one mole of carbon
19. (c) the ratio between the number of molecules in A to number of molecules in B is 2:1
20. (a) 3.59 g
21. (b) 44 g mol^{-1}
22. (c) both (a) & (b)
23. (a) propene
24. (a) relative atomic mass is 12 u
25. (a) ${}_6\text{C}^{12}$



Key to the multiple choice questions



Content	CH_4	O_2	CO_2
Stoichiometric coefficient	1	2	1
Volume of reactants allowed to react	40 mL	80 mL	
Volume of reactant reacted and product formed	40 mL	80 mL	40 mL
Volume of gas after cooling to the room temperature	-	-	-

Since the product was cooled to room temperature, water exists mostly as liquid.
Hence, option (a) is correct

2.

$$= \frac{(200 \times 90) + (199 \times 8) + (202 \times 2)}{100} = 199.96$$

200 u

3. Correct reason: Total number of entities present in one mole of any substance is equal to 6.022×10^{23} .

4. Reaction 1 : $2\text{C} + \text{O}_2 \rightarrow 2\text{CO}$
 2×12 g carbon combines with 32 g of oxygen. Hence,

$$\text{Equivalent mass of carbon} = \frac{2 \times 12}{32} \times 8 = 6$$

Reaction 2 : $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$

12 g carbon combines with 32 g of oxygen. Hence,

$$\text{Equivalent mass of carbon} = \frac{12}{32} \times 8 = 3$$

5. Let the trivalent metal be M^{3+}

Equivalent mass = mass of the metal / valance factor

9 g eq^{-1} = mass of the metal / 3 eq

Mass of the metal = 27 g

Oxide formed M_2O_3 ;

Mass of the oxide = $(2 \times 27) + (3 \times 16)$


= **102 g**

6. Weight of the water drop = 0.018 g

$$\begin{aligned}\text{No. of moles of water in the drop} &= \text{Mass of water} / \text{molar mass} \\ &= 0.018 / 18 = 10^{-3} \text{ mole}\end{aligned}$$

$$\text{No of water molecules present in 1 mole of water} = 6.022 \times 10^{23}$$

$$\begin{aligned}\text{No. water molecules in one drop of water (10}^{-3} \text{ mole)} &= 6.022 \times 10^{23} \times 10^{-3} \\ &= \mathbf{6.022 \times 10^{20}}\end{aligned}$$



$$\text{MgCO}_3 : (1 \times 24) + (1 \times 12) + (3 \times 16) = 84 \text{ g}$$

$$\text{CO}_2 : (1 \times 12) + (2 \times 16) = 44 \text{ g}$$


100 % pure 84 g MgCO_3 on heating gives 44 g CO_2

Given that 1 g of MgCO_3 on heating gives 0.44 g CO_2

Therefore, 84 g MgCO_3 sample on heating gives 36.96 g CO_2

$$\begin{aligned}\text{Percentage of purity of the sample} &= \frac{100 \%}{44 \text{ g CO}_2} \times 36.96 \text{ g CO}_2 \\ &= 84 \%\end{aligned}$$

Percentage of impurity = **16 %**





$$6.3 \text{ g} + 30 \text{ g} \rightarrow 33 \text{ g} + x$$

The amount of CO_2 released, $x = 3.3 \text{ g}$

$$\text{No. of moles of CO}_2 \text{ released} = 3.3 / 44 = 0.075 \text{ mol}$$



Content	$\text{H}_2(\text{g})$	$\text{Cl}_2(\text{g})$	$\text{HCl}(\text{g})$
Stoichiometric coefficient	1	1	2
No. of moles of reactants allowed to react at 273 K and 1 atm pressure	22.4 L (1 mol)	11.2 L (0.5 mol)	-
No. of moles of reactant reacted and product formed	0.5	0.5	1
Amount of HCl formed = 1 mol			



12. The reduction reaction of the oxidising agent (MnO_4^-) involves gain of 3 electrons.
Hence the equivalent mass = (Molar mass of KMnO_4)/3 = $158.1/3 = 52.7$

13. No. of moles of water present in 180 g

$$= \text{Mass of water} / \text{Molar mass of water}$$

$$= 180 \text{ g} / 18 \text{ g mol}^{-1} = 10 \text{ moles}$$

One mole of water contains

$$= 6.022 \times 10^{23} \text{ water molecules}$$

$$10 \text{ mole of water contains} = 6.022 \times 10^{23} \times 10 = 6.022 \times 10^{24} \text{ water molecules}$$

14. 7.5 g of gas occupies a volume of 5.6 liters at 273 K and 1 atm pressure Therefore,
the mass of gas that occupies a volume of 22.4 liters

$$= \frac{7.5 \text{ g}}{5.6 \text{ L}} \times 22.4 \text{ L} = 30 \text{ g}$$

Molar mass of NO (14+16) = 30 g

15. No. of electrons present in one ammonia (NH_3) molecule (7 + 3) = 10



$$\begin{aligned} \text{No. of moles of ammonia} &= \frac{\text{Mass}}{\text{Molar mass}} \\ &= \frac{1.7 \text{ g}}{17 \text{ g mol}^{-1}} \\ &= 0.1 \text{ mol} \end{aligned}$$

No. of molecules present in 0.1 mol of ammonia

$$= 0.1 \times 6.022 \times 10^{23} = 6.022 \times 10^{22}$$

No. of electrons present in 0.1 mol of ammonia

$$= 10 \times 6.022 \times 10^{22} = 6.022 \times 10^{23}$$



18. The mass of one mole of carbon

$$\begin{aligned} 19. \quad \text{No. of moles of oxygen} &= 8 \text{ g} / 32 \text{ g} \\ &= 0.25 \text{ moles of oxygen} \\ \text{No. of moles of sulphur dioxide} &= 8 \text{ g} / 64 \text{ g} \\ &= 0.125 \text{ moles of sulphur dioxide} \\ \text{Ratio between the no. of molecules} &= 0.25 : 0.125 \\ &= 2 : 1 \end{aligned}$$

20. $\text{AgNO}_3 + \text{KCl} \rightarrow \text{KNO}_3 + \text{AgCl}$

50 mL of 8.5 % solution contains 4.25 g of AgNO_3

No. of moles of AgNO_3 present in 50 mL of 8.5 % AgNO_3 solution

$$= \text{Mass} / \text{Molar mass}$$

$$= 4.25 / 170$$

$$= 0.025 \text{ moles}$$

Similarly, No of moles of KCl present in 100 mL of 1.865 % KCl solution

$$= 1.865 / 74.5$$

$$= 0.025 \text{ moles}$$

So total amount of AgCl formed is 0.025 moles (based on the stoichiometry)

Amount of AgCl present in 0.025 moles of AgCl

$$= \text{no. of moles} \times \text{molar mass}$$

$$= 0.025 \times 143.5 = 3.59 \text{ g}$$

21. No. of moles of a gas that occupies a volume of 612.5 mL at room temperature and pressure (25°C and 1 atm pressure)

$$= 612.5 \times 10^{-3} \text{ L} / 24.5 \text{ L mol}^{-1}$$

$$= 0.025 \text{ moles}$$

We know that,

$$\text{Molar mass} = \text{Mass} / \text{no. of moles}$$

$$= 1.1 \text{ g} / 0.025 \text{ mol} = 44 \text{ g mol}^{-1}$$

22. No. of moles of carbon present in 6 g of C-12 = Mass / Molar mass

$$= 6/12 = 0.5 \text{ moles} = 0.5 \times 6.022 \times 10^{23} \text{ carbon atoms.}$$

No. of moles in 8 g of methane = $8 / 16 = 0.5$ moles

$$= 0.5 \times 6.022 \times 10^{23} \text{ carbon atoms.}$$

No. of moles in 7.5 g of ethane = $7.5 / 30 = 0.25$ moles

$$= 2 \times 0.25 \times 6.022 \times 10^{23} \text{ carbon atoms.}$$

23. Percentage of carbon in ethylene (C_2H_4) = $\frac{\text{mass of carbon}}{\text{Molar mass}} \times 100$
$$= \frac{24}{28} \times 100 = 85.71\%$$

$$\text{Percentage of carbon in propene (C}_3\text{H}_6) = \frac{36}{42} \times 100 = 85.71\%$$

24. (a) relative atomic mass of C-12 is 12 u

25. ${}_6\text{C}^{12}$

II Key to brief answer questions:

32. Given :

The density of CO_2 at 273 K and 1 atm pressure = 1.965 kgm^{-3}

Molar mass of CO_2 = ?

At 273 K and 1 atm pressure, 1 mole of CO_2 occupies a volume of 22.4 L

$$\begin{aligned} \text{Mass of 1 mole of CO}_2 &= \frac{1.965 \text{ Kg}}{1 \text{ m}^3} \times 22.4 \text{ L} \\ &= \frac{1.965 \times 10^3 \text{ g} \times 22.4 \times 10^{-3} \text{ m}^3}{1 \text{ m}^3} \\ &= 44.01 \text{ g} \end{aligned}$$

Molar mass of CO_2 = 44 gmol^{-1}

33.

Compound		Given no. of moles	No. of oxygen atoms
Ethanol	- C ₂ H ₅ OH	1	$1 \times 6.022 \times 10^{23}$
Formic acid	- HCOOH	1	$2 \times 6.022 \times 10^{23}$
Water	- H ₂ O	1	$1 \times 6.022 \times 10^{23}$
Answer : Formic acid			

$$\begin{aligned}
 34. \quad \text{Average atomic mass} &= \frac{(78.99 \times 23.99) + (10 \times 24.99) + (11.01 \times 25.98)}{100} \\
 &= \frac{2430.9}{100} \\
 &= 24.31 \text{ u}
 \end{aligned}$$

35. Reaction : $x + y + z_2 \rightarrow xyz_2$

Question	Number of moles of reactants allowed to react			Number of moles of reactants consumed during reaction			Limiting reagent
	x	y	z ₂	x	y	z ₂	
(a)	200 atoms	200 atoms	50 molecules	50 atoms	50 atoms	50 molecules	z ₂
(b)	1 mol	1 mol	3 mol	1 mol	1 mol	1 mol	x and y
(c)	50 atom	25 atom	50 molecules	25 atom	25 atom	25 molecules	y
(d)	2.5 mol	5 mol	5 mol	2.5 mol	2.5 mol	2.5 mol	x

$$\begin{aligned}
 36. \quad \text{Given: mass of one atom} &= 6.645 \times 10^{-23} \text{ g} \\
 \therefore \text{mass of 1 mole of atom} &= 6.645 \times 10^{-23} \text{ g} \times 6.022 \times 10^{23} \\
 &= 40 \text{ g} \\
 \therefore \text{number of moles of element in 0.320 kg} &= \frac{1 \text{ mole}}{40 \text{ g}} \times 0.320 \text{ kg} \\
 &= \frac{1 \text{ mol} \times 320 \text{ g}}{40 \text{ g}} \\
 &= 8 \text{ mol.}
 \end{aligned}$$

38.

Compound	Molecular formula	Empirical formula
Fructose	C ₆ H ₁₂ O ₆	CH ₂ O
Caffeine	C ₈ H ₁₀ N ₄ O ₂	C ₄ H ₅ N ₂ O

39. Given $2\text{Al} + \text{Fe}_2\text{O}_3 \rightarrow \text{Al}_2\text{O}_3 + 2\text{Fe}$

	Reactants		Products	
	Al	Fe_2O_3	Al_2O_3	Fe
Amount of reactant allowed to react	324 g	1.12 kg	-	-
Number of moles allowed to react	$\frac{324}{27} = 12 \text{ mol}$	$\frac{1.12 \times 10^3}{160} = 7 \text{ mol}$	-	-
Stoichiometric Co-efficient	2	1	1	2
Number of moles consumed during reaction	12 mol	6 mol	-	-
Number of moles of reactant unreacted and number of moles of product formed	-	1 mol	6 mol	12 mol

Molar mass of Al_2O_3 formed



$$= 6 \text{ mol} \times 102 \text{ g mol}^{-1} \left[\begin{array}{l} \text{Al}_2\text{O}_3 \\ (2 \times 27) + 3(\times 16) \\ 54 + 48 = 102 \end{array} \right] = 612 \text{ g}$$

Excess reagent = Fe_2O_3

Amount of excess reagent left at the end of the reaction = $1 \text{ mol} \times 160 \text{ g mol}^{-1}$

$$= 160 \text{ g} \left[\begin{array}{l} \text{Fe}_2\text{O}_3 \\ (2 \times 56) + (3 \times 16) \\ 112 + 48 = 160 \end{array} \right] = 160 \text{ g}$$

40. Balanced equation for the combustion of ethane






To produce 4 moles of CO_2 , 2 moles of ethane is required

\therefore To produce 1 mole (44 g) of CO_2 required

$$\begin{aligned}
 \text{number of moles of ethane} &= \frac{2 \text{ mol ethane}}{4 \text{ mol CO}_2} \times 1 \text{ mol CO}_2 \\
 &= \frac{1}{2} \text{ mole of ethane} \\
 &= \mathbf{0.5 \text{ mole of ethane}}
 \end{aligned}$$

41.



42.

Element	Percentage	Atomic mass	Relative number of atoms	Simple ratio	Whole no.
C	76.6	12	$\frac{76.6}{12} = 6.38$	$\frac{6.38}{1.06} = 6$	6
H	6.38	1	$\frac{6.38}{1} = 6.38$	$\frac{6.38}{1.06} = 6$	6
O	17.02	16	$\frac{17.02}{16} = 1.06$	$\frac{1.06}{1.06} = 1$	1

$$\text{Empirical formula} = \text{C}_6 \text{H}_6 \text{O}$$

$$n = \frac{\text{Molar mass}}{\text{Calculated empirical formula mass}}$$

$$= \frac{2 \times \text{vapour density}}{94} = \frac{2 \times 47}{94} = 1$$

$$\therefore \text{molecular formula } (\text{C}_6\text{H}_6\text{O}) \times 1 = \text{C}_6\text{H}_6\text{O}$$

43.

Element	%	Relative no. of atoms	Simple ratio
Na	14.31	$\frac{14.31}{23} = 0.62$	$\frac{0.62}{0.31} = 2$
S	9.97	$\frac{9.97}{32} = 0.31$	$\frac{0.31}{0.31} = 1$
H	6.22	$\frac{6.22}{1} = 6.22$	$\frac{6.22}{0.31} = 20$
O	69.5	$\frac{69.5}{16} = 4.34$	$\frac{4.34}{0.31} = 14$

∴ Empirical formula = Na₂ S H₂₀ O₁₄






$$n = \frac{\text{molar mass}}{\text{calculated empirical formula mass}} = \frac{322}{322} = 1 \quad \left[\begin{array}{l} \text{Na}_2 \text{S H}_{20} \text{O}_{14} \\ = (2 \times 23) + (1 \times 32) + (20 \times 1) + 14(16) \\ = 46 + 32 + 20 + 224 \\ = 322 \end{array} \right]$$

Molecular formula = Na₂ S H₂₀ O₁₄

Since all the hydrogen in the compound present as water





∴ Molecular formula is Na₂ SO₄ · 10H₂O

44. (i)













(ii)






(iii)








(iv)







45. (i) Half reaction are





(ii)







(iii)



(iv) Half reactions are





Unit - 2

Quantum Mechanical Model of Atom

Evaluate yourself:

- 1) Given: accelerated potential = 1 keV

The kinetic energy of the electron =
the energy due to accelerating potential.

$$\frac{1}{2}mv^2 = eV$$

$$mv^2 = 2eV$$

$$m^2v^2 = 2meV \Rightarrow (mv)^2 = 2meV$$

$$\Rightarrow mv = \sqrt{2meV}$$

$$\text{de Broglie wavelength } \lambda = \frac{h}{mv}$$

$$\Rightarrow \lambda = \frac{h}{\sqrt{2meV}}$$

$$m = \text{mass of the electron} = 9.1 \times 10^{-31} \text{ kg}$$

$$h = \text{Planck constant} = 6.626 \times 10^{-34} \text{ Js}$$

$$1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$$

$$\lambda = \frac{6.626 \times 10^{-34} \text{ Js}}{\sqrt{2 \times 9.1 \times 10^{-31} \text{ kg} \times 1 \text{ keV}}}$$

$$\lambda = \frac{6.626 \times 10^{-34} \text{ Js}}{\sqrt{2 \times 9.1 \times 10^{-31} \times 1 \times 10^3 \times 1.6 \times 10^{-19} \text{ kgJ}}}$$

$$\lambda = 3.88 \times 10^{-11} \text{ m}$$

$$[\because \frac{\text{Js}}{\sqrt{\text{J kg}}} = \text{J}^{1/2} \text{kg}^{-1/2} \cdot \text{s}]$$

$$= (\text{kgm}^2 \text{s}^{-2})^{1/2} \cdot \text{kg}^{-1/2} \text{s} \\ = \text{m}]$$

2. Given

$$\Delta v = 5.7 \times 10^5 \text{ ms}^{-1} \quad \Delta x = ?$$

According to Heisenberg's uncertainty principle, $\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$

$$\frac{h}{4\pi} = \frac{6.626 \times 10^{-34}}{4 \times 3.14} \text{ kgm}^2 \text{s}^{-1}$$

$$\frac{h}{4\pi} = 5.28 \times 10^{-35}$$

$$\Delta x \cdot \Delta p \geq 5.28 \times 10^{-35}$$

$$\Delta x \cdot m \cdot \Delta v \geq 5.28 \times 10^{-35}$$

$$\Rightarrow \Delta x \geq \frac{5.28 \times 10^{-35} \text{ kgm}^2 \text{s}^{-1}}{9.1 \times 10^{-31} \text{ kg} \times 5.7 \times 10^5 \text{ ms}^{-1}}$$

$$\Rightarrow \Delta x \geq 1.017 \times 10^{-10} \text{ m}$$

3. $n = 4$

$l = 0, 1, 2, 3$

\therefore 4 sub shells s, p, d & f.

$l = 0 \ m_l = 0 \Rightarrow$ one 4s orbital.

$l = 1 \ m_l = -1, 0, +1 \Rightarrow$ three 4p orbitals.

$l = 2 \ m_l = -2, -1, 0, +1, +2 \Rightarrow$ five 4d orbitals.

$l = 3 \ m_l = -3, -2, -1, 0, +1, +2, +3 \Rightarrow$ seven 4f orbitals.

Over all 16 orbitals are possible.

4.

Orbital	n	l	Radial node $n - l - 1$	Angular node l	Total node $n - 1$
3d	3	2	0	2	2
4f	4	3	0	3	3

5. $E_n = \frac{-13.6}{n^2} \text{eV}$

Second excited state

$n = 3 \quad \therefore E_3 = \frac{-13.6}{9} \text{eV}$
 $E_3 = -1.51 \text{eV}$

6. Electronic configuration of Fe^{3+} is $1s^2 2s^2 2p^6 3s^2 3p^6 4s^0 3d^5$

Five unpaired electrons.

1	1	1	1	1
---	---	---	---	---

$3d^5$


Electronic configuration of Mn^{2+} is $1s^2 2s^2 2p^6 3s^2 3p^6 4s^0 3d^5$

\therefore **Five unpaired electrons.**

Electronic configuration of Ar: $1s^2 2s^2 2p^6 3s^2 3p^6$

no unpaired electrons.

7.





$$n = 4 ; f \text{ orbital } l = 3 \Rightarrow m_l = -3, -2, -1, 0, +1, +2, +3$$

Out of two electrons, one electron occupies 4f orbital with $m_l = -3$ and another electron occupies 4f orbital with $m_l = -2$.

\therefore All the four quantum numbers for the two electrons are

Electron	n	l	m_l	m_s
$1e^-$	4	3	-3	+1/2
$2e^-$	4	3	-2	+1/2

8.

Electronic configuration of Fe^{3+} : $1s^2 2s^2 2p^6 3s^2 3p^6 4s^0 3d^5$

Electronic configuration of Ni^{2+} : $1s^2 2s^2 2p^6 3s^2 3p^6 4s^0 3d^8$

Fe^{3+} has stable $3d^5$ half filled configuration.

EVALUATION





I Choose the best answer:

- (c) 30
- (c) $4.42 \times 10^{-18} \text{ J}$
- (b) $\lambda_1 = 2\lambda_2$
- (d) Stark effect
- (b)
- (d) $n=6$ to $n=5$
- (a)
- (c) $d_{z^2}, d_{x^2-y^2}$
- (b) Spin quantum number
- (b) $[\text{Xe}]4f^7 6s^2$; $[\text{Xe}]4f^7 5d^1 6s^2$; $[\text{Xe}]4f^9 6s^2$
- (c) $4l+2$
- (d) $\sqrt{6} h/2\pi$
- (c) 2
- (c)
- (a) 9
- (a) $ns \rightarrow (n-2)f \rightarrow (n-1)d \rightarrow np$
- (b) (ii),(iv) & (v)
- (b) 17
- (a) Zero
- (c) $1/2m \sqrt{h/\pi}$
- (c) $6.6 \times 10^{-31} \text{ cm}$
- (d) 0.4
- (d) $-9E$
- (a) $\hat{H}\psi = E\psi$
- (d) $\Delta E \cdot \Delta X \geq h/4\pi$

Key to multiple choice questions:

1.



Atomic number = 26

Mass number = 56

No. of neutrons = 56 - 26 = 30

$$2. E = h\nu = hc/\lambda$$

$$= \frac{6.626 \times 10^{-34} \text{ J s} \times 3 \times 10^8 \text{ m s}^{-1}}{45 \times 10^{-9} \text{ m}}$$

$$= 4.42 \times 10^{-18} \text{ J}$$

3.

$$\frac{E_1}{E_2} = \frac{25 \text{ eV}}{50 \text{ eV}} = \frac{1}{2}$$

$$\frac{hc}{\lambda_1} \times \frac{\lambda_2}{hc} = \frac{1}{2}$$

$$2\lambda_2 = \lambda_1$$

4. Splitting of spectral lines in magnetic field is called Zeeman effect and splitting of spectral lines in electric field, is called Stark effect.

5. **Correct statement:** For $n=1$, the electron has more negative energy than it does for $n=6$ which means that the electron is **strongly** bound in the smallest allowed orbit.

6. $n = 6$ to $n=5$

$$E_6 = -13.6 / 6^2 ; \quad E_5 = -13.6 / 5^2$$

$$E_6 - E_5 = (-13.6 / 6^2) - (-13.6 / 5^2)$$

$$= 0.166 \text{ eV atom}^{-1}$$

$$E_5 - E_4 = (-13.6 / 5^2) - (-13.6 / 4^2)$$

$$= 0.306 \text{ eV atom}^{-1}$$

9. Spin quantum number

For the first electron $m_s = +\frac{1}{2}$

For the second electron $m_s = -\frac{1}{2}$

10. Eu : [Xe] $4f^7, 5d^0, 6s^2$

Gd : [Xe] $4f^7, 5d^1, 6s^2$

Tb: [Xe] $4f^9, 5d^0, 6s^2$

$$11. 2(2l+1) = 4l+2$$

12. Orbital angular momentum = $\sqrt{l(l+1)} h/2\pi$ For d orbital = $\sqrt{(2 \times 3)} h/2\pi = \sqrt{6} h/2\pi$

13. $n = 3; l=1; m = -1$

either $3p_x$ or $3p_y$

i.e., Maximum two electrons can be accommodated either in $3p_x$ or in $3p_y$

14. No. of radial node = $n-l-1$

No. of angular node = l

for 3p orbital

No. of angular node = $l = 1$

No. of radial node = $n-l-1 = 3-1-1 = 1$

15. $n = 3; l = 0; m_l = 0$ - one s orbital



$n = 3; l = 1; m_l = -1, 0, 1$ - three p orbitals

$n = 3; l = 2; m_l = -2, -1, 0, 1, 2$ - five d Orbitals

Overall nine orbitals are possible.

16. $n = 6$

According Aufbau principle,



17 (ii) l can have the values from 0 to $n-1$

$n=2$; possible ' l ' values are 0, 1 hence $l = 2$ is not possible.

(iv) for $l=0$; $m=-1$ not possible

(v) for $n=3$ $l = 4$ and $m = 3$ not possible

18. $n+l = 8$

Electronic configuration of atom with atomic number 105 is $[Rn] 5f^{14} 6d^3 7s^2$

Orbital	($n+l$)	No. of electrons
5f	$5+3 = 8$	14
6d	$6+2 = 8$	3
7s	$7+0 = 7$	2
No of electrons = $14 + 3 = 17$		

19. Option (a) - Zero (Refer to Figure 2.9)

20.

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\Delta p \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\Delta p^2 \geq \frac{h}{4\pi}$$

$$m^2 (\Delta v)^2 \geq \frac{h}{4\pi}$$

$$(\Delta v) \geq \sqrt{\frac{h}{4\pi m^2}}$$

$$\Delta v \geq \frac{1}{2m} \sqrt{\frac{h}{\pi}}$$

21.

$$m = 100 \text{ g} = 100 \times 10^{-3} \text{ kg}$$

$$v = 100 \text{ cm s}^{-1} = 100 \times 10^{-2} \text{ m s}^{-1}$$

$$\lambda = \frac{h}{mv}$$

$$= \frac{6.626 \times 10^{-34} \text{ Js}^{-1}}{100 \times 10^{-3} \text{ kg} \times 100 \times 10^{-2} \text{ ms}^{-1}}$$

$$= 6.626 \times 10^{-33} \text{ ms}^{-1}$$

$$= 6.626 \times 10^{-31} \text{ cms}^{-1}$$

23.

$$E_n = \frac{-13.6}{n^2} \text{ eV atom}^{-1}$$

$$E_1 = \frac{-13.6}{(1)^2} = -13.6$$

$$E_3 = \frac{-13.6}{(3)^2} = -\frac{13.6}{9}$$

Given that

$$E_3 = -E$$

$$-\frac{13.6}{9} = -E$$

$$-13.6 = -9E$$

$$\Rightarrow E_1 = -9E$$

II Key to brief answer question:

27. $n = 4$ $l = 0, 1, 2, 3$

four sub-shells \Rightarrow s, p, d, f

$l = 0$ $m_l = 0$; one 4s orbital.

$l = 1$ $m_l = -1, 0, +1$; three 4p orbitals.

$l = 2$ $m_l = -2, -1, 0, +1, +2$; five 4d orbitals.

$l = 3$ $m_l = -3, -2, -1, 0, +1, +2, +3$; seven 4f orbitals

Over all Sixteen orbitals.

28.

Orbital	n	l	Radial node $n - l - 1$	Angular node l
2s	2	0	1	0
4p	4	1	2	1
5d	5	2	2	2
4f	4	3	0	3

30. i) ground state

1	1	1	1	1
---	---	---	---	---

ii) maximum exchange energy

1	1	1	1	1
---	---	---	---	---

32.

Orbital	n	l
3p _x	3	1
4d _{x²-y²}	4	2

34.

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\Delta x \cdot \Delta p \geq 5.28 \times 10^{-35} \text{ Kg m}^2 \text{ s}^{-1}$$

$$\Delta x \cdot (m\Delta v) \geq 5.28 \times 10^{-35} \text{ Kg m}^2 \text{ s}^{-1}$$

$$\text{Given } \Delta v = 0.1\%$$

$$v = 2.2 \times 10^6 \text{ ms}^{-1}$$

$$m = 9.1 \times 10^{-31} \text{ Kg}$$


$$\Delta v = \frac{0.1}{100} \times 2.2 \times 10^6 \text{ ms}^{-1}$$

$$= 2.2 \times 10^3 \text{ ms}^{-1}$$

$$\therefore \Delta x \geq \frac{5.28 \times 10^{-35} \text{ Kg m}^2 \text{ s}^{-1}}{9.1 \times 10^{-31} \text{ Kg} \times 2.2 \times 10^3 \text{ ms}^{-1}}$$

$$\Delta x \geq 2.64 \times 10^{-8} \text{ m}$$


35. Electronic configuration of oxygen



\therefore 8th electron present in 2p_x orbital
and the quantum numbers are

$$n = 2, l = 1, m_l = \text{either } +1 \text{ or } -1 \text{ and } s = -1/2$$

Electronic configuration of chlorine =



15th electron present in 3p_z orbital and
the quantum numbers are $n = 3, l = 1,$
 $m_l = \text{either } +1 \text{ or } -1 \text{ and } m_s = +1/2$

36.

$$E_n = \frac{-13.6}{n^2} \text{ eV atom}^{-1}$$

$$n = 3 \quad E_3 = \frac{-13.6}{3^2} = \frac{-13.6}{9} = -1.51 \text{ eV atom}^{-1}$$

$$n = 4 \quad E_4 = \frac{-13.6}{4^2} = \frac{-13.6}{16} = -0.85 \text{ eV atom}^{-1}$$

$$\Delta E = (E_4 - E_3) = (-0.85) - (-1.51)$$

$$\text{eV atom}^{-1}$$

$$= (-0.85 + 1.51)$$

$$= \mathbf{0.66 \text{ eV atom}^{-1}}$$

$$(1 \text{ eV} = 1.6 \times 10^{-19} \text{ J})$$

$$\Delta E = 0.66 \times 1.6 \times 10^{-19} \text{ J}$$

$$\Delta E = 1.06 \times 10^{-19} \text{ J}$$

$$h\nu = 1.06 \times 10^{-19} \text{ J}$$

$$\frac{hc}{\lambda} = 1.06 \times 10^{-19} \text{ J}$$

$$\therefore \lambda = \frac{hc}{1.06 \times 10^{-19} \text{ J}}$$

$$= \frac{6.626 \times 10^{-34} \text{ Js} \times 3 \times 10^8 \text{ ms}^{-1}}{1.06 \times 10^{-19} \text{ J}}$$

$$\lambda = 1.875 \times 10^{-6} \text{ m}$$

37. Given : de Broglie wavelength of the tennis ball equal to 5400 \AA .

$$m = 54 \text{ g}$$

$$v = ?$$

$$\lambda = \frac{h}{mV}$$

$$V = \frac{h}{m\lambda}$$

$$V = \frac{6.626 \times 10^{-34} \text{ Js}}{54 \times 10^{-3} \text{ Kg} \times 5400 \times 10^{-10} \text{ m}}$$

$$V = 2.27 \times 10^{-26} \text{ ms}^{-1}$$


38.

n	l	Sub Energy levels	m_l values	Number of orbitals
4	2	4d	-2, -1, 0, +1, +2	five 4d orbitals
5	3	5f	-3, -2, -1, 0, +1, +2, +3	seven 5f orbitals
7	0	7s	0	one 7s orbitals

41. no. of electrons : 35 (given)

no. of protons : 35

Electronic configuration



$\uparrow\downarrow$	$\uparrow\downarrow$	\uparrow
----------------------	----------------------	------------

$4P_x \quad 4P_y \quad 4P_z$

last electron present in $4P_y$ orbital

$n = 4 \quad l = 1 \quad m_l = \text{either } +1 \text{ or } -1$
and $s = -1/2$

43. $\text{He}^+ \rightarrow \text{He}^{2+} + e^-$

$$E_n = \frac{-13.6Z^2}{n^2}$$

$$E_1 = \frac{-13.6(2)^2}{(1)^2} = -54.4$$

$$E_\infty = \frac{-13.6(2)^2}{(\infty)^2} = 0$$

\therefore Required Energy for the given process = $E_\infty - E_1 = 0 - (-54.4) = 54.4 \text{ eV}$

44.

	Atom	Uni-negative ion
number of electron	$x - 1$	x
number of protons	$x - 1$	$x - 1$
number of neutrons	y	y

Given that, $y = x + 11.1\% \text{ of } x$

$$= \left(x + \frac{11.1}{100} x \right) = x + 0.111x$$

$$y = 1.111x$$

mass number = 37

number of protons + number of neutrons = 37

$$(x - 1) + 1.111x = 37$$

$$x + 1.111x = 38$$

$$2.111x = 38$$

$$x = \frac{38}{2.11}$$

$$x = 18.009 \quad x = 18 \text{ (whole number)}$$

$$\therefore \text{Atomic number} = x - 1$$

$$= 18 - 1 = 17$$

Mass number = 37

Symbol of the ion ${}^{37}_{17}\text{Cl}^-$

45)

$$r_n = \frac{(0.529)n^2}{z} \text{ \AA} \quad E_n = \frac{-13.6 (z^2)}{n^2} \text{ eV atom}^{-1}$$

for Li^{2+} $z = 3$ Bohr radius for the third orbit (r_3)

$$= \frac{(0.529)(3)^2}{3}$$

$$= 0.529 \times 3$$

$$= 1.587 \text{ \AA}$$

Energy of an electron in the fourth orbit (E_4)

$$= \frac{-13.6 (3)^2}{(4)^2}$$

$$= -7.65 \text{ eV atom}^{-1}$$

46) Given

$$v = 2.85 \times 10^8 \text{ ms}^{-1}$$

$$m_p = 1.673 \times 10^{-27} \text{ Kg}$$

$$\lambda = \frac{h}{mv}$$

$$= \frac{6.626 \times 10^{-34} \text{ kgm}^2 \text{ s}^{-1}}{1.673 \times 10^{-27} \text{ kg} \times 2.85 \times 10^8 \text{ ms}^{-1}}$$

$$\lambda = 1.389 \times 10^{-15} \Rightarrow \lambda = 1.389 \times 10^{-5} \text{ \AA}$$

$$[\because 1 \text{ \AA} = 10^{-10} \text{ m}]$$

47) $m = 160 \text{ g} = 160 \times 10^{-3} \text{ kg}$

$$v = 140 \text{ km hr}^{-1} = \frac{140 \times 10^3}{60 \times 60} \text{ ms}^{-1}$$

$$v = 38.88 \text{ ms}^{-1}$$

$$\lambda = \frac{h}{mv}$$

$$\lambda = \frac{6.626 \times 10^{-34} \text{ Kg m}^2 \text{ s}^{-1}}{160 \times 10^{-3} \text{ Kg} \times 38.88 \text{ ms}^{-1}}$$

$$\lambda = 1.065 \times 10^{-34} \text{ m}$$

48) $\Delta x = 0.6 \text{ \AA} = 0.6 \times 10^{-10} \text{ m}$

$$\Delta p = ?$$

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\Delta x \cdot \Delta p \geq 5.28 \times 10^{-35} \text{ kgm}^2 \text{ s}^{-1}$$

$$(0.6 \times 10^{-10}) \Delta p \geq 5.28 \times 10^{-35}$$

$$\Rightarrow \Delta p \geq \frac{5.28 \times 10^{-35} \text{ kgm}^2 \text{ s}^{-1}}{0.6 \times 10^{-10} \text{ m}}$$

$$\Delta p \geq 8.8 \times 10^{-25} \text{ kgms}^{-1}$$

49)

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\lambda \cdot (m\Delta v) \geq \frac{h}{4\pi}$$

$$\Delta v \geq \frac{h}{4\pi(m\lambda)}$$

$$\Delta v \geq \frac{h}{4\pi \times m \times \frac{h}{mv}} \quad \left[\because \lambda = \frac{h}{mv} \right]$$

$$\Delta v \geq \frac{v}{4\pi}$$

Therefore,

$$\text{minimum uncertainty in velocity} = \frac{v}{4\pi}$$

50) Potential difference = 100V

$$= 100 \times 1.6 \times 10^{-19} \text{ J}$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$= \frac{6.626 \times 10^{-34} \text{ Kg m}^2 \text{ s}^{-1}}{\sqrt{2 \times 9.1 \times 10^{-31} \text{ Kg} \times 100 \times 1.6 \times 10^{-19} \text{ J}}}$$

$$\lambda = 1.22 \times 10^{-10} \text{ m}$$

51)

n	l	m_l	sub energy level
4	2	0	4d
3	1	0	3p
5	1	any one value -1, 0, +1	5p
3	2	-2	3d

Unit - 3

Periodic classification of elements

Evaluate yourself

2. Atomic number : 120

IUPAC temporary symbol :
Unbinilium

IUPAC temporary symbol : Ubn

Possible electronic configuration
:[Og] 8s²

3. Electronic Configuration : (n - 1)d² ns²

for n = 5, the electronic configuration is,

1s² 2s² 2p⁶ 3s² 3p⁶ 4s² 3d¹⁰ 4p⁶ 4d² 5s²
Atomic number : 40

4th group 5th period (d block element)
= Zirconium

4. Electronic Configuration of Aluminium

¹³Al $\underbrace{1s^2}_{(n-2)} \underbrace{2s^2 2p^6}_{(n-1)} \underbrace{3s^2 3p^1}_n$

Group	no. of electrons	Contribution of each electron to 'S' value	Contribution of a particular group
n	2	0.35	0.70
(n - 1)	8	0.85	6.80
(n - 2)	2	1	2.00
			9.50

∴ Effective nuclear charge = Z - S

$$= 13 - 9.5$$

$$(Z_{\text{eff}})_{\text{Al}} = 3.5$$

Electronic Configuration of chlorine

$\underbrace{1s^2}_{n-2} \underbrace{2s^2 2p^6}_{n-1} \underbrace{3s^2 3p^5}_n$

Group	no. of electrons	Contribution of each electron to 'S' value	Contribution of a particular group
n	6	0.35	2.1
(n - 1)	8	0.85	6.8
(n - 2)	2	1	2
		S =	10.9

∴ Effective nuclear charge = Z - S

$$= 17 - 10.9$$

$$(Z_{\text{eff}})_{\text{Cl}} = 6.1$$

$(Z_{\text{eff}})_{\text{Cl}} > (Z_{\text{eff}})_{\text{Al}}$ and. Hence $r_{\text{Cl}} < r_{\text{Al}}$.

5.

X³⁺, Y²⁺, Z⁻ are isoelectronic.

∴ Effective nuclear charge is in the order $(Z_{\text{eff}})_{\text{Z}^-} < (Z_{\text{eff}})_{\text{Y}^{2+}} < (Z_{\text{eff}})_{\text{X}^{3+}}$ and

Hence, ionic radii should be in the order

$$r_{\text{Z}^-} > r_{\text{Y}^{2+}} > r_{\text{X}^{3+}}$$

∴ The correct values are,

Species	Ionic radii
Z ⁻	136
Y ²⁺	64
X ³⁺	49


6. **Noble gases** : Ionization energy ranging from 2372 KJmol⁻¹ to 1037 kJ mol⁻¹.

For element X, the IE₁ value is in the

range of noble gas, moreover for this element both IE_1 and IE_2 are higher and hence X is the noble gas.

For Y, the first ionisation energy is low and second ionisation energy is very high and hence Y is most reactive metal.

For Z, both IE_1 and IE_2 are higher and hence it is least reactive.




For one mole (35.5g) 348 kJ is released.

\therefore For 17.75g chlorine,
 $\frac{348 \text{ kJ}}{35.5 \text{ g}} \times 17.75 \text{ g}$ energy leased.


$$\therefore \text{The amount of energy released} = \frac{348}{2} = 174 \text{ kJ}$$

Activitiy : 3.1


Covalent radii - 2nd group elements




Covalent radii - 17th group elements



Covalent radii - 3rd period elements



Covalent radii - 4th period elements





EVALUATION

I Choose the best answer:

1. (d) bibibium
2. (b) AB_2
3. (d) f-block elements
4. (a) $I < Br < Cl < F$ (increasing electron gain enthalpy)
5. (d) fluorine
6. (c) Aluminium
7. (b) $Na < Al < Mg < Si < P$
8. (a)
9. (d) $Ca < Al < C < O < F$
10. (c) $Cl > F > Br > I$
11. (d) Hydrogen
12. (c) Argon
13. (a) $Y > Z > X > A$
14. (c)
15. (a) $1s^2, 2s^2, 2p^6, 3s^1$
16. (c) Oxygen
17. (c) $+527 \text{ kcal mol}^{-1}$
18. (a) $s > p > d > f$
19. (d) None of these
20. (b) 575 kJ mol^{-1}
21. (a)
22. (a) Generally increases
23. (d) Be and Al

Unit - 4

Hydrogen

EVALUATION




I Choose the best answer:

- (c)
- (c) $\text{CO} + \text{H}_2$
- (b)
- (d) group one elements
- (c) $1p+2n$
- (a) Palladium, Vanadium
- (a)
- (a) 1.2 g
- (d) EDTA
- (c) CaCl_2
- (a) sodium aluminium silicate
- (a)
- (c) $\text{CrO}(\text{O}_2)_2$
- (c) $5/2$
- (d) 8.4
- (d) sp^3 and sp^3
- (c) monobasic acid
- (a) tetrahedrally by 4-H atoms
- (b) intra-molecular hydrogen bonding and inter molecular hydrogen bonding
- (c) both (a) & (b)
- (c) amphoteric oxide

Key for multiple choice questions:


- Option (c)
Correct statement : Hydrogen has three isotopes of which **protium** is the most common.
- Option (c)
 $\text{CO} + \text{H}_2$ - Water gas
- Option (b)
Correct statement : Ortho isomer - one nuclear spin
Para isomer - zero nuclear spin
- Option (d)
eg : Sodium hydride ($\text{Na}^+ \text{H}^-$)
- Option (c)
 ${}_1\text{T}^3 (1e^-, 1p, 2n)$
- Option (a)
 $\text{Ca}^{2+} + \text{Na}_2\text{CO}_3 \rightarrow \text{CaCO}_3 \downarrow + 2\text{Na}^+$
- mass of deuterium = $2 \times$ mass of protium
If all the 1.2 g hydrogen is replaced with deuterium, the weight will become 2.4g. Hence the increase in body weight is $(2.4 - 1.2 = 1.2 \text{ g})$
Option (a)
- EDTA (option (d))
- Permanent hardness if water is due to the presence of the chlorides, nitrates and sulphates of Ca^{2+} and Mg^{2+} ions.
Option (c) CaCl_2

- 11) Zeolite is sodium aluminium silicate.




Option (a)

- 12) (a) 1 mL of H_2O_2 will give 100ml O_2 at STP.



Option (c)




Option (c) 5/2 moles

- 15) Volume strength of hydrogen peroxide = Normality of hydrogen peroxide $\times 5.6$

$$= 1.5 \times 5.6$$

$$= 8.4$$

Option (d)



Volume strength of Hydrogen peroxide

$$\frac{\text{Normality} \times \text{Equivalent weight of } \text{H}_2\text{O}_2}{68} \times 22.4$$

$$= \text{Normality} \times \left(\frac{17 \times 22.4}{68} \right)$$


Volume strength of hydrogen peroxide

$$= \text{Normality} \times 5.6$$

- 16) sp^3 and sp^3

Option (d)

- 17) Hypophosphorus acid on reaction with D_2O , only one hydrogen is replaced by deuterium and hence it is mono basic




Option (c) monobasic acid


- 18) (a) tetrahedrally surrounded by 4 hydrogen atoms (refer 4.6 (a) Structure of ice)

19)

o-nitro phenol



p-nitro phenol



- 20) Option (c)

Heavy water is used as moderator as well as coolant in nuclear reactions.

- 21) Water is a amphoteric oxide.

Option (c)

Unit - 5

Alkali and Alkaline Earth Metals

EVALUATION




I Choose the best answer.

1. (c) Density: $\text{Li} < \text{K} < \text{Na} < \text{Rb} < \text{Cs}$
2. (a) Li^+ has minimum degree of hydration among alkali metal cations
3. (d) none of these
4. (b) Li
5. (c) kerosene
6. (a) superoxide and paramagnetic
7. (c) Potassium carbonate can be prepared by solvay process
8. (b) Magnesium
9. (b) $\text{MI} < \text{MBr} < \text{MCl} < \text{MF}$
10. (a) Castners process
11. (c) $\text{Ca}(\text{CN})_2$
12. (a) MgCl_2
13. (a) p-2, q-1, r-4, s-5, t-6, u-3
14. (d) both assertion and reason are false
15. (a)
16. (b) $\text{MgCO}_3 > \text{CaCO}_3 > \text{SrCO}_3 > \text{BaCO}_3$
17. (c) Its salts are rarely hydrolysed
18. (c) milk of lime
19. (b) NaHCO_3
20. (b) $\text{Ca}(\text{OH})_2$
21. (a) Ca^{2+} ions are not important in maintaining the regular beating of the heart.
22. (b) CaF_2
23. (a) $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$
24. (b) CaNCN
25. (d) Li_2CO_3

Keys to multiple choice questions:

- 1) Option (c)
Potassium is lighter than sodium
(Refer table 5.3)
The correct order of density is
 $\text{Li} < \text{K} < \text{Na} < \text{Rb} < \text{Cs}$
 $0.54 < 0.86 < 0.97 < 1.53 < 1.90$ (in g cm^{-3})
- 2) Option (a)
 Li^+ has maximum degree of hydration among alkali metal cations.
 $\text{Li}^+ > \text{Na}^+ > \text{K}^+ > \text{Rb}^+ > \text{Cs}^+$
- 3) All these compounds reacts with alkali metals to evolve hydrogen gas.
(d) none of these.
- 4) hydration energy of Li^+ is more and hence Li^+ is stabilized in aqueous medium.
(b) Li
- 5) (c) Kerosene

20)



21) Ca^{2+} ion plays an important role in maintaining regular heart beat.


Option (a)

22) 'Blue john' – CaF_2

(A variety of fluorite)

\therefore Option (b)

23) $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ \therefore Option (a)



\therefore Option (b)

25) Li_2CO_3 is least stable.

Option (d)

Unit - 6

Gaseous State

Evaluate yourself:

1. Volume of freon (V_1) = 1.5 dm^3

Pressure (P_1) = 0.3 atm

'T' is constant

$P_2 = 1.2 \text{ atm}$

$V_2 = ?$

$\therefore P_1 V_1 = P_2 V_2$

$$\Rightarrow V_2 = \frac{P_1 V_1}{P_2} = \frac{0.3 \text{ atm} \times 1.5 \text{ dm}^3}{1.2 \text{ atm}} = 0.375 \text{ dm}^3$$

\therefore Volume decreased from 1.5 dm^3 to 0.375 dm^3

2. $V_1 = 0.375 \text{ dm}^3$ $V_2 = 0.125$

$P_1 = 1.05 \text{ atm}$ $P_2 = ?$

'T' - Constant

$P_1 V_1 = P_2 V_2$

$$\therefore P_2 = \frac{P_1 V_1}{V_2} = \frac{1.05 \times 0.375}{0.125} = 3.15 \text{ atm}$$

3. $V_1 = 3.8 \text{ dm}^3$ $T_2 = 0^\circ\text{C} = 273 \text{ K}$

$T_1 = ?$ $V_2 = 2.27 \text{ dm}^3$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \quad T_1 = \left(\frac{T_2}{V_2} \right) \times V_1 = \frac{273 \text{ K}}{2.27 \text{ dm}^3} \times 3.8 \text{ dm}^3 = 457 \text{ K}$$

4. $V_1 = 7.05 \text{ dm}^3$ $V_2 = 2.35 \text{ dm}^3$

$n_1 = 0.312 \text{ mol}$ $n_2 = ?$

'P' and 'T' are constant

$$\therefore \frac{V_1}{n_1} = \frac{V_2}{n_2} \Rightarrow n_2 = \left(\frac{n_1}{V_1} \right) \times V_2 = \frac{0.312 \text{ mol}}{7.05 \text{ dm}^3} \times 2.35 \text{ dm}^3 = 0.104 \text{ mol}$$

Number of moles exhaled =

$0.312 - 0.104 = 0.208 \text{ moles}$

- 5) $T_1 = 8^\circ\text{C} = 8 + 273 = 281 \text{ K}$

$P_1 = 6.4 \text{ atm}$ $V_1 = 2.1 \text{ ml}$

$T_2 = 25^\circ\text{C} = 25 + 273 = 298 \text{ K}$

$P_2 = 1 \text{ atm}$ $V_2 = ?$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow V_2 = \left(\frac{P_1 V_1}{T_1} \right) \times \frac{T_2}{P_2} = \frac{6.4 \text{ atm} \times 2.1 \text{ ml}}{281 \text{ K}} \times \frac{298 \text{ K}}{1 \text{ atm}} = 14.25 \text{ ml}$$

6(a)

$V_{\text{O}_2} = 12 \text{ dm}^3$ $P = 1 \text{ atm}$

$V_{\text{He}} = 46 \text{ dm}^3$

$V_{\text{total}} = 5 \text{ dm}^3$

$$P_{\text{O}_2} = x_{\text{O}_2} \times P_{\text{total}} \quad n_{\text{O}_2} = \frac{1 \text{ mol}}{22.4 \text{ L}} \times 12 \text{ L} = 0.54 \text{ mol}$$

$$x_{\text{O}_2} = \frac{n_{\text{O}_2}}{n_{\text{O}_2} + n_{\text{He}}} = \frac{0.54}{0.54 + 2.05} = 0.21$$

$$n_{\text{He}} = \frac{1 \text{ mol}}{22.4 \text{ L}} \times 46 \text{ L} = 2.05 \text{ mol}$$



$$P_{\text{total}} \times V_{\text{total}} = 1 \text{ atm} \times 22.4 \text{ l}$$

$$\therefore P_{\text{total}} = \frac{1 \text{ atm} \times 22.4 \cancel{\text{l}}}{5 \cancel{\text{l}}}$$

$$P_{\text{total}} = 4.48 \text{ atm}$$

$$\begin{aligned} \therefore P_{\text{O}_2} &= 0.21 \times 4.48 \text{ atm} \\ &= 0.94 \text{ atm} \end{aligned}$$


$$P_{\text{He}} = x_{\text{He}} \times P_{\text{total}}$$

$$\begin{aligned} x_{\text{He}} &= \frac{n_{\text{He}}}{n_{\text{O}_2} + n_{\text{He}}} \\ &= \frac{2.05}{0.54 + 2.05} \end{aligned}$$

$$x_{\text{He}} = \frac{2.05}{2.59} = 0.79$$

$$\therefore P_{\text{He}} = 0.79 \times 4.48 \text{ atm}$$

$$P_{\text{He}} = 3.54 \text{ atm}$$



$$P_{\text{total}} = 772 \text{ mm Hg}$$

$$P_{\text{H}_2\text{O}} = 26.7 \text{ mm Hg}$$

$$P_{\text{total}} = P_{\text{O}_2} + P_{\text{H}_2\text{O}}$$

$$\therefore P_{\text{O}_2} = P_{\text{total}} - P_{\text{H}_2\text{O}}$$

$$P_1 = 26.7 \text{ mm Hg} \quad T_2 = 295 \text{ K}$$

$$T_1 = 300 \text{ K} \quad P_2 = ?$$

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$\Rightarrow P_2 = \left(\frac{P_1}{T_1} \right) T_2 = \frac{26.7 \text{ mm Hg}}{300 \cancel{\text{K}}} \times 295 \cancel{\text{K}}$$

$$P_2 = 26.26 \text{ mm Hg}$$

$$\begin{aligned} \therefore P_{\text{O}_2} &= 772 - 26.26 \\ &= 745.74 \text{ mm Hg} \end{aligned}$$

$$7) \quad t_1 = 1.5 \text{ minutes} \quad (\text{gas})_{\text{hydro carbon}}$$

$$t_2 = 4.73 \text{ minutes} \quad (\text{gas})_{\text{Bromine}}$$

$$\frac{\gamma_{\text{Hydrocarbon}}}{\gamma_{\text{Bromine}}} = \frac{t_{\text{Bromine}}}{t_{\text{Hydrocarbon}}}$$

(\because Volume is constant)

$$= \frac{4.73 \text{ minutes}}{1.5 \text{ minutes}}$$

$$= 3.15$$

$$\frac{\gamma_{\text{Hydrocarbon}}}{\gamma_{\text{Bromine}}} = \sqrt{\frac{M_{\text{Bromine}}}{M_{\text{hydrocarbon}}}}$$

$$3.15 = \sqrt{\frac{159.8 \text{ g mol}^{-1}}{M_{\text{hydro carbon}}}}$$

Squaring on both sides and rearranging,

$$M_{\text{hydro carbon}} = \frac{159.8 \text{ g mol}^{-1}}{(3.15)^2}$$

$$M_{\text{hydro carbon}} = 16.1 \text{ g mol}^{-1}$$

$$\begin{aligned} n(12) + (2n+2)1 &= 16 & \left[\begin{array}{l} \because \text{general formula} \\ \text{for hydrocarbon} \end{array} \right] \\ 12n + 2n + 2 &= 16 & \left[\begin{array}{l} \text{C}_n\text{H}_{2n+2} \quad (\text{alkane}) \end{array} \right] \end{aligned}$$

$$14n = 16 - 2$$

$$14n = 14$$

$$n = 1$$

$$\therefore \text{The hydro carbon is } \text{C}_1\text{H}_{2(1)+2} = \text{CH}_4$$

8) Critical temperature of a gas is defined as the temperature above which it cannot be liquified even at high pressures.

\therefore When cooling starts from 700 K, H_2O will liquify first, then followed by ammonia and finally carbondioxide will liquify.



EVALUATION



I Choose the correct answer:

- (d) at high pressure the intermolecular interactions become significant
- (d) inversely proportional to the square root of its molecular weight
- (c) $\left(P + \frac{an^2}{V^2}\right)(V - nb) = nRT$
- (b) exert no attractive forces on each other
- (a) $1/3$
- (b) Boyle temperature
- (c) diffusion
- (b) near the hydrogen chloride bottle
- (d) units of pressure and volume
- (c) $8.3 \text{ J mol}^{-1} \text{ K}^{-1}$
- (a) Boyle's Law
- (c) NH_3
- (d) I, II and III
- (c) 0.41 dm^3
- (c) P
- (b) 4
- (c) $1/8$
- (b) $1/T$
- (a) P
- (b) NH_3
- (c) $\text{mol}^{-1} \text{ L}$ and $\text{L}^2 \text{ atm mol}^{-2}$

- (d) both assertion and reasons are false
- (c) 3.41 g L^{-1}
- (c)
- (d) HI

Keys to multiple choice questions:

- mass of methane
= mass of oxygen = a

$$\text{number of moles of methane} = \frac{a}{16}$$

$$\text{number of moles of Oxygen} = \frac{a}{32}$$

$$\begin{aligned}\text{mole fraction of Oxygen} &= \frac{\frac{a}{32}}{\frac{a}{16} + \frac{a}{32}} \\ &= \frac{\frac{a}{32}}{\frac{2a}{32} + \frac{a}{32}} \\ &= \frac{\frac{a}{32}}{\frac{3a}{32}} \\ &= \frac{1}{3}\end{aligned}$$

$$\begin{aligned}\text{Partial pressure of oxygen} &= \text{mole fraction} \times \text{Total Pressure} \\ &= \frac{1}{3}P\end{aligned}$$

- The temperature at which real gases obey the ideal gas laws over a wide range of pressure is called Boyle temperature

8. Rate of diffusion $\propto 1/\sqrt{M}$

$$M_{\text{NH}_3} = 17 ; M_{\text{HCl}} = 36.5$$

$$\gamma_{\text{NH}_3} > \gamma_{\text{HCl}}$$

Hence white fumes first formed near hydrogen chloride

12. Higher the value of 'a', greater the intermolecular force of attraction, easier the liquefaction.

option (c) is correct

- 14.

$$\text{Compressibility factor (z)} = \frac{PV}{nRT}$$

$$V = \frac{z \times nRT}{P}$$

$$\begin{aligned} & 0.8697 \times 1 \times 8.314 \\ & \times 10^{-2} \text{ bar dm}^3 \text{ K}^{-1} \text{ mol}^{-1} \\ & = \frac{\times 400 \text{ K}}{71 \text{ bar}} \end{aligned}$$

$$V = 0.41 \text{ dm}^3$$

15. $T_1 \quad T_2 = 2T_1$

$$V_1 \quad V_2 = 2V_1$$

$$P_1 \quad P_2 = ?$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_2 = \frac{P_1 V_1}{T_1} \times \frac{T_2}{V_2} = \frac{P_1 V_1}{T_1} \times \frac{2T_1}{2V_1}$$

$$P_2 = P_1$$

Option (c)

- 16.

$$\begin{aligned} \frac{\gamma_{\text{H}_2}}{\gamma_{\text{C}_n\text{H}_{2n+2}}} &= \sqrt{\frac{M_{\text{C}_n\text{H}_{2n+2}}}{M_{\text{H}_2}}} \\ 3\sqrt{3} &= \sqrt{\frac{M_{\text{C}_n\text{H}_{2n+2}}}{2}} \end{aligned}$$

Squaring on both sides and rearranging

$$27 \times 2 = M_{\text{C}_n\text{H}_{2n+2}}$$

$$54 = n(12) + (2n+2)(1)$$

$$54 = 12n + 2n + 2$$

$$54 = 14n + 2$$

$$n = (54+2)/14 = 56/14 = 4$$

- 17.

$$\begin{aligned} \frac{\gamma_{\text{O}_2}}{\gamma_{\text{H}_2}} &= \sqrt{\frac{M_{\text{H}_2}}{M_{\text{O}_2}}} = \sqrt{\frac{2}{32}} = \frac{1}{4} \\ \gamma_{\text{O}_2} &= \frac{1}{4} \gamma_{\text{H}_2} \end{aligned}$$

The fraction of oxygen that escapes in the time required for one half of the hydrogen to escape is 1/8

- 18.

$$\begin{aligned} & \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_P \quad [\text{For an ideal gas } PV = nRT] \\ &= \frac{1}{V} \left(\frac{\partial \left(\frac{nRT}{P} \right)}{\partial T} \right)_P \\ &= \frac{nR}{PV} \left(\frac{\partial T}{\partial T} \right) = \frac{nR}{nRT} = \frac{1}{T} \end{aligned}$$

19. Greater the 'a' value, easier the liquefaction

21.

$$an^2/V^2 = \text{atm}$$

$$a = \text{atm L}^2/\text{mol}^2 = \text{L}^2 \text{ mol}^{-2} \text{ atm}$$

$$nb = \text{L}$$

$$b = \text{L/mol} = \text{L mol}^{-1}$$

22. **Correct Statement:** Critical temperature of CO_2 is 304 K. It means that CO_2 cannot be liquefied above 304 K, whatever the pressure may applied.

Pressure is inversely proportional to volume

23.

$$\begin{aligned} \text{Density} &= \frac{\text{Mass}}{\text{Volume}} \\ &= \frac{m}{\left(\frac{nRT}{P}\right)} = \left(\frac{m}{n}\right) \frac{P}{RT} \\ &= \text{Molar mass} \times \frac{P}{RT} \\ &= \frac{28 \text{ g mol}^{-1} \times 5 \text{ atm}}{0.082 \text{ L atm K}^{-1} \text{ mol}^{-1} \times 500 \text{ K}} \\ &= 3.41 \text{ g L}^{-1} \end{aligned}$$

24. For a fixed mass of an ideal gas $V \propto T$

$$P \propto 1/V$$

$$\text{and } PV = \text{Constant}$$

25. At a given temperature and pressure

Volume \propto no. of moles

Volume \propto Mass / Molar mass

Volume \propto 28 / Molar mass

i.e. if molar mass is more, volume is less. Hence HI has the least volume

II Key to short answer questions:

46. $T_1 = 15^\circ \text{C} + 273$ $T_2 = 38 + 273$

$$T_1 = 288 \text{ K} \quad T_2 = 311 \text{ K}$$

$$V_1 = 2.58 \text{ dm}^3 \quad V_2 = ?$$

(P = 1 atm constant)

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\begin{aligned} V_2 &= \left(\frac{V_1}{T_1}\right) \times T_2 \\ &= \frac{2.58 \text{ dm}^3}{288 \text{ K}} \times 311 \text{ K} \end{aligned}$$

$V_2 = 2.78 \text{ dm}^3$ i.e. volume increased from 2.58 dm^3 to 2.78 dm^3

47. $V_1 = 8.5 \text{ dm}^3$ $V_2 = 6.37 \text{ dm}^3$

$$T_1 = ? \quad T_2 = 0^\circ \text{C} = 273 \text{ K}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$V_1 \times \left(\frac{T_2}{V_2}\right) = T_1$$

$$T_1 = 8.5 \text{ dm}^3 \times \frac{273 \text{ K}}{6.37 \text{ dm}^3}$$

$$T_1 = 364.28 \text{ K}$$

48. $n_A = 1.5 \text{ mol}$ $n_B = ?$

$$V_A = 37.6 \text{ dm}^3 \quad V_B = 16.5 \text{ dm}^3$$

(T = 298 K constant)

$$\frac{V_A}{n_A} = \frac{V_B}{n_B}$$

$$n_A = \left(\frac{n_A}{V_A}\right) V_B$$



$$\begin{aligned} &= \frac{1.5 \text{ mol}}{37.6 \text{ dm}^3} \times 16.5 \text{ dm}^3 \\ &= 0.66 \text{ mol} \end{aligned}$$

49. $n = 1.82 \text{ mole}$

$$V = 5.43 \text{ dm}^3$$

$$T = 69.5 + 273 = 342.5$$

$$P = ?$$

$$PV = nRT$$

$$P = \frac{nRT}{V}$$

$$1.82 \text{ mol} \times$$

$$0.0821 \text{ dm}^3 \text{ atm mol}^{-1} \text{ K}^{-1}$$

$$P = \frac{\times 342.5 \text{ K}}{5.43 \text{ dm}^3}$$

$$P = 9.425 \text{ atm}$$

50.

$$P_1 = 1.2 \text{ atm}$$

$$T_1 = 18^\circ \text{C} + 273 = 291 \text{ K}$$

$$T_2 = 85^\circ \text{C} + 273 = 358 \text{ K}$$

$$P_2 = ?$$

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$P_2 = \left(\frac{P_1}{T_1} \right) \times T_2$$

$$= \frac{1.2 \text{ atm}}{291 \text{ K}} \times 358 \text{ K}$$

$$P_2 = 1.48 \text{ atm}$$

51. $T_1 = 6^\circ \text{C} + 273 = 279 \text{ K}$

$$P_1 = 4 \text{ atm} \quad V_1 = 1.5 \text{ ml}$$

$$T_2 = 25^\circ \text{C} + 273 = 298 \text{ K}$$

$$P_2 = 1 \text{ atm} \quad V_2 = ?$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$V_2 = \frac{P_1 V_1}{T_1} \times \frac{T_2}{P_2}$$

$$= \frac{4 \text{ atm} \times 1.5 \text{ ml} \times 298 \text{ K}}{279 \text{ K} \times 1 \text{ atm}}$$

$$V_2 = 6.41 \text{ ml}$$

52. Given,

$$V = 154.4 \times 10^{-3} \text{ dm}^3,$$

$$P = 742 \text{ mm of Hg}$$

$$T = 298 \text{ K} \quad m = ?$$

$$\begin{aligned} n &= \frac{PV}{RT} = \frac{742 \text{ mm Hg} \times 154.4 \times 10^{-3} \text{ L}}{62 \text{ mm Hg L K}^{-1} \text{ mol}^{-1} \times 298 \text{ K}} \\ &= 0.006 \text{ mol} \end{aligned}$$

$$n = \frac{\text{Mass}}{\text{Molar mass}}$$

$$\text{Mass} = n \times \text{Molar mass}$$

$$= 0.006 \times 2.016$$

$$= 0.0121 \text{ g} = 12.1 \text{ mg}$$

53.

$$\frac{\gamma_{\text{unknown}}}{\gamma_{\text{N}_2}} = \frac{t_{\text{N}_2}}{t_{\text{unknown}}} = \sqrt{\frac{M_{\text{N}_2}}{M_{\text{unknown}}}}$$

$$\frac{84 \text{ sec}}{192 \text{ sec}} = \sqrt{\frac{28 \text{ g mol}^{-1}}{M_{\text{unknown}}}}$$

On squaring both sides and rearranging

$$\left(\frac{84 \text{ sec}}{192 \text{ sec}} \right)^2 = \frac{28 \text{ g mol}^{-1}}{M_{\text{unknown}}}$$

$$M_{\text{unknown}} = 28 \text{ g mol}^{-1} \times \left(\frac{192 \text{ sec}}{84 \text{ sec}} \right)^2$$

$$M_{\text{unknown}} = 146 \text{ g mol}^{-1}$$

$$\begin{aligned}
 54. \quad m_{\text{O}_2} &= 52.5 \text{ g} & P_{\text{O}_2} &= ? \\
 m_{\text{CO}_2} &= 65.1 \text{ g} & P_{\text{CO}_2} &= ? \\
 T &= 300 \text{ K} & P &= 9.21 \text{ atm}
 \end{aligned}$$

$$P_{\text{O}_2} = X_{\text{O}_2} \times \text{Total Pressure}$$

$$X_{\text{O}_2} = \frac{n_{\text{O}_2}}{n_{\text{O}_2} + n_{\text{CO}_2}}$$

$$\begin{aligned}
 n_{\text{O}_2} &= \frac{\text{Mass of O}_2}{\text{Molar mass of O}_2} \\
 &= \frac{52.5 \text{ g}}{32 \text{ g mol}^{-1}} = 1.64 \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 n_{\text{CO}_2} &= \frac{\text{Mass of CO}_2}{\text{Molar mass of CO}_2} \\
 &= \frac{65.1 \text{ g}}{44 \text{ g mol}^{-1}} = 1.48 \text{ mol}
 \end{aligned}$$

$$X_{\text{O}_2} = \frac{n_{\text{O}_2}}{n_{\text{O}_2} + n_{\text{CO}_2}} = \frac{1.64}{3.12} = 0.53$$

$$X_{\text{CO}_2} = \frac{n_{\text{CO}_2}}{n_{\text{O}_2} + n_{\text{CO}_2}} = \frac{1.48}{3.12} = 0.47$$

$$\begin{aligned}
 P_{\text{O}_2} &= X_{\text{O}_2} \times \text{Total pressure} \\
 &= 0.53 \times 9.21 \text{ atm} = 4.88 \text{ atm}
 \end{aligned}$$

$$\begin{aligned}
 P_{\text{CO}_2} &= X_{\text{CO}_2} \times \text{Total pressure} \\
 &= 0.47 \times 9.21 \text{ atm} = 4.33 \text{ atm}
 \end{aligned}$$

55. Pressure of the gas in the tank at its melting point

$$T_1 = 298 \text{ K}; P_1 = 2.98 \text{ atm}; T_2 = 1100 \text{ K}; P_2 = ?$$

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$\begin{aligned}
 \Rightarrow P_2 &= \frac{P_1}{T_1} \times T_2 \\
 &= \frac{2.98 \text{ atm}}{298 \text{ K}} \times 1100 \text{ K} = 11 \text{ atm}
 \end{aligned}$$

At 1100 K the pressure of the gas inside the tank will become 11 atm. Given that tank can withstand a maximum pressure of 12 atm, the tank will start melting first.

Unit - 7

Thermodynamics

Evaluation Yourself

1.


Solution :

Given

$$\Delta H_f^0 \text{ CO}_2 = -393.5 \text{ kJ mol}^{-1}$$

$$\Delta H_f^0 \text{ CO} = -111.31 \text{ kJ mol}^{-1}$$

$$\Delta H_f^0 (\text{H}_2\text{O}) = -242 \text{ kJ mol}^{-1}$$



$$\Delta H_r^0 = ?$$

$$\Delta H_r^0 = \sum (\Delta H_f^0)_{\text{products}} - \sum (\Delta H_f^0)_{\text{reactants}}$$

$$\Delta H_r^0 = [\Delta H_f^0 (\text{CO}) + \Delta H_f^0 (\text{H}_2\text{O})] - [\Delta H_f^0 (\text{CO}_2) + \Delta H_f^0 (\text{H}_2)]$$

$$\Delta H_r^0 = [-111.31 + (-242)] - [-393.5 + (0)]$$

$$\Delta H_r^0 = [-353.31] + 393.5$$

$$\Delta H_r^0 = 40.19$$

$$\Delta H_r^0 = +40.19 \text{ kJ mol}^{-1}$$

2.

Solution :

Given :

number of moles of water n

$$= \frac{180 \text{ g}}{18 \text{ g mol}^{-1}} = 10 \text{ mol}$$

molar heat capacity of water

$$C_p = 75.3 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$T_2 = 100^\circ \text{C} = 373 \text{ K}$$

$$T_1 = 25^\circ \text{C} = 298 \text{ K}$$

$$\Delta H = ?$$


$$\Delta H = nC_p (T_2 - T_1)$$

$$\Delta H = 10 \text{ mol} \times 75.3 \text{ J mol}^{-1} \text{ K}^{-1} \times (373 - 298) \text{ K}$$

$$\Delta H = 56475 \text{ J}$$

$$\Delta H = 56.475 \text{ kJ}$$

3.



$$\Delta U \text{ at } 25^\circ \text{C} = -3268.12 \text{ kJ}$$

Solution :

Given

$$T = 25^\circ \text{C} = 298 \text{ K} ;$$

$$\Delta U = -3268.12 \text{ kJ mol}^{-1}$$

$$\Delta H = ?$$

$$\Delta H = \Delta U + \Delta n_g RT$$

$$\Delta H = \Delta U + (n_p - n_r) RT$$

$$\Delta H = -3268.12 + \left(6 - \frac{15}{2}\right) \times 8.314 \times 10^{-3} \times 298$$

$$\Delta H = -3268.12$$

$$- (1.5 \times 8.314 \times 10^{-3} \times 298)$$


$$\Delta H = -3268.12 - 3.72$$


$$\Delta H = -3271.84 \text{ kJ mol}^{-1}$$

4.


Solution :

Given :







Sublimation :




Ionisation :





Vapourisation :




Dissociation :



Electron affinity :



Solution :



$$-524 = 148 + 2188 + 31 + 193$$

$$+ (2 \times -324.5) + u$$

$$-524 = 1911 + u$$

$$u = -524 - 1911$$

$$u = -2435 \text{ kJ mol}^{-1}$$

5.

Solution:

Given

$$T_h = 127^\circ \text{C} = 127 + 273 = 400 \text{ K}$$

$$T_c = 47^\circ \text{C} = 47 + 273 = 320 \text{ K}$$

$$\% \text{ efficiency } \eta = ?$$

$$\eta = \left[\frac{T_h - T_c}{T_h} \right] \times 100$$

$$\eta = \left[\frac{400 - 320}{400} \right] \times 100$$

$$\eta = \left[\frac{80}{400} \right] \times 100$$

$$\eta = 20\%$$

6.

Solution:


Given:

$$S^0(\text{urea}) = 173.8 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$S^0(\text{H}_2\text{O}) = 70 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$S^0(\text{CO}_2) = 213.5 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$S^0(\text{NH}_3) = 192.5 \text{ J mol}^{-1} \text{ K}^{-1}$$



$$\Delta S_r^0 = \sum (S^0)_{\text{products}} - \sum (S^0)_{\text{reactants}}$$

$$\Delta S_r^0 = [2 S^0(\text{NH}_3) + S^0(\text{CO}_2)]$$



$$- [S^0 (\text{urea}) + S^0 (\text{H}_2\text{O})]$$

$$\Delta S_r^0 = [2 \times 192.5 + 213.5]$$

$$- [173.8 + 70]$$

$$\Delta S_r^0 = [598.5] - [243.8]$$

$$\Delta S_r^0 = 354.7 \text{ J mol}^{-1} \text{ K}^{-1}$$

7.

Solution:

Given:

$$T_b = 351 \text{ K}$$

$$\Delta H_{\text{vap}} = 39840 \text{ J mol}^{-1}$$

$$\Delta S_v = ?$$

$$\Delta S_v = \frac{\Delta H_{\text{vap}}}{T_b}$$

$$\Delta S_v = \frac{39840}{351}$$

$$\Delta S_v = 113.5 \text{ J K}^{-1} \text{ mol}^{-1}$$

8.

Solution:

Given:

$$\Delta H = -10 \text{ kJ mol}^{-1} = -10000 \text{ J mol}^{-1}$$

$$\Delta S = -20 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$T = 300 \text{ K}$$

$$\Delta G = ?$$

$$\Delta G = \Delta H - T\Delta S$$

$$\Delta G = -10 \text{ kJ mol}^{-1} - 300 \text{ K} \times (-20 \times 10^{-3}) \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$\Delta G = (-10 + 6) \text{ kJ mol}^{-1}$$

$$\Delta G = -4 \text{ kJ mol}^{-1}$$

$$\text{At } 600 \text{ K}$$

$$\Delta G = -10 \text{ kJ mol}^{-1} - 600 \text{ K}$$

$$\times (-20 \times 10^{-3}) \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$\Delta G = (-10 + 12) \text{ kJ mol}^{-1}$$

$$\Delta G = +2 \text{ kJ mol}^{-1}$$

The value of ΔG is negative at 300K and the reaction is spontaneous, but at 600K the value ΔG becomes positive and the reaction is non spontaneous.

EVALUATION



I Choose the best answer:

- (b) ΔH
- (d) decrease in free energy
- (b) $q = 0$
- (d) $= 0$
- (a) $w = -\Delta U$
- (d) $\frac{\text{mass}}{\text{volume}}$
- (a) -900 J
- (b) negative
- (b) -67.6 kcal
- (a) graphite is more stable than diamond
- (d) -462 kJ
- (d) frictional energy
- (d) $\Delta H < \Delta U$
- (c) $+3 \text{ kJ}$
- (a) -2.48 kJ
- (b) -500 R
- (d) $\frac{b-2a}{2}$



18. (d) -635.66 kJ
19. (c) 80 kJ mol^{-1}
20. (a) $\Delta H < 0$ and $\Delta S > 0$
21. (c) adiabatic expansion
22. (d) $(-, -, +)$
23. (b) 27°C
24. (d) $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
25. (a) 300K

Keys to multiple choice questions:

7.

$$w = -P \Delta V$$

$$w = -(1 \times 10^5 \text{ Nm}^{-2})$$

$$(1 \times 10^{-2} \text{ m}^3 - 1 \times 10^{-3} \text{ m}^3)$$

$$w = -10^5 (10^{-2} - 10^{-3}) \text{ Nm}$$


$$w = -10^5 (10 - 1) 10^{-3} \text{ J}$$

$$w = -10^5 (9 \times 10^{-3}) \text{ J}$$

$$w = -9 \times 10^2 \text{ J}$$

$$w = -900 \text{ J}$$

9.




$$\Delta H_C^0(\text{CO}) = \Delta H_f(\text{CO}_2) - \Delta H_f(\text{CO}) + \Delta H_f(\text{O}_2)]$$

$$\Delta H_C^0(\text{CO}) = -94 \text{ KCal} - [-26.4 \text{ KCal} + 0]$$

$$\Delta H_C^0(\text{CO}) = -94 \text{ KCal} + 26.4 \text{ KCal}$$

$$\Delta H_C^0(\text{CO}) = -67.6 \text{ KCal}$$

11.



$$\Delta H_r^0 = [2\Delta H_f(\text{Cr}) + \Delta H_f(\text{Al}_2\text{O}_3)] - [2\Delta H_f(\text{Al}) + \Delta H_f(\text{Cr}_2\text{O}_3)]$$

$$\Delta H_r^0 = [0 + (-1596 \text{ kJ})]$$

$$- [0 + (-1134)]$$

$$\Delta H_r^0 = -1596 \text{ kJ} + 1134 \text{ kJ}$$

$$\Delta H_r^0 = -462 \text{ kJ}$$


14.

$$\Delta U = q + w$$

$$\Delta U = -1 \text{ kJ} + 4 \text{ kJ}$$

$$\Delta U = +3 \text{ kJ}$$

15.



1 mole of Iron liberates 1 mole of Hydrogen gas

$$55.85 \text{ g Iron} = 1 \text{ mole Iron}$$

$$\therefore n = 1$$

$$T = 25^\circ\text{C} = 298 \text{ K}$$

$$w = -P \Delta V$$

$$w = -P \left(\frac{nRT}{P} \right)$$

$$w = -nRT$$

$$w = -1 \times 8.314 \times 298 \text{ J}$$

$$w = -2477.57 \text{ J}$$

$$w = -2.48 \text{ kJ}$$

16.

$$T_i = 125^\circ\text{C} = 398 \text{ K}$$

$$T_f = 25^\circ\text{C} = 298 \text{ K}$$




$$\Delta H = nC_p (T_f - T_i)$$

$$\Delta H = 2 \times \frac{5}{2} R (298 - 398)$$

$$\Delta H = - 500 R$$


17.



(i) $\times 2$



Reverse of equation (ii) will be



(iii) + (iv)



(v) $\div 2$




18.

Given :


$$\Delta H_C (CH_4) = - 890 \text{ kJ mol}^{-1}$$

$$\Delta H_C (C_3H_8) = - 2220 \text{ kJ mol}^{-1}$$

Let the mixture contain x lit of CH_4 and $(15.68 - x)$ lit of propane.



x $2x$



$(15.68 - x)$ $5 (15.68 - x)$

$$\text{Volume of oxygen consumed} = 2x + 5 (15.68 - x) = 32 \text{ lit}$$



$$2x + 78.4 - 5x = 32 \text{ l}$$

$$78.4 - 3x = 32$$

$$3x = 46.4 \text{ l}$$

$$x = 15.47 \text{ l}$$

Given mixture contains 15.47 lit of methane and 0.213 lit of propane, hence

$$\Delta H_c = \left[\left(\frac{\Delta H_c (\text{CH}_4)}{22.4 \text{ lit}} \right) x \text{ lit} \right] + \left[\left(\frac{\Delta H_c (\text{C}_3\text{H}_8)}{22.4 \text{ lit}} \right) (15.68 - x) \text{ lit} \right]$$

$$\Delta H_c = \left[\left(\frac{-890 \text{ kJ mol}^{-1}}{22.4 \text{ lit}} \right) 15.47 \text{ lit} \right] + \left[\left(\frac{-2220}{22.4 \text{ lit}} \right) 0.21 \text{ lit} \right]$$

$$\Delta H_c = [-611.87 \text{ kJ mol}^{-1}] + [-20.81 \text{ kJ mol}^{-1}]$$

$$\Delta H_c = -632.68 \text{ KJ mol}^{-1}$$

19.

$$4E_{\text{C-H}} = 360 \text{ kJ mol}^{-1}$$

$$E_{\text{C-H}} = 90 \text{ kJ mol}^{-1}$$

$$E_{\text{C-C}} + 6 E_{\text{C-H}} = 620 \text{ kJ mol}^{-1}$$

$$E_{\text{C-C}} + 6 \times 90 = 620 \text{ kJ mol}^{-1}$$

$$E_{\text{C-C}} + 540 = 620 \text{ kJ mol}^{-1}$$

$$E_{\text{C-C}} = 80 \text{ kJ mol}^{-1}$$

22. During compression, energy of the system increases, in isothermal condition, to maintain temperature constant, heat is liberated from the system. Hence q is negative.

During compression entropy decreases.

During compression work is done on the system, hence w is positive

23.

$$\Delta S_v = \frac{\Delta H_v}{T_b}$$

$$T_b = \frac{\Delta H_v}{\Delta S_v} = \frac{4800 \text{ J mol}^{-1}}{16 \text{ J mol}^{-1} \text{ K}^{-1}} = 300 \text{ K} = 27^\circ \text{C}$$

24. In $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$, entropy change is positive. **in option d, A solid reactant gives a gaseous product. Hence the entropy change is expected to be maximum for this process.**

25.

$$\Delta G = \Delta H - T \Delta S$$

At 300K

$$\Delta G = 30000 \text{ J mol}^{-1} - 300 \text{ K} \\ \times 100 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta G = 0$$

above 300 K ; ΔG will be negative and reaction becomes spontaneous.

I Keys to the short answer questions:

53.

SOLUTION :

Given :

$$n = 2 \text{ moles}$$

$$V_i = 500 \text{ ml} = 0.5 \text{ lit}$$

$$V_f = 2 \text{ lit}$$

$$T = 25^\circ\text{C} = 298 \text{ K}$$

$$w = -2.303 nRT \log \left(\frac{V_f}{V_i} \right)$$

$$w = -2.303 \times 2 \times 8.314 \times 298 \\ \times \log \left(\frac{2}{0.5} \right)$$

$$w = -2.303 \times 2 \times 8.314 \\ \times 298 \times \log(4)$$

$$w = -2.303 \times 2 \times 8.314 \\ \times 298 \times 0.6021$$

$$w = -6871 \text{ J}$$

$$w = -6.871 \text{ kJ}$$

54.

SOLUTION :**Given :**

$$T_i = 298 \text{ K}$$

$$T_f = 298.45 \text{ K}$$

$$k = 2.5 \text{ kJ K}^{-1}$$

$$m = 3.5 \text{ g}$$

$$M_m = 28$$

$$\text{heat evolved} = k \Delta T$$

$$= k (T_f - T_i)$$

$$= 2.5 \text{ kJ K}^{-1} (298.45 - 298) \text{ K}$$

$$= 1.125 \text{ kJ}$$

$$\Delta H_c = \frac{1.125}{3.5} \times 28 \text{ kJ mol}^{-1}$$

$$\Delta H_c = 9 \text{ kJ mol}^{-1}$$

55.

SOLUTION :**Given :**

$$T_{\text{sys}} = 77^\circ \text{C} = (77 + 273) = 350 \text{ K}$$

$$T_{\text{surr}} = 33^\circ \text{C} = (33 + 273) = 306 \text{ K}$$

$$q = 245 \text{ J}$$

$$\Delta S_{\text{sys}} = \frac{q}{T_{\text{sys}}} = \frac{-245}{350} = -0.7 \text{ JK}^{-1}$$

$$\Delta S_{\text{surr}} = \frac{q}{T_{\text{sys}}} = \frac{+245}{306} = +0.8 \text{ JK}^{-1}$$

$$\Delta S_{\text{univ}} = \Delta S_{\text{sys}} + \Delta S_{\text{surr}}$$

$$\Delta S_{\text{univ}} = -0.7 \text{ JK}^{-1} + 0.8 \text{ JK}^{-1}$$

$$\Delta S_{\text{univ}} = 0.1 \text{ JK}^{-1}$$

56.

SOLUTION :**Given :**

$$n = 1 \text{ mole}$$

$$P = 4.1 \text{ atm}$$

$$V = 2 \text{ Lit}$$

$$T = ?$$

$$q = 3710 \text{ J}$$

$$\Delta S = \frac{q}{T}$$

$$\Delta S = \frac{q}{\left(\frac{PV}{nR}\right)}$$

$$\Delta S = \frac{nRq}{PV}$$

$$\Delta S = \frac{1 \times 0.082 \text{ lit atm K}^{-1} \times 3710 \text{ J}}{4.1 \text{ atm} \times 2 \text{ lit}}$$

$$\Delta S = \frac{1 \times 0.082 \text{ lit atm K}^{-1} \times 3710 \text{ J}}{4.1 \text{ atm} \times 2 \text{ lit}}$$

$$\Delta S = 37.10 \text{ JK}^{-1}$$

57.

SOLUTION:**Given :**

$$\Delta H_f(\text{NaCl}) = 30.4 \text{ kJ} = 30400 \text{ J mol}^{-1}$$

$$\Delta S_f(\text{NaCl}) = 28.4 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$T_f = ?$$

$$\Delta S_f = \frac{\Delta H_f}{T_f}$$

$$T_f = \frac{\Delta H_f}{\Delta S_f}$$




$$T_f = \frac{30400 \text{ J mol}^{-1}}{28.4 \text{ J K}^{-1} \text{ mol}^{-1}}$$

$$T_f = 1070.4 \text{ K}$$


58.

SOLUTION :


Given




$$\Delta H_c^\circ = -2220.2 \text{ kJ mol}^{-1} \text{ -----(1)}$$




$$\Delta H_f^\circ = -393.5 \text{ kJ mol}^{-1} \text{ -----(2)}$$




$$\Delta H_f^\circ = -285.8 \text{ kJ mol}^{-1} \text{ ----- (3)}$$




$$\Delta H_c^\circ = ?$$




$$\Delta H_f^\circ = -1180.5 \text{ kJ} \text{ -----(4)}$$



$$\Delta H_f^\circ = -1143.2 \text{ kJ} \text{ -----(5)}$$



$$\Delta H_f^\circ = -1180.5 - 1143.2 \\ - (-2220.2) \text{ kJ}$$



$$\Delta H_f^\circ = -103.5 \text{ kJ}$$

Standard heat of formation of propane is $\Delta H_f^\circ (\text{C}_3\text{H}_8) = -103.5 \text{ kJ}$

59.

S. No	Liquid	Boiling points ($^\circ\text{C}$)	ΔH (kJ mol $^{-1}$)
1.	Ethanol	78.4	+ 42.4
2.	Toluene	110.6	+ 35.2

SOLUTION :

For ethanol :

Given :

$$T_b = 78.4^\circ \text{C} = (78.4 + 273)$$

$$= 351.4 \text{ K}$$

$$\Delta H_v(\text{ethanol}) = + 42.4 \text{ kJ mol}^{-1}$$

$$\Delta S_v = \frac{\Delta H_v}{T_b}$$

$$\Delta S_v = \frac{+ 42.4 \text{ kJ mol}^{-1}}{351.4 \text{ K}}$$

$$\Delta S_v = \frac{+ 42400 \text{ J mol}^{-1}}{351.4 \text{ K}}$$

$$\Delta S_v = +120.66 \text{ J K}^{-1} \text{ mol}^{-1}$$

For Toluene :

Given :

$$T_b = 110.6^\circ \text{C} = (110.6 + 273)$$

$$= 383.6 \text{ K}$$

$$\Delta H_v(\text{toluene}) = + 35.2 \text{ kJ mol}^{-1}$$

$$\Delta S_v = \frac{\Delta H_v}{T_b}$$

$$\Delta S_v = \frac{+ 35.2 \text{ kJ mol}^{-1}}{383.6 \text{ K}}$$



$$\Delta S_v = \frac{+35200 \text{ J mol}^{-1}}{383.6 \text{ K}}$$

$$\Delta S_v = +91.76 \text{ J K}^{-1} \text{ mol}^{-1}$$

60.

Solution :

Given :

$$\Delta H = 30.56 \text{ kJ mol}^{-1}$$

$$= 30560 \text{ J mol}^{-1}$$

$$\Delta S = 6.66 \times 10^{-3} \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$T = ? \text{ at which } \Delta G = 0$$

$$\Delta G = \Delta H - T \Delta S$$

$$0 = \Delta H - T \Delta S$$

$$T = \frac{\Delta H}{\Delta S}$$

$$T = \frac{30.56 \text{ kJ mol}^{-1}}{6.66 \times 10^{-3} \text{ kJ K}^{-1} \text{ mol}^{-1}}$$

$$T = 4589 \text{ K}$$

(i) At 4589K ; $\Delta G = 0$ the reaction is in equilibrium.

(ii) at temperature below 4598 K , $\Delta H > T \Delta S$

$\Delta G = \Delta H - T \Delta S > 0$, the reaction in the forward direction, is non spontaneous. In other words the reaction occurs in the backward direction.

61.

Solution :

Given

$$T = 400 \text{ K}; \Delta H^0 = 77.2 \text{ kJ mol}^{-1} = 77200$$

$$\text{J mol}^{-1}; \Delta S^0 = 122 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta G^0 = -2.303 RT \log K_{eq}$$

$$\log K_{eq} = \frac{-\Delta G^0}{2.303 RT}$$

$$\log K_{eq} = -\frac{(\Delta H^0 - T \Delta S^0)}{2.303 RT}$$

$$\log K_{eq} = -\left(\frac{77200 - 400 \times 122}{2.303 \times 8.314 \times 400} \right)$$

$$\log K_{eq} = -\left(\frac{28400}{7659} \right)$$

$$\log K_{eq} = -3.7080$$

$$K_{eq} = \text{antilog}(-3.7080)$$

$$K_{eq} = 1.95 \times 10^{-4}$$

62.

Solution :

Given

$$T = 298 \text{ K}; \Delta U = -742.4 \text{ kJ mol}^{-1}$$

$$\Delta H = ?$$

$$\Delta H = \Delta U + \Delta n_g RT$$

$$\Delta H = \Delta U + (n_p - n_r) RT$$

$$\Delta H = -742.4 + \left(2 - \frac{3}{2} \right) \times 8.314 \times 10^{-3} \times 298$$

$$= -742.4 + (0.5 \times 8.314 \times 10^{-3} \times 298)$$

$$= -742.4 + 1.24$$

$$= -741.16 \text{ kJ mol}^{-1}$$

63.


Solution :**Given :**

$$E_{\text{C-H}} = 414 \text{ kJ mol}^{-1}$$

$$E_{\text{C-C}} = 347 \text{ kJ mol}^{-1}$$

$$E_{\text{C=C}} = 618 \text{ kJ mol}^{-1}$$

$$E_{\text{H-H}} = 435 \text{ kJ mol}^{-1}$$



$$\Delta H_r = \Sigma (\text{Bond energy})_r - \Sigma (\text{Bond energy})_p$$


$$\Delta H_r = (E_{\text{C=C}} + 4E_{\text{C-H}} + E_{\text{H-H}}) - (E_{\text{C-C}} + 6E_{\text{C-H}})$$

$$\Delta H_r = (618 + (4 \times 414) + 435) - (347 + (6 \times 414))$$

$$\Delta H_r = 2709 - 2831$$

$$\Delta H_r = -122 \text{ kJ mol}^{-1}$$

64.

Solution :

$$+ 2 \Delta H_4 + u$$

$$-795 = 121 + 2422 + 242.8$$

$$+ (2 \times -355) + u$$

$$-795 = 2785.8 - 710 + u$$

$$-795 = 2075.8 + u$$

$$u = -795 - 2075.8$$

$$u = -2870.8 \text{ kJ mol}^{-1}$$


65.

Solution :**Given :**

$$\Delta H_f(\text{Fe}_2\text{O}_3) = -741 \text{ kJ mol}^{-1}$$

$$\Delta H_f(\text{CO}) = -137 \text{ kJ mol}^{-1}$$

$$\Delta H_f(\text{CO}_2) = -394.5 \text{ kJ mol}^{-1}$$



$$\Delta H_r = ?$$

$$\Delta H_r = \Sigma (\Delta H_f)_{\text{products}}$$

$$- \Sigma (\Delta H_f)_{\text{reactants}}$$

$$\Delta H_r = [2 \Delta H_f(\text{Fe}) + 3 \Delta H_f(\text{CO}_2)]$$

$$- [\Delta H_f(\text{Fe}_2\text{O}_3) + 3 \Delta H_f(\text{CO})]$$

$$\Delta H_r = [0 + 3(-394.5)]$$

$$- [-741 + 3(-137)]$$

$$\Delta H_r = [-1183.5] - [-1152]$$

$$\Delta H_r = -1183.5 + 1152$$

$$\Delta H_r = -31.5 \text{ kJ mol}^{-1}$$

66.

Solution:**Given :**

$$T = 175^{\circ}\text{C} = 175 + 273 = 448\text{K}$$

Concentration of 1-pentyne

$$[A] = 1.3\%$$


Concentration of 2-pentyne

$$[B] = 95.2\%$$

Concentration of 1, 2-pentadiene


$$[C] = 3.5\%$$

At equilibrium



$$95.2\% \quad 1.3\% \Rightarrow$$

$$K_1 = \frac{1.3}{95.2} = 0.0136$$



$$95.2\% \quad 3.5\% \Rightarrow$$

$$K_2 = \frac{3.5}{95.2} = 0.0367$$

$$\Rightarrow \Delta G_1^{\circ} = -2.303 RT \log K_1$$

$$\Delta G_1^{\circ} = -2.303 \times 8.314 \times 448 \times \log 0.0136$$

$$\Delta G_1^{\circ} = +16010 \text{ J}$$

$$\Delta G_1^{\circ} = +16 \text{ kJ}$$

$$\Rightarrow \Delta G_2^{\circ} = -2.303 RT \log K_2$$

$$\Delta G_2^{\circ} = -2.303 \times 8.314 \times 448 \times \log 0.0367$$


$$\Delta G_2^{\circ} = +12312 \text{ J}$$

$$\Delta G_2^{\circ} = +12.312 \text{ kJ}$$

67.

Solution:**Given :**

$$T = 33\text{K}$$




	N_2O_4	\rightleftharpoons	2NO_2
Initial no of moles	100		0
Number of moles dissociated	50		-
No of moles remaining	50		100
Total no of moles	150 moles		
$P_{\text{N}_2\text{O}_4} = \frac{n_{\text{N}_2\text{O}_4}}{n_{\text{N}_2\text{O}_4} + n_{\text{NO}_2}} \cdot P$	$K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}}$		
$P_{\text{N}_2\text{O}_4} = \frac{50 \text{ mol}}{150 \text{ mol}} \cdot 1 \text{ atm}$	$K_p = \frac{(0.667)^2 \text{ atm}^2}{0.333 \text{ atm}}$		
$P_{\text{N}_2\text{O}_4} = 0.333 \text{ atm}$	$K_p = 1.336 \text{ atm}$		
$P_{\text{NO}_2} = \frac{n_{\text{NO}_2}}{n_{\text{N}_2\text{O}_4} + n_{\text{NO}_2}} \cdot P$	$\Delta G^{\circ} = -2.303 RT \log K_p$		
$P_{\text{NO}_2} = \frac{100 \text{ mol}}{150 \text{ mol}} \cdot 1 \text{ atm}$	$\Delta G^{\circ} = -2.303 \times 8.314 \times 33 \times \log 1.336$		
$P_{\text{NO}_2} = 0.667 \text{ atm}$	$\Delta G^{\circ} = -79.49 \text{ J mol}^{-1}$		

68.

Solution :**Given :**

$$\Delta H_f^{\circ} (\text{SO}_2) = -297 \text{ kJ mol}^{-1}$$

$$\Delta H_f^{\circ} (\text{SO}_3) = -396 \text{ kJ mol}^{-1}$$



$$\Delta H_r^{\circ} = (\Delta H_f^{\circ})_{\text{compound}}$$

$$- \sum (\Delta H_f^{\circ})_{\text{elements}}$$

$$\Delta H_r^{\circ} = \Delta H_f^{\circ} (\text{SO}_3) - \left(\Delta H_f^{\circ} (\text{SO}_2) + \frac{1}{2} \Delta H_f^{\circ} (\text{O}_2) \right)$$

$$\Delta H_r^{\circ} = -396 \text{ kJ mol}^{-1}$$

$$- (-297 \text{ kJ mol}^{-1} + 0)$$

$$\Delta H_r^{\circ} = -396 \text{ kJ mol}^{-1} + 297$$

$$\Delta H_r^{\circ} = -99 \text{ kJ mol}^{-1}$$

69.

Solution:

Given :

$$T = 298 \text{ K}$$

$$\Delta H = 400 \text{ KJ mol}^{-1}$$

$$\Delta S = 0.2 \text{ KJ K}^{-1} \text{ mol}^{-1}$$

$$\Delta G = \Delta H - T\Delta S$$

$$\text{if } T = 2000 \text{ K}$$

$$\Delta G = 400 - (0.2 \times 2000) = 0$$

$$\Delta H = 400 \text{ KJ mol}^{-1} \text{ if } T > 2000 \text{ K}$$

ΔG will be negative.

The reaction would be spontaneous only beyond 2000K

70.

Solution :

Given :

$$T = 298\text{K}$$

$$\Delta G_r^0 = -13.6 \text{ kJ mol}^{-1}$$

$$= -13600 \text{ J mol}^{-1}$$

$$\Delta G^0 = -2.303 RT \log K_{eq}$$

$$\log K_{eq} = \frac{-\Delta G^0}{2.303 RT} \quad \log K_{eq} = \frac{13.6 \text{ kJ mol}^{-1}}{2.303 \times 8.314 \times 10^{-3} \text{ J K}^{-1} \text{ mol}^{-1} \times 298 \text{ K}}$$

$$\log K_{eq} = 2.38$$

$$K_{eq} = \text{antilog}(2.38)$$

$$K_{eq} = 239.88$$

71.


Solution :

Given :


$$\Delta H_C(\text{CH}_4) = -890 \text{ kJ mol}^{-1}$$

$$\Delta H_C(\text{C}_2\text{H}_4) = -1423 \text{ kJ mol}^{-1}$$

Let the mixture contain x lit of CH_4 and $(3.67 - x)$ lit of ethylene.



x lit x lit



$(3.67 - x)$ lit 2 $(3.67 - x)$ lit

Volume of Carbondioxide formed = $x + 2(3.67 - x) = 6.11$ lit

$$x + 7.34 - 2x = 6.11$$

$$7.34 - x = 6.11$$

$$x = 1.23 \text{ lit}$$

Given mixture contains 1.23 lit of methane and 2.44 lit of ethylene, hence

$$\begin{aligned} \Delta H_C &= \left[\frac{\Delta H_C(\text{CH}_4)}{22.4 \text{ lit}} \times (x) \text{ lit} \right] \\ &+ \left[\frac{\Delta H_C(\text{C}_2\text{H}_4)}{22.4 \text{ lit}} \times (3.67 - x) \text{ lit} \right] \\ \Delta H_C &= \left[\frac{-890 \text{ kJ mol}^{-1}}{22.4 \text{ lit}} \times 1.23 \text{ lit} \right] \\ &+ \left[\frac{-1423}{22.4 \text{ lit}} \times (3.67 - 1.23) \text{ lit} \right] \end{aligned}$$

$$\Delta H_C = [-48.87 \text{ kJ mol}^{-1}]$$

$$+ [-155 \text{ kJ mol}^{-1}]$$


$$\Delta H_C = -203.87 \text{ kJ mol}^{-1}$$

Books for Reference

1. Basic concept of chemistry, L. J. Malone, T. O. Dolter, 8th Edition.
2. Chemistry in your life, Colin Baird, 2nd Edition.
3. Chemistry structure and properties, N. J. Tro, 2nd Edition.
4. General chemistry, R. Chang, 5th Edition.
5. Introductory chemistry for today, S. L. Seagal, M. R. Slabaugh, 8th Edition.
6. Basic Inorganic Chemistry, F. A. cotton, G. Wilkinson and P. L. Gaus, 3rd Edition.
7. Inorganic chemistry principles structure and reactivity, O. K. Medhi, E. A. Keiter, J. E. Huheey, R. L. Keiter, 4th Edition.
8. Inorganic chemistry, A. K. De.
9. Inorganic chemistry, Holleman-wiberg, 1st Edition.
10. Elements of physical chemistry, P. Atkins, 7th Edition.
11. Physical chemistry, I. Levine, 6th Edition..
12. Physical chemistry, G. Mortimer, 3rd Edition.
13. Advanced organic chemistry, Reactions, Mechanism and structure, J. March, 4th Edition.
14. Advanced organic chemistry, A. Bahl.
15. Introduction to organic chemistry, J. McMurry, 7th Edition.
16. Organic Chemistry, I.L Finar, 5th Edition.
17. Organic chemistry, R. T. Morrison, R. N. Boyd, 7th Edition.
18. Organic chemistry, L. G. Wade, 8th Edition.
19. Organic chemistry, J. McMurry, 9th Edition.
20. Organic chemistry, F. A.Carey, 10th Edition.
21. Environmental chemistry, R. P. Schwarzenbach, P. M. Gschwend, D. M. Imboden, 3rd Edition.

GLOSSARY

A-Z



Absolute zero	தனிச்சூழி
Absorption spectra	உட்கவர் நிறமாலை
Adiabatic process	வெப்பம் மாறாச் செயல்முறை
Alkali metals	கார உலோகங்கள்
Alkaline earth metals	காரமண் உலோகங்கள்
Allotropes	புறவேற்றுமை வடிவங்கள்
Amphoteric	ஈரியல்புள்ள
amu	அணு நிறை அலகு
Anion	எதிர்மின் அயனி
Anode ray	நேர்மின்வாய்க் கதிர்
Anomalous	அசாதாரண
Aqueous solution	நீர்க் கரைசல்
Armor plates	கவச தட்டு
Artificial silk	செயற்கைப் பட்டு
Baking soda	சமையல் சோடா
Boundary	எல்லை
Bunsen flame	புன்சன் சுடர்
Catalyst	வினையூக்கி
Cathode ray	எதிர்மின்வாய்க் கதிர்
Cation	நேர் மின் அயனி
Caustic soda	எரிசோடா
Combustion	எரிதல்
Compression	சுருங்கச் செய்தல்
Concept	கொள்கை
Covalent radius	சகப்பிணைப்பு ஆரம்
Critical constants	நிலைமாறு மாறிலிகள்
Decomposition	சிதைதல்
Diagonal relation	மூலைவிட்டத் தொடர்பு
Diffusion	விரவுதல்
Dipole moment	இருமுனைத் திருப்புத்திறன்

Direct proportion	நேர்விகிதத் தொடர்பு
Discharge tube	மின்னிறக்க குழாய்
Disproportionation	விகிதச் சிதைவு
Dual behavior	ஈரியல்புத் தன்மை
Efficiency	திிறன்
Electron affinity	எலக்ட்ரான் நாட்டம்
Electronegativity	எலக்ட்ரான் கவர்திறன்
Electrostatic attraction	நிலைமின்னியல் கவர்ச்சி
Emission spectra	உமிழ் நிறமாலை
Empirical formula	எளிய விகித வாய்ப்பாடு
Equivalent mass	சமான நிறை
Excited state	கிளர்வு நிலை
Expansion	விரிவடைதல்
Fertilizers	உரங்கள்
Formation	உருவாக்கம்
Gaseous state	வாயு நிலை
Ground state	அடிநிலை (சாதாரண நிலை)
Hard water	கடின நீர்
Hardness	கடினத்தன்மை
Heat capacity	வெப்ப ஏற்புத்திறன்
Heavy water	கன நீர்
Heterogeneous mixture	பலபடித்தான கலவை
Homogeneous mixture	ஒருபடித்தான கலவை
Hydrogen bonding	ஹைட்ரஜன் பிணைப்பு
Ideal gases	நல்லியல்பு வாயுக்கள்
Inter molecular	மூலக்கூறுகளுக்கிடையிட்ட
Inner transition metals	உள் இடைநிலைத் தனிமங்கள்
Insoluble	கரையாத்தன்மை
Internal energy	அக ஆற்றல்
Ion exchange	அயனிப் பரிமாற்றம்
Ionic radius	அயனி ஆரம்
Ionisation energy	அயனியாக்கும் ஆற்றல்
Irreversible	மீளாச் செயல்முறை
Isobaric process	அழுத்தம் மாறாச் செயல்முறை
Isochoric process	கனஅளவு மாறாச் செயல்முறை
Isoelectronic species	ஒத்த எலக்ட்ரான் அமைப்பு கொண்ட கூறுகள்
IUPAC	International Union of Pure and Applied Chemistry
Lather	நுரை

Lattice energy	படிகக் கூடு ஆற்றல்
Laundering	சலவை செய்தல்
Lime stone	சுண்ணாம்புக் கல் $CaCO_3$
Linear	நேர்கோட்டு வடிவம்
Lone pair	தனித்த இரட்டை
Matter	பொருண்மை
Milk of lime	சுண்ணப்பால்- $Ca(OH)_2$
Milk of magnesia	மெக்னீஷியா பால்மம் $Mg(OH)_2$
Naphthalene balls	நாஃப்தலீன் உருண்டை (இரசக் கற்பூரம்)
Negligible	புறக்கணிக்கத்தக்க
Neutralisation	நடுநிலையாக்கல்
nm	நேநோ மீட்டர்
Non-spontaneous process	தன்னிச்சையற்ற செயல்முறை
Octahedron	எண்முகி
Octaves	எட்டு எண்ணிக்கை கொண்ட தொகுதி (a group of eight)
Octet rule	எண்ம விதி
Operator	செயலி
Oxidation	ஆக்ஸிஜனேற்றம்
Oxidation number	ஆக்ஸிஜனேற்ற எண்
Path function	வழிச் சார்பு
Periodic table	ஆவர்த்தன அட்டவணை
Photoelectric effect	ஒளிமின் விளைவு
Plaster of Paris	பாரீஸ் சாந்து
pm	பிகோ மீட்டர்
Polarisation	முனைவாக்கல்
Polarisability	முனைவாக்கும் திறன்
Polyatomic molecule	பல்லணு மூலக்கூறு
Precipitate	வீழ்ப்படிவு
Quantity	பொருளின் அளவு
Quick lime	சுட்ட சுண்ணாம்பு- CaO
Radioactive elements	கதிரியக்கத் தனிமங்கள்
Reagent	வினைக் கரணி
Redox	ஆக்ஸிஜனேற்ற-ஒடுக்கம்
Reduction	ஒடுக்கம்
Relative atomic mass	ஒப்பு அணு நிறை
Reversible process	மீள் செயல்முறை
Saturated solution	நிறைவுற்ற கரைசல்
Scattering	சிதறடித்தல்



Shielding effect	மறைத்தல் விளைவு
SI unit	System International அலகு
Slaked lime - Ca(OH)_2	நீர்த்த சுண்ணாம்பு
Soluble	கரையக்கூடிய
Sparingly soluble	பகுதியாக கரையும் தன்மை
Spin	தற்சுழற்சி
Spontaneous process	தன்னிச்சைச் செயல்முறை
State function	நிலைச்சார்பு
Sublimation	பதங்கமாதல்
Surroundings	சுற்றுப்புறம்
Synthesis	தொகுத்தல்
Tetrahedron	நான்முகி
Thermal equilibrium	வெப்பச் சமநிலை
Trans uranium elements	யுரேனியத்தை தொடர்ந்து வரும் தனிமங்கள்
Transition metals	இடைநிலைத் தனிமங்கள்
Triads	மும்மைகள்
Uncertainty	நிச்சயமற்ற தன்மை
Unified mass	வரையறுக்கப்பட்ட நிறை
Universe	அண்டம்
Unsaturated compound	நிறைவுறாச் சேர்மம்
Valence electrons	இணைதிறன் எலக்ட்ரான்கள்
Valency	இணைதிறன்
Vapour pressure	ஆவி அழுத்தம்
Vapourisation	ஆவியாதல்
Viscosity	பாகுநிலைத்தன்மை
Wave function	அலைச் சார்பு





Chemistry – Class XI

List of Authors and Reviewers

Domain Experts / Reviewers

Mentor

Dr. E. Murugan

Professor & Head
Department of Physical Chemistry
School of Chemical Sciences
University of Madras,
Guindy Campus, Guindy, Chennai.

Dr. M. Palanichamy

Professor (Retd.)
Department of Chemistry
Anna University, Chennai.

Dr. V. Subramaniam

Professor (Retd.)
Department of Chemistry
Pachaiyappa's College, Chennai.

Dr. P. Selvam

Professor
Department of Chemistry &
National Centre for Catalysis Research
Indian Institute of Technology-Madras,
Chennai.

Dr. Mangala Sunder Krishnan

Professor
Department of Chemistry
Indian Institute of Technology-Madras,
Chennai.

Prof. B. Viswanathan

Professor (Retd.)
Department of Chemistry
Indian Institute of Technology-Madras,
Chennai.

Prof. V.R. Vijayaraghavan

Professor & Head (Retd.)
Department of Physical Chemistry
School of Chemical Sciences
University of Madras.

Dr. U. Venkatasubramanian

Senior Assistant Professor
School of Chemical and Biotechnology
Sastra Deemed to be University, Thanjavur.

Content Development Team

Dr. S. Chidambara Vinayagam

Principal
Muthukalathur Government Arts College,
Ramanathapuram.

Dr. S. Arul Antony

Head of the Department of Chemistry
Presidency College, Chennai.

S. Kannan

Assistant Professor and Head Post Graduate and
Research Department of Chemistry
LG Government College (Autonomous)
Ponneri.

Dr. A. Syed Mohamed

Assistant Professor
Research Department of Chemistry
Sadakathullah Appa College (Autonomous)
Tirunelveli.

Dr. K. Muruga Poopathi Raja

Assistant Professor
Department of Physical Chemistry
Madurai Kamaraj University, Madurai.

Dr. V.S. Gayathri

Professor and Head
Department of Chemistry
SSN Institution, Kalavakkam, Chennai.

Dr. P. Kamaraj

Professor
Department of Chemistry
Vel Tech, Avadi, Chennai.

C.E. Ruckmani Jayanthi

Post Graduate Assistant
C. Kalyanam Higher Secondary School,
Chintadrapet, Chennai.

D. Jagannathan

Post Graduate Assistant
SGR Government Higher Secondary School,
Kosavanpudur, Vellore District.

R. Subramanian

Post Graduate Assistant
Government Higher Secondary School,
Malayappalayam, Gobichettipalayam, Erode Dt.

Expert & Co-ordinator

Boopathi Rajendran

Deputy Director
Directorate of Elementary Education,
Chennai - 6.

C. Mahalakshmi

Post Graduate Assistant
Presidency Higher Secondary School, Chennai.

A. Helen Jayanthi

Post Graduate Assistant
St. Raphael's Girls Higher Secondary School,
Santhome, Chennai.

Dr. S. Sankar

Post Graduate Assistant
Government Higher Secondary School,
Thirupputkuzhi, Kancheepuram District.

S. Balagurunathan

Head Master (Retd.)
NLC Boys Higher Secondary School,
Neyveli, Cuddalore District.

L. Balthasar

Post Graduate Assistant
Government Higher Secondary School
Thottiyam, Trichy District.

K. Durai Chandran

Post Graduate Assistant
Chinnmaya Vidhyalaya Higher Secondary School,
Chennai.

S. Sasikumar

Post Graduate Assistant
Green Garden Girls Matriculation Higher
Secondary School, Perundurai, Erode.

ICT Co-ordinators

Dr. K. Rajendra Kumar

Assistant Professor (Sr)
Chemistry Division
School of Advanced Sciences, VIT - Chennai.

D. Vasuraj

B.T. Assistant
P.U.M.S Kosapur
Puzhal Block, Thiruvallur District.

S. Shyamala

B.T. Assistant
Govt. ADW High School,
Pullianthope, Chennai.

Content Readers

R.C. Saraswathi

Headmistress
Government Girls Higher Secondary School,
Ashok Nagar, Chennai.

Dr. M. Alphonse

Headmaster (Retd.)
Government Higher Secondary School,
Sadurangapatnam, Kancheepuram District.

G. Rajendran

Headmaster (Retd.)
Government Periyar Higher Secondary School,
Puzhuthivakkam, Chennai.

Co-ordinator – Logistics

P. Sankaralingam

Post Graduate Assistant (Chemistry)
Government Higher Secondary School,
Melakkottai, Madurai District.

A. Palanivel Raj

Assistant Professor
SCERT, Chennai.

This book has been printed on 80 G.S.M.
Elegant Maplitho paper.

Printed by offset at:

Art and Design Team

Layout

A Adison Raj,
Mathan Raj,
C. Prasanth,
Kamatchi Balan

In-House - QC

Rajesh Thangappan,
Jerald wilson

Co-ordinator

Ramesh Munisamy