

Chapter 9. Sequences and Series

Question-1

Write the first 5 terms of each of the following sequences:

(i) $a_n = (-1)^{n-1} 5^{n+1}$

(ii) $a_n = \frac{n(n^2+5)}{4}$

(iii) $a_n = -11n + 10$

(iv) $a_n = \frac{n+1}{n+2}$

(v) $a_n = \frac{1-(-1)^n}{3}$

(vi) $a_n = \frac{n^2}{3^n}$

Solution:

(i) $a_n = (-1)^{n-1} 5^{n+1}$

$a_1 = (-1)^0 5^2 = 5^2;$

$a_2 = (-1)^1 5^3 = -5^3$

$a_3 = (-1)^2 5^4 = 5^4;$

$a_4 = (-1)^3 5^5 = -5^5$

$a_5 = (-1)^4 5^6 = 5^6$

(ii) $a_n = \frac{n(n^2+5)}{4}$

$$a_1 = \frac{1(1^2 + 5)}{4} = \frac{6}{4} = \frac{3}{2}$$

$$a_2 = \frac{2(4 + 5)}{4} = \frac{18}{4} = \frac{9}{2}$$

$$a_3 = \frac{3(9 + 5)}{4} = \frac{21}{2}$$

$$a_4 = \frac{4(16 + 5)}{4} = 21$$

$$a_5 = \frac{5(25 + 5)}{4} = \frac{75}{2}$$

$$(iii) a_n = -11n + 10$$

$$a_1 = -11 + 10 = -1$$

$$a_2 = -22 + 10 = -12$$

$$a_3 = -33 + 10 = -23$$

$$a_4 = -44 + 10 = -34$$

$$a_5 = -55 + 10 = -45$$

$$(iv) a_n = \frac{1+1}{1+2} = \frac{2}{3}$$

$$a_2 = \frac{2+1}{2+2} = \frac{3}{4}$$

$$a_3 = \frac{3+1}{3+2} = \frac{4}{5}$$

$$a_4 = \frac{4+1}{4+2} = \frac{5}{6}$$

$$a_5 = \frac{5+1}{5+2} = \frac{6}{7}$$

$$(v) a_n = \frac{1 - (-1)^n}{3}$$

$$a_1 = \frac{1 - (-1)^1}{3} = \frac{2}{3}$$

$$a_2 = \frac{1 - (-1)^2}{3} = 0$$

$$a_3 = \frac{1 - (-1)^3}{3} = \frac{2}{3}$$

$$a_4 = \frac{1 - (-1)^4}{3} = 0$$

$$a_5 = \frac{1 - (-1)^5}{3} = \frac{2}{3}$$

$$(vi) a_n = \frac{n^2}{3^n}$$

$$a_1 = \frac{1^2}{3^1}; a_2 = \frac{2^2}{3^2}; a_3 = \frac{3^2}{3^3}; a_4 = \frac{4^2}{3^4}; a_5 = \frac{5^2}{3^5}$$

Question-2

Find the first terms of the following sequences whose n^{th} term is

(i) $a_n = 2 + \frac{1}{n}$; a_5, a_7

(ii) $a_n = \cos\left[\frac{n\pi}{2}\right]$; a_4, a_5

(iii) $a_n = \frac{n+1}{n}$; a_7, a_{10}

(iv) $a_n = (-1)^{n-1} 2^{n+1}$; a_5, a_8

Solution:

(i) $a_n = 2 + \frac{1}{n}$

$$a_5 = 2 + \frac{1}{5}$$

$$= \frac{11}{5};$$

$$a_7 = 2 + \frac{1}{7}$$

$$= \frac{15}{7}$$

(ii) $a_n = \cos\left[\frac{n\pi}{2}\right];$

$$a_4 = \cos\left(\frac{4\pi}{2}\right) = \cos 2\pi = 1$$

$$a_5 = \cos\left[\frac{5\pi}{2}\right] = \cos\left[2\pi + \frac{\pi}{2}\right] = \cos \frac{\pi}{2} = 0$$

(iii) $a_n = \frac{(n+1)^2}{n}$

$$a_7 = \frac{(7+1)^2}{7}$$

$$= \frac{64}{7};$$

$$a_{10} = \frac{(10+1)^2}{10}$$

$$= \frac{121}{10}$$

(iv) $a_n = (-1)^{n-1} 2^{n+1}$

$$a_5 = (-1)^4 2^{5+1}$$

$$= 2^6$$

$$= 64;$$

$$a_8 = (-1)^7 2^{8+1}$$

$$= -2^9$$

$$= -512$$

Question-3

Find the first 6 terms of the sequence whose general term is

$$a_n = \{n^2 - 1 \text{ if } n \text{ is odd } \frac{n^2 + 1}{2} \text{ if } n \text{ is even}\}$$

Solution:

$$a_1 = 1^2 - 1 = 0$$

$$a_2 = \frac{2^2 + 1}{2} = \frac{5}{2}$$

$$a_3 = 3^2 - 1 = 8$$

$$a_4 = \frac{4^2 + 1}{2} = \frac{17}{2}$$

$$a_5 = 5^2 - 1 = 24$$

$$a_6 = \frac{6^2 + 1}{2} = \frac{37}{2}$$

Question-4

Write the first five terms of the sequence given by

(i) $a_1 = a_2 = 2, a_n = a_{n-1} - 1, n > 2$

(ii) $a_1 = 1, a_2 = 2, a_n = a_{n-1} + a_{n-2}, n > 2$

(iii) $a_1 = 1, a_n = na_{n-1}, n \geq 2$

(iv) $a_1 = a_2 = 1, a_n = 2a_{n-1} + 3a_{n-2}, n > 2$

Solution:

(i) Put $n = 3 \Rightarrow a_3 = a_2 - 1 = 2 - 1 = 1$

$$n = 4 \Rightarrow a_4 = a_3 - 1 = 1 - 1 = 0$$

$$n = 5 \Rightarrow a_5 = a_4 - 1 = 0 - 1 = -1$$

(ii) Put $n = 3 \Rightarrow a_3 = a_2 + a_1 = 2 + 1 = 3$

$$n = 4 \Rightarrow a_4 = a_3 + a_2 = 3 + 2 = 5$$

$$n = 5 \Rightarrow a_5 = a_4 + a_3 = 5 + 3 = 8$$

(iii) Put $n = 2 \Rightarrow a_2 = 2 a_1 = 2.1 = 2$

$$n = 3 \Rightarrow a_3 = 3 a_2 = 3.2 = 6$$

$$n = 4 \Rightarrow a_4 = 4 a_3 = 4.6 = 24$$

$$n = 5 \Rightarrow a_5 = 5 a_4 = 5.24 = 120$$

(iv) Put $n = 3 \Rightarrow a_3 = 2a_2 + 3a_1 = 2(1) + 3(1) = 5$

$$n = 4 \Rightarrow a_4 = 2a_3 + 3a_2 = 2(5) + 3(1) = 13$$

$$n = 5 \Rightarrow a_5 = 2a_4 + 3a_3 = 2(13) + 3(5) = 41$$

Question-5

Find the n^{th} partial sum of the series $\sum_{n=1}^{\infty} \frac{1}{3^n}$

Solution:

$$\sum_{n=1}^{\infty} \frac{1}{3^n}$$

$$S_n = \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots + \frac{1}{3^n}$$

$$S_{n+1} = \frac{1}{3} + \frac{1}{3^2} + \dots + \frac{1}{3^n} + \frac{1}{3^{n+1}}$$

$$S_{n+1} = S_n + \frac{1}{3^{n+1}}$$

$$\begin{aligned} S_{n+1} &= \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots + \frac{1}{3^n} + \frac{1}{3^{n+1}} \\ &= \frac{1}{3} \left[1 + \frac{1}{3} + \frac{1}{3^2} + \dots + \frac{1}{3^n} \right] = \frac{1}{3} [1 + S_n] \end{aligned}$$

$$S_n + \frac{1}{3^{n+1}} = \frac{1}{3} + \frac{1}{3} S_n$$

$$3 S_n + \frac{1}{3^n} = 1 + S_n$$

$$2S_n = 1 - \frac{1}{3^n}$$

$$S_n = \frac{1}{2} \left[1 - \frac{1}{3^n} \right]$$

Question-6

Find the sum of first n terms of the series $\sum_{n=1}^{\infty} 5^n$

Solution:

$$\sum_{n=1}^{\infty} 5^n = 5 + 5^2 + 5^3 + \dots + 5^n + \dots$$

$$S_n = 5 + 5^2 + 5^3 + \dots + 5^n$$

$$\begin{aligned} S_{n+1} &= 5 + 5^2 + 5^3 + \dots + 5^n + 5^{n+1} \\ &= S_n + 5^{n+1} \end{aligned}$$

$$\begin{aligned} \text{Also } S_{n+1} &= 5 + 5^2 + 5^3 + \dots + 5^n + 5^{n+1} \\ &= 5[1 + 5 + 5^2 + \dots + 5^n] \\ &= 5[1 + S_n] \end{aligned}$$

$$S_n + 5^{n+1} = 5 + 5 S_n$$

$$4S_n = 5^{n+1} - 5$$

$$\therefore S_n = \frac{5(5^n - 1)}{4}$$

Question-7

Find the sum of 101th term to 200th term of the series $\sum_{n=1}^{\infty} \frac{1}{2^n}$

Solution:

$$\sum_{n=1}^{\infty} \frac{1}{2^n}$$

To find $S_{200} - S_{100}$

$$\text{To find } S_{200}: S_{200} = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{200}}$$

$$\begin{aligned} S_{201} &= \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{200}} + \frac{1}{2^{201}} \\ &= S_{200} + \frac{1}{2^{201}} \end{aligned}$$

$$\begin{aligned} \text{also, } S_{201} &= \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{200}} + \frac{1}{2^{201}} \\ &= \frac{1}{2} \left[1 + \frac{1}{2} + \frac{1}{2^2} + \dots + \frac{1}{2^{200}} \right] \end{aligned}$$

$$S_{200} + \frac{1}{2^{201}} = \frac{1}{2} [1 + s_{200}]$$

$$2S_{200} + \frac{1}{2^{200}} = 1 + S_{200}$$

$$S_{200} = 1 - \frac{1}{2^{200}}$$

$$\text{Similarly } S_{100} = 1 - \frac{1}{2^{100}}$$

$$\begin{aligned} \text{Hence } S_{200} - S_{100} &= \left[1 - \frac{1}{2^{200}} \right] - \left[1 - \frac{1}{2^{100}} \right] \\ &= \frac{1}{2^{100}} - \frac{1}{2^{200}} \end{aligned}$$

Question-8

Find five arithmetic means between 1 and 19.

Solution:

Let 1, x_1 , x_2 , x_3 , x_4 , x_5 , 19 be in A.P.

Let d be the common difference

$$19 = 1 + (n-1)d$$

$$19 = 1 + 6d$$

$$\therefore d = 3$$

$$\therefore x_1 = 1 + 3 = 4$$

$$x_2 = 4 + 3 = 7$$

$$x_3 = 7 + 3 = 10$$

$$x_4 = 10 + 3 = 13$$

$$x_5 = 13 + 3 = 16$$

The arithmetic means are 4, 7, 10, 13, 16.

Question-9

Find six arithmetic mean between 3 and 17.

Solution:

Let 3, x_1 , x_2 , x_3 , x_4 , x_5 , x_6 , 17 be in A.P

Then $17 = 3 + (n-1)d$

$$17 = 3 + 7d$$

$$14 = 7d$$

$$d = 2$$

$$x_1 = 3 + 2 = 5$$

$$x_2 = 5 + 2 = 7$$

$$x_3 = 7 + 2 = 9$$

$$x_4 = 9 + 2 = 11$$

$$x_5 = 11 + 2 = 13$$

$$x_6 = 13 + 2 = 15$$

The arithmetic means are 5, 7, 9, 11, 13, 15.

Question-10

Find the single A.M. between

(i) 7 and 13

(ii) 5 and -3

(iii) $(p + q)$ and $(p - q)$

Solution:

(i) A.M. between 7 and 13 = $\frac{7+13}{2} = 10$

(ii) A.M. between 5 and -3 = $\frac{5-3}{2} = 1$

(iii) A.M. between $(p + q)$ and $(p - q)$ = $\frac{p+q+p-q}{2} = p$

Question-11

If b is the G.M. of a and c and x is the A.M of a and b and y is the A.M of b and c , prove that $\frac{a}{x} + \frac{c}{y} = 2$.

Solution:

$$b = \text{G.M. of } a \text{ and } c \Rightarrow \sqrt{ac} = b \dots\dots\dots(1)$$

$$x = \text{A.M. of } a \text{ and } b \Rightarrow x = \frac{a+b}{2} \dots\dots\dots(2)$$

$$y = \text{A.M. between } b \text{ and } c \Rightarrow y = \frac{b+c}{2} \dots\dots\dots(3)$$

To prove that $\frac{a}{x} + \frac{b}{y} = 2$

$$\text{From (1) } b^2 = ac \Rightarrow c = \frac{b^2}{a}$$

$$\begin{aligned} \frac{a}{x} + \frac{c}{y} &= \frac{2a}{a+b} + \frac{2c}{b+c} \\ &= \frac{2a}{a+b} + \frac{\frac{2b^2}{a}}{b + \frac{b^2}{a}} \\ &= \frac{2a}{a+b} + \frac{2bb}{b(a+b)} \\ &= \frac{2a}{a+b} + \frac{2b}{a+b} \\ &= \frac{2(a+b)}{a+b} \\ &= 2 \end{aligned}$$

Question-12

The first and second terms of H.P are $\frac{1}{3}$ and $\frac{1}{5}$ respectively, find the 9th term.

Solution:

Let the H.P are $\frac{1}{a}, \frac{1}{a+d}, \frac{1}{a+2d} + \dots\dots\dots$

$$\frac{1}{a} = \frac{1}{3} \Rightarrow a = 3$$

$$\frac{1}{a+d} = \frac{1}{5} \Rightarrow 5 = a + d; d = 2$$

$$9^{\text{th}} \text{ term} = \frac{1}{a+8d} = \frac{1}{3+16} = \frac{1}{19}$$

Question-13

If a, b, c are in H.P., prove that $\frac{b+a}{b-a} + \frac{b+c}{b-c} = 2$.

Solution:

If a, b, c are in H.P then $b = \frac{2ac}{a+c}$

$$\frac{b}{a} = \frac{2c}{a+c}$$
$$\Rightarrow \frac{b+a}{b-a} = \frac{2c+a+c}{2c-a-c} = \frac{3c+a}{c-a} \dots\dots\dots(1)$$

$$\text{Also } \frac{b}{c} = \frac{2a}{a+c}$$

$$\frac{b+c}{b-c} = \frac{2a+a+c}{2a-a-c} = \frac{3a+c}{a-c} \dots\dots\dots(2)$$

Adding (1) and (2)

$$\begin{aligned} \therefore \frac{b+a}{b-a} + \frac{b+c}{b-c} &= \frac{3c+a}{c-a} + \frac{3a+c}{a-c} \\ &= \frac{3c+a}{c-a} - \frac{3a+c}{c-a} \\ &= \frac{3c+a-3a-c}{c-a} \\ &= \frac{2c-2a}{c-a} \\ &= 2 \end{aligned}$$

Hence $\frac{b+a}{b-a} + \frac{b+c}{b-c} = 2$.

Question-14

The difference between two positive numbers is 18, and 4 times their G.M. is equal to 5 times their H.M. find the numbers.

Solution:

Let the two numbers be a and b

$$b - a = 18$$

$$4\sqrt{ab} = 5 \left[\frac{2ab}{a+b} \right]$$

$$2\sqrt{ab} = \frac{5ab}{a+b}$$

$$2(a+b) = 5\sqrt{ab}$$

$$4(a+b)^2 = 25ab$$

$$4(a^2 + b^2 + 2ab) = 25ab$$

$$4a^2 + 4b^2 - 17ab = 0$$

$$4a^2 + 4(18+a)^2 - 17a(18+a) = 0$$

$$4a^2 + 4(324 + 36a + a^2) - 306a - 17a^2 = 0$$

$$4a^2 + 1296 + 144a + 4a^2 - 306a - 17a^2 = 0$$

$$-9a^2 - 162a + 1296 = 0$$

$$a^2 + 18a - 144 = 0$$

$$(a+24)(a-6) = 0$$

$$a = -24 \text{ (or) } 6$$

If $a = 6$ then b is 24.

Therefore the numbers are 6 and 24.

Question-15

If the A.M. between two numbers is 1, prove that their H.M. is the square of their G.M.

Solution:

A.M. between two numbers a and b is 1.

$$\frac{a+b}{2} = 1$$

$$\Rightarrow a + b = 2$$

$$HM = (GM)^2$$

$$HM = \frac{2ab}{a+b} = \frac{2ab}{2} = ab$$

$$GM = \sqrt{ab}$$

$$\therefore (GM)^2 = ab$$

$$\text{Hence } HM = GM^2$$

Question-16

If a, b, c are in A.P and a, mb, c are in G.P then prove that a, m^2b, c are in H.P.

Solution:

Given

a, b, c are in A.P.

$$\Rightarrow b = \frac{a+c}{2} \dots\dots\dots(1)$$

a, mb, c are in G.P.

$$\Rightarrow mb = \sqrt{ac} \dots\dots\dots(2)$$

To prove

a, m^2b, c are in H.P.

$$\text{i.e., } m^2b = \frac{2ac}{a+c}$$

Proof

$$\begin{aligned} \text{R.H.S} &= \frac{2ac}{a+c} = \frac{2m^2b^2}{2b} \text{ from (2) and (1)} \\ &= m^2 b = \text{LHS} \end{aligned}$$

Question-17

If the p^{th} and q^{th} terms of a H.P. are q and p respectively, show that $(pq)^{\text{th}}$ term is 1.

Solution:

Given

p^{th} and q^{th} terms of a H.P. are q and p .

$$\text{Therefore } \frac{1}{a+(p-1)d} = q \dots\dots\dots(1)$$

$$\text{and } \frac{1}{a+q-1d} = p \dots\dots\dots(2)$$

To prove

$$pq^{\text{th}} \text{ term, i.e., } \frac{1}{a+(pq-1)d} = 1$$

Proof

$$\text{From (1) } a + pd - d = \frac{1}{q}$$

$$\text{From (2) } a + qd - d = \frac{1}{p}$$

$$\text{Subtracting } (p - q)d = \frac{1}{q} - \frac{1}{p} = \frac{p - q}{pq}$$

$$\therefore d = \frac{1}{pq}$$

$$\therefore a + p \frac{1}{pq} - \frac{1}{pq} = \frac{1}{q}$$

$$a = \frac{1}{pq}$$

$$\therefore a + (pq - 1)d = \frac{1}{pq} + (pq - 1) \frac{1}{pq}$$

$$= \frac{1 + pq - 1}{pq}$$

$$= \frac{pq}{pq} = 1$$

$$\therefore \frac{1}{a + (pq - 1)d} = 1$$

$\Rightarrow pq^{\text{th}}$ term is 1.

Question-18

Three number form a H.P. the sum of the numbers is 11 and the sum of the reciprocals is one. Find the numbers.

Solution:

Let $\frac{1}{a-d}, \frac{1}{a}, \frac{1}{a+d}$ be in H.P.

Their sum is $\frac{1}{a-d} + \frac{1}{a} + \frac{1}{a+d} = 11$

The sum of their reciprocal is $a - d + a + a + d = 1$

$$3a = 1$$

$$\Rightarrow a = \frac{1}{3}$$

$$\therefore \frac{1}{\frac{1}{3}-d} + \frac{1}{\frac{1}{3}} + \frac{1}{\frac{1}{3}+d} = 11$$

$$\frac{3}{1-3d} + 3 + \frac{3}{1+3d} = 11$$

$$\frac{3}{1-3d} + \frac{3}{1+3d} = 8$$

$$\frac{3(1+3d+1-3d)}{1-9d^2} = 8$$

$$\frac{6}{1-9d^2} = 8$$

$$6 = 8 - 72d^2$$

$$72d^2 = 2$$

$$\Rightarrow d^2 = \frac{1}{36}$$

$$\Rightarrow d = \frac{1}{6}$$

$$\text{The numbers are } \frac{1}{\frac{1}{3}-\frac{1}{6}} + \frac{1}{\frac{1}{3}} + \frac{1}{\frac{1}{3}+\frac{1}{6}} = \frac{1}{\frac{1}{6}}, \frac{1}{\frac{1}{3}}, \frac{1}{\frac{1}{2}}$$

The numbers are 6, 3, 2.

Question-19

Write the first four terms in the expansions of the following:

(i) $\frac{1}{(2+x)^4}$ where $|x| > 2$

(ii) $\frac{1}{\sqrt[3]{6-3x}}$ where $|x| < 2$

Solution:

$$(i) \frac{1}{(2+x)^4} = \frac{1}{16\left(1+\frac{x}{2}\right)^4} = \frac{1}{16} \left[1+\frac{x}{2}\right]^{-4}$$

$$= \frac{1}{16} \left[1 - 4\left(\frac{x}{2}\right) + \frac{4 \cdot 5}{1 \cdot 2} \left(\frac{x}{2}\right)^2 - \frac{4 \cdot 5 \cdot 6}{1 \cdot 2 \cdot 3} \left(\frac{x}{2}\right)^3 + \dots\right]$$

$$= \frac{1}{16} \left[1 - 2x + \frac{5}{2}x^2 - \frac{5}{2}x^3 + \dots\right]$$

$$(ii) \frac{1}{\sqrt[3]{6-3x}} = \frac{1}{(6-3x)^{\frac{1}{3}}} = \frac{1}{6^{\frac{1}{3}}} \left[1 - \frac{x}{2}\right]^{\frac{1}{3}}$$

$$= \frac{1}{6^{\frac{1}{3}}} \left[1 + \frac{1}{3}\left(\frac{x}{2}\right) + \frac{\left(\frac{1}{3}\right)\left(\frac{4}{3}\right)}{1 \cdot 2} \left(\frac{x}{2}\right)^2 + \frac{\left[\frac{1}{3}\right]\left[\frac{4}{3}\right]\left[\frac{7}{3}\right]}{1 \cdot 2 \cdot 3} \left[\frac{x}{2}\right]^3\right]$$

$$= \frac{1}{6^{\frac{1}{3}}} \left[1 + \frac{x}{6} + \frac{x^2}{18} + \frac{7}{326}x^3 + \dots\right]$$

Question-20

Evaluate the following:

(i) $\sqrt[3]{1003}$ correct to 4 places of decimals.

(ii) $\frac{1}{\sqrt[3]{128}}$ correct to 4 places of decimals.

(iii) $\sqrt[3]{1003} - \sqrt[3]{997}$ correct to 3 place of decimals.

Solution:

$$\begin{aligned}\text{(i)} \quad \sqrt[3]{1003} &= (1003)^{\frac{1}{3}} \\&= (1000 + 3)^{\frac{1}{3}} \\&= (1000)^{1/3} \left[1 + \frac{3}{1000} \right]^{\frac{1}{3}} \\&= 10 [1 + 0.003]^{\frac{1}{3}} \\&= 10 \left[1 + \frac{1}{3}(0.003) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)}{1.2}(0.003)^2 + \dots \right] \\&= 10 [1 + 0.001 - 0.000001 + \dots] \\&= 10.00999\end{aligned}$$

$$\begin{aligned}\text{(ii)} \quad \frac{1}{\sqrt[3]{128}} &= \frac{1}{(128)^{\frac{1}{3}}} = \frac{1}{(125+3)^{\frac{1}{3}}} = \frac{1}{5 \left[1 + \frac{3}{125} \right]^{\frac{1}{3}}} = \frac{1}{5} \left[1 + \frac{3}{125} \right]^{-\frac{1}{3}} \\&= \frac{1}{5} (1 + 0.024)^{-\frac{1}{3}} = \frac{1}{5} \left[1 - \frac{1}{3}(0.024) + \frac{\frac{1}{3}\left(\frac{4}{3}\right)}{1.2}(0.024)^2 + \dots \right] \\&= \frac{1}{5} [1 - 0.008 + 0.000128] = 0.1984256\end{aligned}$$

$$\begin{aligned}\text{(iii)} \quad \sqrt[3]{1003} - \sqrt[3]{997} &= (1000 + 3)^{1/3} - (1000 - 3)^{1/3} \\&= 10 \left[1 + \frac{3}{1000} \right]^{\frac{1}{3}} - 10 \left[1 - \frac{3}{1000} \right]^{\frac{1}{3}} \\&= 10 [1 + 0.003]^{\frac{1}{3}} - 10 [1 - 0.003]^{\frac{1}{3}} \\&= 10 \left[1 + \frac{1}{3}(0.003) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)}{1.2}(0.003)^2 + \dots \right] \\&\quad - 10 \left[1 - \frac{1}{3}(0.003) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)}{1.2}(0.003)^2 + \dots \right] \\&= 10 \left[1 + \frac{0.003}{3} - \frac{(0.003)^2}{9} - 1 + \frac{0.003}{3} - \frac{(0.003)^2}{9} + \dots \right] \\&= 10[0.002] = 0.02\end{aligned}$$

Question-21

If x is so small show that

$$(i) \frac{\sqrt{1-x}}{1+x} = 1 - x + \frac{x^2}{2} \text{ (app)}$$

$$(ii) \frac{1}{(1+x)^2 \sqrt{1+4x}} = 1 - 4x \text{ (app.)}$$

Solution:

$$\begin{aligned} (i) \frac{\sqrt{1-x}}{1+x} &= (1-x)^{\frac{1}{2}} (1+x)^{-\frac{1}{2}} \\ &= \left[1 - \frac{1}{2}x + \frac{\left[\frac{1}{2}\right]\left[\frac{-1}{2}\right]}{1.2}x^2 + \dots \right] \left[1 - \frac{1}{2}x + \frac{\frac{1}{2}\frac{3}{2}}{1.2}x^2 + \dots \right] \\ &= \left[1 - \frac{x}{2} - \frac{x^2}{8} \right] \left[1 - \frac{x}{2} + \frac{3x^2}{8} \right] \\ &= 1 - \frac{x}{2} - \frac{x^2}{8} - \frac{x}{2} + \frac{x^2}{4} + \frac{3x^2}{8} + \dots \\ &= 1 - x + \frac{x^2}{2} \text{ (app.)} \\ (ii) \frac{1}{(1+x)^2 \sqrt{1+4x}} &= (1+x)^{-2} (1+4x)^{-1/2} \\ &= \left[1 - 2x + \frac{2.3}{1.2}x^2 + \dots \right] \left[1 - \frac{1}{2}(4x) + \dots \right] \\ &= (1 - 2x + \dots)(1 - 2x + \dots) \\ &= 1 - 2x - 2x + 4x^2 + \dots = 1 - 4x \text{ (app.)} \end{aligned}$$

Question-22

If x is so large prove that $\sqrt{x^2+25} - \sqrt{x^2+9} = \frac{8}{x}$ nearly.

Solution:

$$\begin{aligned} \sqrt{x^2+25} - \sqrt{x^2+9} &= x \left[1 + \frac{25}{x^2} \right]^{\frac{1}{2}} - x \left[1 + \frac{9}{x^2} \right]^{\frac{1}{2}} \\ &= x \left[1 + \frac{1}{2} \left[\frac{25}{x^2} \right] + \frac{\frac{1}{2} \left[\frac{-1}{2} \right]}{1.2} \left[\frac{25}{x^2} \right]^2 + \dots \right] - x \left[1 + \frac{1}{2} \left[\frac{9}{x^2} \right] + \frac{\frac{1}{2} \left[\frac{-1}{2} \right]}{1.2} \left[\frac{9}{x^2} \right]^2 + \dots \right] \\ &= x + \frac{25}{2x} - \frac{625}{8x^3} + \dots - x - \frac{9}{2x} + \frac{81}{8x^3} + \dots \\ &= \frac{16}{2x} + \frac{81}{8x^3} + \dots \\ &= \frac{16}{2x} = \frac{8}{x} \text{ approximately} \end{aligned}$$

Question-23

If c is small compared to l , show that $\left[\frac{1}{1+c}\right]^{\frac{1}{2}} + \left[\frac{1}{1-c}\right]^{\frac{1}{2}} = 2 + \frac{3c^2}{4l^2}$ (app)

Solution:

$$\left[\frac{1}{1+c}\right]^{\frac{1}{2}} + \left[\frac{1}{1-c}\right]^{\frac{1}{2}} = \frac{1}{\left[1+\frac{c}{l}\right]^{\frac{1}{2}}} + \frac{1}{\left[1-\frac{c}{l}\right]^{\frac{1}{2}}}$$

$$= \left[1+\frac{c}{l}\right]^{\frac{1}{2}} + \left[1-\frac{c}{l}\right]^{\frac{-1}{2}}$$

Since c is small in comparison with l then $\left|\frac{c}{l}\right| < 1$, $\therefore$ binomial expansion is valid.

$$= 1 + \left[\frac{-1}{2}\right]\left[\frac{c}{l}\right] + \frac{\left[\frac{-1}{2}\right]\left[\frac{-1}{2}-1\right]}{1.2} \left[\frac{-c}{l}\right]^2 + \dots$$

$$+ 1 + \left[\frac{-1}{2}\right]\left[\frac{-c}{l}\right] + \frac{\left[\frac{-1}{2}\right]\left[\frac{-1}{2}-1\right]}{1.2} \left[\frac{-c}{l}\right]^2 + \dots$$

$$= 1 - \frac{c}{2l} + \frac{3c^2}{8l^2} + \dots + 1 + \frac{c}{2l} + \frac{3c^2}{8l^2} + \dots$$

$$= 2 + \frac{3c^2}{4l^2} \text{ approximately.}$$

Question-24

Find the 5th term in the expansion of $(1 - 2x^3)^{11/2}$.

Solution:

$$(1 - 2x^3)^{11/2} = 1 + \left[\frac{11}{2}\right](-2x^3) + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]}{1.2}(-2x^3)^2 + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]}{1.2.3}(-2x^3)^3 + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]\left[\frac{5}{2}\right]}{1.2.3.4}(-2x^3)^4$$

$$5^{\text{th}} \text{ term is } \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]\left[\frac{5}{2}\right]}{1.2.3.4}(-2x^3)^4 = \frac{1155}{8}x^{12}$$

Question-25

Find the $(r+1)^{\text{th}}$ term in the expansion of $(1-x)^{-4}$.

Solution:

T_{r+1} in $(1-x)^{-4}$

$$(1-x)^{-4} = \frac{1}{6} [1.2.3 + 2.3.4.x + \dots (r+1)(r+2)(r+3)x^r + \dots]$$

$$\therefore T_{r+1} = \frac{(r+1)(r+2)(r+3)}{6} x^r$$

Question-26

Show that $x^n = 1 + n\left[1 - \frac{1}{x}\right] + \frac{n(n+1)}{1.2}\left[1 - \frac{1}{x}\right]^2 + \dots$

Solution:

$$\text{R.H.S} = 1 + n\left[1 - \frac{1}{x}\right] + \frac{n(n+1)}{1.2}\left[1 - \frac{1}{x}\right]^2 + \dots$$

$$\text{Put } y = 1 - \frac{1}{x}$$

$$= 1 + ny + \frac{n(n+1)}{1.2}y^2 + \dots$$

$$= (1 - y)^{-n}$$

$$= \left[1 - \left[1 - \frac{1}{x}\right]\right]^{-n}$$

$$= \left[\frac{1}{x}\right]^{-n}$$

$$= x^n$$

$$= \text{L.H.S}$$

Question-27

Find the sum to infinity of the series

$$(i) 1 + \frac{9}{8} + \frac{9.15}{8.16} + \frac{9.15.21}{8.16.24} + \dots$$

$$(ii) 1 - \frac{1}{5} + \frac{1.4}{5.10} - \frac{1.4.7}{5.10.15} + \dots$$

$$(iii) \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots$$

Solution:

$$(i) \text{ Let } S = 1 + \frac{9}{8} + \frac{9.15}{8.16} + \frac{9.15.21}{8.16.24} + \dots$$

$$= 1 + \frac{9}{6}\left(\frac{6}{8}\right) + \frac{\left(\frac{9}{6}\right)\left(\frac{5}{6}\right)}{1.2}\left(\frac{6}{8}\right)^2 + \dots$$

$$= \left[1 - \frac{6}{8}\right]^{-9}$$

$$\left[1 - \frac{3}{4}\right]^{-9} = \left[\frac{1}{4}\right]^{-\frac{3}{2}} = 4^{\frac{3}{2}} = 4^1 \cdot 4^{\frac{1}{2}} = \sqrt[4]{4} = 4(2) = 8$$

$$\begin{aligned}
 \text{(ii) Let } S &= 1 - \frac{1}{5} + \frac{1.4}{5.10} - \frac{1.4.7}{5.10.15} + \dots \\
 &= 1 - \left(\frac{1}{3} \right) \left(\frac{3}{5} \right) + \frac{\left[\frac{1}{3} \right] \left[\frac{4}{3} \right]}{1.2} \left(\frac{3}{5} \right)^2 + \dots \\
 &= \left[1 + \frac{3}{5} \right]^{-\frac{1}{3}} \\
 &= \left[\frac{8}{5} \right]^{-\frac{1}{3}} \\
 &= \left[\frac{5}{8} \right]^{\frac{1}{3}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) Let } S &= \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots \\
 S + 1 &= 1 + \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots \\
 &= 1 + \left(\frac{3}{2} \right) \left(\frac{2}{4} \right) + \frac{\left(\frac{3}{2} \right) \left(\frac{5}{2} \right)}{1.2} \left(\frac{2}{4} \right)^2 + \dots \\
 &= \left(1 - \frac{2}{4} \right)^{-3/2} = \left(\frac{1}{2} \right)^{-3/2} = 2^{3/2} \\
 S + 1 &= 2^{3/2}
 \end{aligned}$$

$$\text{Therefore } S = 2^{3/2} - 1$$

Question-28

Show that the coefficient of x^n in the infinite series $1 +$

$$\frac{b+ax}{1!} + \frac{b+ax)^2}{2!} + \frac{b+ax)^3}{3!} + \dots \text{ is } \frac{e^b a^n}{n!}.$$

$$\text{(ii) Show that } \left(1 + \frac{1}{2!} + \frac{1}{4!} + \dots \right)^2 = 1 + \left(1 + \frac{1}{3!} + \frac{1}{5!} + \dots \right)^2.$$

$$\text{(iii) Show that } 2 \left\{ 1 + \frac{(\log n)^2}{2} + \frac{(\log n)^4}{4!} + \dots \right\} = n + c.$$

Solution:

$$\text{(i) } 1 + \frac{y}{1!} + \frac{y^2}{2!} + \frac{y^3}{3!} + \dots = e^y = e^{b+ax} = e^b \cdot e^{ax} = e^b \left(1 + \frac{ax}{1!} + \frac{(ax)^2}{2!} + \dots \right)$$

$$\text{Coefficient of } x^n = e^b \cdot \left(\frac{a^n}{n!} \right)$$

$$\begin{aligned}
 \text{(ii) L.H.S} &= \left(1 + \frac{1}{2!} + \frac{1}{4!} + \dots\right)^2 = \left(\frac{e + e^{-1}}{2}\right)^2 \\
 &= 1 + \frac{e^2 + e^{-2} - 2}{4} \\
 &= \frac{4 + e^2 + e^{-2} - 2}{4} \\
 &= \frac{e^2 + e^{-2} + 2}{4} \\
 &= \text{R.H.S}
 \end{aligned}$$

$$\text{(iii) L.H.S} = 2 \left\{ 1 + \frac{(\log n)^2}{2!} + \frac{(\log n)^4}{4!} + \dots \right\}$$

Put $\log n = y$

$$\begin{aligned}
 2 \left\{ 1 + \frac{(\log n)^2}{2!} + \frac{(\log n)^4}{4!} + \dots \right\} &= 2 \left(\frac{e^y + e^{-y}}{2} \right) \\
 &= e^y + e^{-y} \\
 &= e^{\log n} + e^{-\log n} \\
 &= e^{\log n} + e^{\log 1/n} \\
 &= n + \frac{1}{n}
 \end{aligned}$$

Question-29

Show that $\log a - \log b = \frac{a-b}{a} + \frac{1}{2} \left(\frac{a-b}{a} \right)^2 + \frac{1}{3} \left(\frac{a-b}{a} \right)^3 + \dots$

Solution:

$$\text{R.H.S} = \frac{a-b}{a} + \frac{1}{2} \left(\frac{a-b}{a} \right)^2 + \frac{1}{3} \left(\frac{a-b}{a} \right)^3 + \dots$$

$$\text{Put } y = \frac{a-b}{a}$$

$$y + \frac{y^2}{2} + \frac{y^3}{3} + \dots = -\log(1-y)$$

$$= -\log \left(1 - \frac{a-b}{a} \right)$$

$$= -\log \left(\frac{b}{a} \right)$$

$$= \log \left(\frac{a}{b} \right)$$

$$= \log a - \log b$$

$$= \text{L.H.S}$$

Question-30

Prove that $\log \frac{n+1}{n-1} = \frac{2n}{n^2+1} + \frac{1}{3} \left(\frac{2n}{n^2+1} \right)^3 + \dots$

Solution:

$$\text{R.H.S} = \frac{2n}{n^2+1} + \frac{1}{3} \left(\frac{2n}{n^2+1} \right)^3 + \dots$$

$$\text{Put } \frac{2n}{n^2+1} = y$$

$$y + \frac{y^3}{3} + \frac{y^5}{5} + \dots = \frac{1}{2} \log \left(\frac{1+y}{1-y} \right)$$

$$= \frac{1}{2} \log \left(\frac{1 + \frac{2n}{n^2+1}}{1 - \frac{2n}{n^2+1}} \right)$$

$$= \frac{1}{2} \log \left(\frac{n^2+1+2n}{n^2+1-2n} \right)$$

$$= \frac{1}{2} \log \left(\frac{n+1}{n-1} \right)^2$$

$$= \log \left(\frac{n+1}{n-1} \right)$$

$$= \text{L.H.S}$$

Question-31

Find the sum to infinity the series $\frac{1}{x-1} + \frac{1}{3} \frac{1}{(x-1)^3} + \frac{1}{5} \frac{1}{(x-1)^5} + \dots$

Solution:

$$\frac{1}{x-1} + \frac{1}{3} \frac{1}{(x-1)^3} + \frac{1}{5} \frac{1}{(x-1)^5} + \dots$$

$$\text{Put } y = \frac{1}{x-1}$$

$$\frac{1}{y} + \frac{1}{3} \frac{1}{y^3} + \frac{1}{5} \frac{1}{y^5} + \dots = \frac{1}{2} \log \left(\frac{1+y}{1-y} \right)$$

$$= \frac{1}{2} \log \left(\frac{1 + \frac{1}{1-x}}{1 - \frac{1}{1-x}} \right)$$

$$= \frac{1}{2} \log \left(\frac{1-x+1}{1-x-1} \right)$$

$$= \frac{1}{2} \log \left(\frac{2-x}{-x} \right)$$

$$= \frac{1}{2} \log \left(\frac{x+2}{x} \right)$$

Question-32

If x is so small show that

$$(i) \frac{\sqrt{1-x}}{1+x} = 1 - x + \frac{x^2}{2} \text{ (app.)}$$

$$(ii) \frac{1}{(1+x)^2 \sqrt{1+4x}} = 1 - 4x \text{ (app.)}$$

Solution:

$$(i) \frac{\sqrt{1-x}}{1+x} = (1-x)^{\frac{1}{2}} (1+x)^{-\frac{1}{2}}$$

$$= \left[1 - \frac{1}{2}x + \frac{\left[\frac{1}{2}\right]\left[\frac{-1}{2}\right]}{1.2}x^2 + \dots \right] \left[1 - \frac{1}{2}x + \frac{\frac{1}{2} \cdot \frac{3}{2}}{1.2}x^2 + \dots \right]$$

$$= \left[1 - \frac{x}{2} - \frac{x^2}{8} \right] \left[1 - \frac{x}{2} + \frac{3x^2}{8} \right]$$

$$= 1 - \frac{x}{2} - \frac{x^2}{8} - \frac{x}{2} + \frac{x^2}{4} + \frac{3x^2}{8} + \dots$$

$$= 1 - x + \frac{x^2}{2} \text{ (app.)}$$

$$(ii) \frac{1}{(1+x)^2 \sqrt{1+4x}} = (1+x)^{-2} (1+4x)^{-1/2}$$

$$= \left[1 - 2x + \frac{2 \cdot 3}{1.2}x^2 + \dots \right] \left[1 - \frac{1}{2}(4x) + \dots \right]$$

$$= (1 - 2x + \dots)(1 - 2x + \dots)$$

$$= 1 - 2x - 2x + 4x^2 + \dots = 1 - 4x \text{ (app.)}$$

Question-33

If x is so large prove that $\sqrt{x^2+25} - \sqrt{x^2+9} = \frac{8}{x}$ nearly.

Solution:

$$\sqrt{x^2+25} - \sqrt{x^2+9} = x \left[1 + \frac{25}{x^2} \right]^{\frac{1}{2}} - x \left[1 + \frac{9}{x^2} \right]^{\frac{1}{2}}$$

$$= x \left[1 + \frac{1}{2} \left[\frac{25}{x^2} \right] + \frac{\frac{1}{2} \left[\frac{-1}{2} \right]}{1.2} \left[\frac{25}{x^2} \right]^2 + \dots \right] - x \left[1 + \frac{1}{2} \left[\frac{9}{x^2} \right] + \frac{\frac{1}{2} \left[\frac{-1}{2} \right]}{1.2} \left[\frac{9}{x^2} \right]^2 + \dots \right]$$

$$= x + \frac{25}{2x} - \frac{625}{8x^3} + \dots - x - \frac{9}{2x} + \frac{81}{8x^3} + \dots$$

$$= \frac{16}{2x} + \frac{81}{8x^3} + \dots$$

$$= \frac{16}{2x} = \frac{8}{x} \text{ approximately}$$

Question-34

If c is small compared to l , show that $\left[\frac{1}{1+c}\right]^{\frac{1}{2}} + \left[\frac{1}{1-c}\right]^{\frac{1}{2}} = 2 + \frac{3c^2}{4l^2}$ (app)

Solution:

$$\begin{aligned}\left[\frac{1}{1+c}\right]^{\frac{1}{2}} + \left[\frac{1}{1-c}\right]^{\frac{1}{2}} &= \frac{1}{\left[1+\frac{c}{l}\right]^{\frac{1}{2}}} + \frac{1}{\left[1-\frac{c}{l}\right]^{\frac{1}{2}}} \\ &= \left[1+\frac{c}{l}\right]^{\frac{1}{2}} + \left[1-\frac{c}{l}\right]^{\frac{1}{2}}\end{aligned}$$

Since c is small in comparison with l then $\left|\frac{c}{l}\right| < 1$, $\therefore$ binomial expansion is valid.

$$\begin{aligned}&= 1 + \left[\frac{-1}{2}\right]\left[\frac{c}{l}\right] + \frac{\left[\frac{-1}{2}\right]\left[\frac{-1}{2}-1\right]}{1.2}\left[\frac{c}{l}\right]^2 + \dots\dots\dots \\ &+ 1 + \left[\frac{-1}{2}\right]\left[\frac{-c}{l}\right] + \frac{\left[\frac{-1}{2}\right]\left[\frac{-1}{2}-1\right]}{1.2}\left[\frac{-c}{l}\right]^2 + \dots\dots\dots \\ &= 1 - \frac{c}{2l} + \frac{3c^2}{8l^2} + \dots\dots + 1 + \frac{c}{2l} + \frac{3c^2}{8l^2} + \dots\dots \\ &= 2 + \frac{3c^2}{4l^2} \text{ approximately.}\end{aligned}$$

Question-35

Find the 5th term in the expansion of $(1 - 2x^3)^{11/2}$.

Solution:

$$\begin{aligned}(1 - 2x^3)^{11/2} &= 1 + \left[\frac{11}{2}\right](-2x^3) + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]}{1.2}(-2x^3)^2 + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]}{1.2.3}(-2x^3)^3 + \frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]\left[\frac{5}{2}\right]}{1.2.3.4}(-2x^3)^4 \\ \text{5th term is } &\frac{\left[\frac{11}{2}\right]\left[\frac{2}{9}\right]\left[\frac{7}{2}\right]\left[\frac{5}{2}\right]}{1.2.3.4}(-2x^3)^4 = \frac{1155}{8}x^{12}\end{aligned}$$

Question-36

Find the $(r + 1)^{\text{th}}$ term in the expansion of $(1 - x)^{-4}$.

Solution:

$$\begin{aligned}&T_{r+1} \text{ in } (1 - x)^{-4} \\ (1-x)^{-4} &= \frac{1}{6} [1.2.3 + 2.3.4.x + \dots\dots\dots (r+1)(r+2)(r+3)x^r + \dots\dots\dots] \\ \therefore T_{r+1} &= \frac{(r+1)(r+2)(r+3)}{6} x^r\end{aligned}$$

Question-37

Show that $x^n = 1 + n\left[1 - \frac{1}{x}\right] + \frac{n(n+1)}{1.2}\left[1 - \frac{1}{x}\right]^2 + \dots$

Solution:

$$\text{R.H.S} = 1 + n\left[1 - \frac{1}{x}\right] + \frac{n(n+1)}{1.2}\left[1 - \frac{1}{x}\right]^2 + \dots$$

$$\text{Put } y = 1 - \frac{1}{x}$$

$$= 1 + ny + \frac{n(n+1)}{1.2}y^2 + \dots$$

$$= (1 - y)^{-n}$$

$$= \left[1 - \left[1 - \frac{1}{x}\right]\right]^{-n}$$

$$= \left[\frac{1}{x}\right]^{-n}$$

$$= x^n$$

$$= \text{L.H.S}$$

Question-38

Find the sum to infinity of the series

$$(i) 1 + \frac{9}{8} + \frac{9.15}{8.16} + \frac{9.15.21}{8.16.24} + \dots$$

$$(ii) 1 - \frac{1}{5} + \frac{1.4}{5.10} - \frac{1.4.7}{5.10.15} + \dots$$

$$(iii) \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots$$

Solution:

$$(i) \text{ Let } S = 1 + \frac{9}{8} + \frac{9.15}{8.16} + \frac{9.15.21}{8.16.24} + \dots$$

$$= 1 + \frac{9}{6}\left(\frac{6}{8}\right) + \frac{\left(\frac{9}{6}\right)\left(\frac{5}{6}\right)}{1.2}\left(\frac{6}{8}\right)^2 + \dots$$

$$= \left[1 - \frac{6}{8}\right]^{-\frac{9}{6}}$$

$$\left[1 - \frac{3}{4}\right]^{-\frac{3}{2}} = \left[\frac{1}{4}\right]^{-\frac{3}{2}} = 4^{\frac{3}{2}} = 4^1 \cdot 4^{\frac{1}{2}} = \sqrt[4]{4} = 4(2) = 8$$

$$\begin{aligned}
 \text{(ii) Let } S &= 1 - \frac{1}{5} + \frac{1.4}{5.10} - \frac{1.4.7}{5.10.15} + \dots \\
 &= 1 - \left(\frac{1}{3} \right) \left(\frac{3}{5} \right) + \frac{\left[\frac{1}{3} \right] \left[\frac{4}{3} \right]}{1.2} \left(\frac{3}{5} \right)^2 + \dots \\
 &= \left[1 + \frac{3}{5} \right]^{-\frac{1}{3}} \\
 &= \left[\frac{8}{5} \right]^{-\frac{1}{3}} \\
 &= \left[\frac{5}{8} \right]^{\frac{1}{3}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) Let } S &= \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots \\
 S + 1 &= 1 + \frac{3}{4} + \frac{3.5}{4.8} + \frac{3.5.7}{4.8.12} + \dots \\
 &= 1 + \left(\frac{3}{2} \right) \left(\frac{2}{4} \right) + \frac{\left(\frac{3}{2} \right) \left(\frac{5}{2} \right)}{1.2} \left(\frac{2}{4} \right)^2 + \dots \\
 &= \left(1 - \frac{2}{4} \right)^{-3/2} = \left(\frac{1}{2} \right)^{-3/2} = 2^{3/2} \\
 S + 1 &= 2^{3/2}
 \end{aligned}$$

$$\text{Therefore } S = 2^{3/2} - 1$$

Question-39

(i) Show that the coefficient of x^n in the infinite series $1 +$

$$\frac{b+ax}{1!} + \frac{b+ax)^2}{2!} + \frac{b+ax)^3}{3!} + \dots \text{ is } \frac{e^b a^n}{n!}.$$

(ii) Show that $\left(1 + \frac{1}{2!} + \frac{1}{4!} + \dots \right)^2 = 1 + \left(1 + \frac{1}{3!} + \frac{1}{5!} + \dots \right)^2.$

(iii) Show that $2 \left[1 + \frac{(\log n)^2}{2!} + \frac{(\log n)^4}{4!} + \dots \right] = n + c.$

Solution:

$$(i) 1 + \frac{y}{1!} + \frac{y^2}{2!} + \frac{y^3}{3!} + \dots = e^y = e^{b+ax} = e^b \cdot e^{ax} = e^b \left(1 + \frac{ax}{1!} + \frac{(ax)^2}{2!} + \dots \right)$$

$$\text{Coefficient of } x^n = e^b \cdot \left(\frac{a^n}{n!} \right)$$

$$(ii) \text{ L.H.S} = \left(1 + \frac{1}{2!} + \frac{1}{4!} + \dots \right)^2 = \left(\frac{e+e^{-1}}{2} \right)^2$$

$$\begin{aligned}
 &= 1 + \frac{e^2 + e^{-2} - 2}{4} \\
 &= \frac{4 + e^2 + e^{-2} - 2}{4} \\
 &= \frac{e^2 + e^{-2} + 2}{4} \\
 &= \text{R.H.S}
 \end{aligned}$$

$$(iii) \text{ L.H.S} = 2 \left\{ 1 + \frac{(\log n)^2}{2!} + \frac{(\log n)^4}{4!} + \dots \right\}$$

Put $\log n = y$

$$\begin{aligned}
 2 \left\{ 1 + \frac{(\log n)^2}{2!} + \frac{(\log n)^4}{4!} + \dots \right\} &= 2 \left(\frac{e^y + e^{-y}}{2} \right) \\
 &= e^y + e^{-y} \\
 &= e^{\log n} + e^{-\log n} \\
 &= e^{\log n} + e^{\log 1/n} \\
 &= n + \frac{1}{n}
 \end{aligned}$$

Question-40

Show that $\log a - \log b = \frac{a-b}{a} + \frac{1}{2} \left(\frac{a-b}{a} \right)^2 + \frac{1}{3} \left(\frac{a-b}{a} \right)^3 + \dots$

Solution:

$$\text{R.H.S} = \frac{a-b}{a} + \frac{1}{2} \left(\frac{a-b}{a} \right)^2 + \frac{1}{3} \left(\frac{a-b}{a} \right)^3 + \dots$$

$$\text{Put } y = \frac{a-b}{a}$$

$$y + \frac{y^2}{2} + \frac{y^3}{3} + \dots = -\log(1-y)$$

$$= -\log \left(1 - \frac{a-b}{a} \right)$$

$$= -\log \left(\frac{b}{a} \right)$$

$$= \log \left(\frac{a}{b} \right)$$

$$= \log a - \log b$$

$$= \text{L.H.S}$$

Question-41

Prove that $\log \frac{n+1}{n-1} = \frac{2n}{n^2+1} + \frac{1}{3} \left(\frac{2n}{n^2+1} \right)^3 + \dots$

Solution:

$$\text{R.H.S} = \frac{2n}{n^2+1} + \frac{1}{3} \left(\frac{2n}{n^2+1} \right)^3 + \dots$$

$$\text{Put } \frac{2n}{n^2+1} = y$$

$$\begin{aligned} y + \frac{y^3}{3} + \frac{y^5}{5} + \dots &= \frac{1}{2} \log \left(\frac{1+y}{1-y} \right) \\ &= \frac{1}{2} \log \left(\frac{1 + \frac{2n}{n^2+1}}{1 - \frac{2n}{n^2+1}} \right) \\ &= \frac{1}{2} \log \left(\frac{n^2+1+2n}{n^2+1-2n} \right) \\ &= \frac{1}{2} \log \left(\frac{n+1}{n-1} \right)^2 \\ &= \log \left(\frac{n+1}{n-1} \right) \\ &= \text{L.H.S} \end{aligned}$$

Question-42

Find the sum to infinity the series $\frac{1}{x-1} + \frac{1}{3} \frac{1}{(x-1)^3} + \frac{1}{5} \frac{1}{(x-1)^5} + \dots$

Solution:

$$\frac{1}{x-1} + \frac{1}{3} \frac{1}{(x-1)^3} + \frac{1}{5} \frac{1}{(x-1)^5} + \dots$$

$$\text{Put } y = \frac{1}{x-1}$$

$$\begin{aligned} \frac{1}{y} + \frac{1}{3} \frac{1}{y^3} + \frac{1}{5} \frac{1}{y^5} + \dots &= \frac{1}{2} \log \left(\frac{1+y}{1-y} \right) \\ &= \frac{1}{2} \log \left(\frac{1 + \frac{1}{1-x}}{1 - \frac{1}{1-x}} \right) \\ &= \frac{1}{2} \log \left(\frac{1-x+1}{1-x-1} \right) \\ &= \frac{1}{2} \log \left(\frac{2-x}{-x} \right) \\ &= \frac{1}{2} \log \left(\frac{x+2}{x} \right) \end{aligned}$$