

DAY TWENTY TWO

Inverse Trigonometric Function

Learning & Revision for the Day

• Inverse Trigonometric Function

• Properties of Inverse Trigonometric Function

Inverse Trigonometric Function

Trigonometric functions are not one-one and onto on their natural domains and ranges, so their inverse do not exists in the whole domain. If we restrict their domain and range, then their inverse may exists.

$y = f(x) = \sin x$. Then, its inverse is $x = \sin^{-1} y$.

NOTE • $\sin^{-1} y \neq (\sin y)^{-1}$ • $\sin^{-1} y \neq \sin\left(\frac{1}{y}\right)$

The value of an inverse trigonometric functions which lies in its principal value branch is called the principal value of that inverse trigonometric function.

Domain and range of inverse trigonometric functions

Function	Domain	Range (Principal Value Branch)
$\sin^{-1} x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$\cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1} x$	R	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$\cot^{-1} x$	R	$(0, \pi)$
$\sec^{-1} x$	$R - (-1, 1)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$
$\operatorname{cosec}^{-1} x$	$R - (-1, 1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

Properties of Inverse Trigonometric Functions

$$1. \quad (i) \quad \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}; \quad (-1 \leq x \leq 1)$$

$$(ii) \quad \tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}; \quad x \in R$$

$$(iii) \quad \sec^{-1} x + \operatorname{cosec}^{-1} x = \frac{\pi}{2}; \quad (x \leq -1 \text{ or } x \geq 1)$$

$$2. \quad (i) \quad \sin^{-1}(-x) = -\sin^{-1} x; \quad (-1 \leq x \leq 1)$$

$$(ii) \quad \cos^{-1}(-x) = \pi - \cos^{-1} x; \quad (-1 \leq x \leq 1)$$

$$(iii) \quad \tan^{-1}(-x) = -\tan^{-1} x; \quad (-\infty < x < \infty)$$

$$(iv) \quad \cot^{-1}(-x) = \pi - \cot^{-1} x; \quad (-\infty < x < \infty)$$

$$(v) \quad \sec^{-1}(-x) = \pi - \sec^{-1} x; \quad x \leq -1 \text{ or } x \geq 1$$

$$(vi) \quad \operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1} x; \quad (x \leq -1 \text{ or } x \geq 1)$$

$$3. \quad (i) \quad \sin^{-1}(\sin x) \text{ is a periodic function with period } 2\pi.$$

$$\sin^{-1}(\sin x) = \begin{cases} x, & x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \\ \pi - x, & x \in \left[\frac{\pi}{2}, \frac{3\pi}{2}\right] \\ x - 2\pi, & x \in \left[\frac{3\pi}{2}, \frac{5\pi}{2}\right] \\ 3\pi - x, & x \in \left[\frac{5\pi}{2}, \frac{7\pi}{2}\right] \end{cases}$$

$$(ii) \quad \cos^{-1}(\cos x) \text{ is a periodic function with period } 2\pi.$$

$$\cos^{-1}(\cos x) = \begin{cases} x, & x \in [0, \pi] \\ 2\pi - x, & x \in [\pi, 2\pi] \\ x - 2\pi, & x \in [2\pi, 3\pi] \\ 4\pi - x, & x \in [3\pi, 4\pi] \end{cases}$$

$$(iii) \quad \tan^{-1}(\tan x) \text{ is a periodic function with period } \pi.$$

$$\tan^{-1}(\tan x) = \begin{cases} x, & x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \\ x - \pi, & x \in \left[\frac{\pi}{2}, \frac{3\pi}{2}\right] \\ x - 2\pi, & x \in \left[\frac{3\pi}{2}, \frac{5\pi}{2}\right] \\ x - 3\pi, & x \in \left[\frac{5\pi}{2}, \frac{7\pi}{2}\right] \end{cases}$$

$$(iv) \quad \cot^{-1}(\cot x) \text{ is a periodic function with period } \pi.$$

$$\cot^{-1}(\cot x) = x; \quad 0 < x < \pi$$

$$(v) \quad \sec^{-1}(\sec x) \text{ is a periodic function with period } 2\pi.$$

$$\sec^{-1}(\sec x) = x; \quad 0 \leq x < \frac{\pi}{2} \text{ or } \frac{\pi}{2} < x \leq \pi$$

$$(vi) \quad \operatorname{cosec}^{-1}(\operatorname{cosec} x) \text{ is a periodic function with period } 2\pi.$$

$$\operatorname{cosec}^{-1}(\operatorname{cosec} x) = x; \quad -\frac{\pi}{2} \leq x < 0 \text{ or } 0 < x \leq \frac{\pi}{2}$$

$$4. \quad (i) \quad \sin^{-1}\left(\frac{1}{x}\right) = \operatorname{cosec}^{-1} x, \text{ if } x \in (-\infty, -1] \cup [1, \infty)$$

$$(ii) \quad \cos^{-1}\left(\frac{1}{x}\right) = \sec^{-1} x, \text{ if } x \in (-\infty, -1] \cup [1, \infty)$$

$$(iii) \quad \tan^{-1}\left(\frac{1}{x}\right) = \begin{cases} \cot^{-1} x, & \text{if } x > 0 \\ -\pi + \cot^{-1} x, & \text{if } x < 0 \end{cases}$$

$$5. \quad (i) \quad \sin^{-1} x = \cos^{-1} \sqrt{1-x^2}$$

$$= \tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) = \cot^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right)$$

$$= \sec^{-1}\left(\frac{1}{\sqrt{1-x^2}}\right) = \operatorname{cosec}^{-1}\left(\frac{1}{x}\right), \text{ if } x \in (0, 1)$$

$$(ii) \quad \cos^{-1} x = \sin^{-1} \sqrt{1-x^2}$$

$$= \tan^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right) = \cot^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) = \sec^{-1}\left(\frac{1}{x}\right)$$

$$= \operatorname{cosec}^{-1}\left(\frac{1}{\sqrt{1-x^2}}\right), \text{ if } x \in (0, 1)$$

$$(iii) \quad \tan^{-1} x = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right)$$

$$= \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right) = \cot^{-1}\left(\frac{1}{x}\right)$$

$$= \operatorname{cosec}^{-1}\left(\frac{\sqrt{1+x^2}}{x}\right) = \sec^{-1}(\sqrt{1+x^2}), \text{ if } x \in (0, \infty)$$

$$6. \quad (i) \quad \sin^{-1} x + \sin^{-1} y$$

$$= \begin{cases} \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2}); & |x|, |y| \leq 1 \text{ and } x^2 + y^2 \leq 1 \text{ or } (xy < 0 \text{ and } x^2 + y^2 > 1) \\ \pi - \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2}); & 0 < x, y \leq 1 \text{ and } x^2 + y^2 > 1 \\ -\pi - \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2}); & -1 \leq x, y < 0 \text{ and } x^2 + y^2 > 1 \end{cases}$$

$$(ii) \quad \sin^{-1} x - \sin^{-1} y$$

$$= \begin{cases} \sin^{-1}(x\sqrt{1-y^2} - y\sqrt{1-x^2}); & |x|, |y| \leq 1 \text{ and } x^2 + y^2 \leq 1 \text{ or } (xy > 0 \text{ and } x^2 + y^2 > 1) \\ \pi - \sin^{-1}(x\sqrt{1-y^2} - y\sqrt{1-x^2}); & 0 < x \leq 1, -1 \leq y < 0 \text{ and } x^2 + y^2 > 1 \\ -\pi - \sin^{-1}(x\sqrt{1-y^2} - y\sqrt{1-x^2}); & -1 \leq x < 0, 0 < y \leq 1 \text{ and } x^2 + y^2 > 1 \end{cases}$$

$$(iii) \quad \cos^{-1} x + \cos^{-1} y$$

$$= \begin{cases} \cos^{-1}\{xy - \sqrt{(1-x^2)}\sqrt{(1-y^2)}\}; & |x|, |y| \leq 1 \text{ and } x + y \geq 0 \\ 2\pi - \cos^{-1}\{xy - \sqrt{(1-x^2)}\sqrt{(1-y^2)}\}; & |x|, |y| \leq 1 \text{ and } x + y < 0 \end{cases}$$

$$(iv) \quad \cos^{-1} x - \cos^{-1} y$$

$$= \begin{cases} \cos^{-1}\{xy + \sqrt{(1-x^2)}\sqrt{(1-y^2)}\}; & |x|, |y| \leq 1 \text{ and } x \leq y \\ -\cos^{-1}\{xy + \sqrt{(1-x^2)}\sqrt{(1-y^2)}\}; & -1 \leq y \leq 0, 0 < x \leq 1 \text{ and } x \geq y \end{cases}$$

$$(v) \tan^{-1} x + \tan^{-1} y = \begin{cases} \tan^{-1} \left(\frac{x+y}{1-xy} \right); & xy < 1 \\ \pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right); & x > 0, y > 0, xy > 1 \\ -\pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right); & x < 0, y < 0, xy > 1 \end{cases}$$

$$(vi) \tan^{-1} x - \tan^{-1} y = \begin{cases} \tan^{-1} \left(\frac{x-y}{1+xy} \right); & xy > -1 \\ \pi + \tan^{-1} \left(\frac{x-y}{1+xy} \right); & xy < -1, x > 0, y < 0 \\ -\pi + \tan^{-1} \left(\frac{x-y}{1+xy} \right); & xy < -1, x < 0, y > 0 \end{cases}$$

$$7. (i) 2 \sin^{-1} x = \begin{cases} \sin^{-1} \{2x \sqrt{1-x^2}\}; & -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ \pi - \sin^{-1} (2x \sqrt{1-x^2}); & \frac{1}{\sqrt{2}} \leq x \leq 1 \\ -\pi - \sin^{-1} (2x \sqrt{1-x^2}); & -1 \leq x < -\frac{1}{\sqrt{2}} \end{cases}$$

$$(ii) 2 \cos^{-1} x = \begin{cases} \cos^{-1} (2x^2 - 1); & 0 \leq x \leq 1 \\ 2\pi - \cos^{-1} (2x^2 - 1); & -1 \leq x < 0 \end{cases}$$

$$(iii) 2 \tan^{-1} x = \begin{cases} \tan^{-1} \left(\frac{2x}{1-x^2} \right); & -1 < x < 1 \\ \pi + \tan^{-1} \left(\frac{2x}{1-x^2} \right); & x > 1 \\ -\pi + \tan^{-1} \left(\frac{2x}{1-x^2} \right); & x < -1 \end{cases}$$

$$(iv) 2 \tan^{-1} x = \begin{cases} \sin^{-1} \left(\frac{2x}{1+x^2} \right); & -1 \leq x \leq 1 \\ \pi - \sin^{-1} \left(\frac{2x}{1+x^2} \right); & x > 1 \\ -\pi - \sin^{-1} \left(\frac{2x}{1+x^2} \right); & x < -1 \end{cases}$$

$$(v) 2 \tan^{-1} x = \begin{cases} \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right); & 0 \leq x < \infty \\ -\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right); & -\infty < x \leq 0 \end{cases}$$

NOTE

- If $\sin^{-1} x + \sin^{-1} y = \theta$, then $\cos^{-1} x + \cos^{-1} y = \pi - \theta$
- If $\cos^{-1} x + \cos^{-1} y = \theta$, then $\sin^{-1} x + \sin^{-1} y = \pi - \theta$

$$8. (i) 3 \sin^{-1} x = \begin{cases} \sin^{-1} (3x - 4x^3), & \text{if } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ \pi - \sin^{-1} (3x - 4x^3), & \text{if } \frac{1}{2} < x \leq 1 \\ -\pi - \sin^{-1} (3x - 4x^3), & \text{if } -1 \leq x < -\frac{1}{2} \end{cases}$$

$$(ii) 3 \cos^{-1} x = \begin{cases} \cos^{-1} (4x^3 - 3x), & \text{if } \frac{1}{2} \leq x \leq 1 \\ 2\pi - \cos^{-1} (4x^3 - 3x), & \text{if } -\frac{1}{2} \leq x < \frac{1}{2} \\ 2\pi + \cos^{-1} (4x^3 - 3x), & \text{if } -1 \leq x < -\frac{1}{2} \end{cases}$$

$$(iii) 3 \tan^{-1} x = \begin{cases} \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right), & \text{if } -\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}} \\ \pi + \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right), & \text{if } x > \frac{1}{\sqrt{3}} \\ -\pi + \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right), & \text{if } x < -\frac{1}{\sqrt{3}} \end{cases}$$

DAY PRACTICE SESSION 1

FOUNDATION QUESTIONS EXERCISE

1 The principal value of $\sin^{-1} \left(\cos \frac{33\pi}{5} \right)$ is **→ NCERT Exemplar**

- (a) $\frac{3\pi}{5}$ (b) $\frac{7\pi}{5}$ (c) $\frac{\pi}{10}$ (d) $-\frac{\pi}{10}$

2 If $\sum_{i=1}^{20} \sin^{-1} x_i = 10\pi$, then $\sum_{i=1}^{20} x_i$ is equal to

- (a) 20 (b) 10 (c) 0 (d) None of these

3 The domain of the function defined by $f(x) = \sin^{-1} \sqrt{x-1}$ is **→ NCERT Exemplar**

- (a) [1, 2] (b) [-1, 1] (c) [0, 1] (d) None of these

4 The value of $\cos (2 \cos^{-1} x + \sin^{-1} x)$ at $x = \frac{1}{5}$ is

- (a) 1 (b) 3 (c) 0 (d) $-\frac{2\sqrt{6}}{5}$

5 The value of $\cos[\tan^{-1}\{\sin(\cot^{-1} x)\}]$ is

- (a) $\frac{1}{\sqrt{x^2+2}}$ (b) $\sqrt{\frac{x^2+2}{x^2+1}}$ (c) $\sqrt{\frac{x^2+1}{x^2+2}}$ (d) $\frac{1}{\sqrt{x^2+1}}$

6 The equation $\tan^{-1} x - \cot^{-1} x = \tan^{-1} \left(\frac{1}{\sqrt{3}} \right)$ has

- (a) no solution **→ NCERT Exemplar**
 (b) unique solution
 (c) infinite number of solutions
 (d) two solutions

7 Let $\tan^{-1} y = \tan^{-1} x + \tan^{-1} \left(\frac{2x}{1-x^2} \right)$, where $|x| < \frac{1}{\sqrt{3}}$. Then,

a value of y is

→ JEE Mains 2015

- (a) $\frac{3x-x^3}{1-3x^2}$ (b) $\frac{3x+x^3}{1-3x^2}$
(c) $\frac{3x-x^3}{1+3x^2}$ (d) $\frac{3x+x^3}{1+3x^2}$

8 If $\theta = \tan^{-1} a$, $\phi = \tan^{-1} b$ and $ab = -1$, then $(\theta - \phi)$ is equal to

- (a) 0 (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{2}$ (d) None of these

9 The range of

$$f(x) = |3 \tan^{-1} x - \cos^{-1}(0)| - \cos^{-1}(-1)$$

- (a) $[-\pi, \pi]$ (b) $(-\pi, \pi)$ (c) $[-\pi, \pi]$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

10 The number of solutions of the equation

$$\cos(\cos^{-1} x) = \operatorname{cosec}(\operatorname{cosec}^{-1} x)$$

- (a) 2 (b) 3 (c) 4 (d) 1

11 The value of $\cot \left(\operatorname{cosec}^{-1} \frac{5}{3} + \tan^{-1} \frac{2}{3} \right)$ is

→ AIEEE 2008

- (a) $\frac{5}{17}$ (b) $\frac{6}{17}$ (c) $\frac{3}{17}$ (d) $\frac{4}{17}$

12 If $\tan(\cos^{-1} x) = \sin \left(\cot^{-1} \frac{1}{2} \right)$, then x is equal to

- (a) $\pm \frac{5}{3}$ (b) $\pm \frac{\sqrt{5}}{3}$ (c) $\pm \frac{5}{\sqrt{3}}$ (d) None of these

13 If $\cos^{-1} x + \cos^{-1} y = \frac{\pi}{2}$ and $\tan^{-1} x - \tan^{-1} y = 0$, then

$$x^2 + xy + y^2 \text{ is equal to}$$

- (a) 0 (b) $\frac{1}{\sqrt{2}}$ (c) $\frac{3}{2}$ (d) $\frac{1}{8}$

14 If $\sin^{-1} \left(\frac{x}{5} \right) + \operatorname{cosec}^{-1} \left(\frac{5}{4} \right) = \frac{\pi}{2}$, then the value of x is

→ AIEEE 2007

- (a) 1 (b) 3 (c) 4 (d) 5

15 If $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$, then the value of

$$\sum \frac{(x^{101} + y^{101})(x^{202} + y^{202})}{(x^{303} + y^{303})(x^{404} + y^{404})}$$

- (a) 0 (b) 1 (c) 2 (d) 3

16 The root of the equation

$$\tan^{-1} \left(\frac{x-1}{x+1} \right) + \tan^{-1} \left(\frac{2x-1}{2x+1} \right) = \tan^{-1} \left(\frac{23}{36} \right)$$

- (a) $-\frac{3}{8}$ (b) $-\frac{1}{2}$ (c) $\frac{3}{4}$ (d) $\frac{4}{3}$

17 The number of solutions of

$$\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2 + x + 1} = \frac{\pi}{2}$$

→ NCERT Exemplar

- (a) 0 (b) 1 (c) 2 (d) infinite

18 The maximum value of $(\sec^{-1} x)^2 + (\operatorname{cosec}^{-1} x)^2$ is

- (a) $\frac{\pi^2}{2}$ (b) $\frac{5\pi^2}{4}$
(c) π^2 (d) None of these

19 The trigonometric equation $\sin^{-1} x = 2 \sin^{-1} a$, has a solution for

→ AIEEE 2003

- (a) $\frac{1}{2} < |a| < \frac{1}{\sqrt{2}}$ (b) all real values of a
(c) $|a| \leq \frac{1}{\sqrt{2}}$ (d) $|a| \geq \frac{1}{\sqrt{2}}$

20 If $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$, then the value of

$$\tan^{-1} \left(\frac{\tan x}{4} \right) + \tan^{-1} \left(\frac{3 \sin 2x}{5 + 3 \cos 2x} \right)$$

- (a) $\frac{x}{2}$ (b) $2x$ (c) $3x$ (d) x

21 If $\tan \theta + \tan \left(\frac{\pi}{3} + \theta \right) + \tan \left(-\frac{\pi}{3} + \theta \right) = a \tan 3\theta$, then a

is equal to

- (a) $1/3$ (b) 1
(c) 3 (d) None of these

22 If $\cos^{-1} x = \tan^{-1} x$, then $\sin(\cos^{-1} x)$ is equal to

- (a) $-x$ (b) x^2 (c) x^3 (d) $-\frac{1}{x^2}$

23 The real solution of

$$\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2 + x + 1} = \frac{\pi}{2}$$

- (a) 2, 3 (b) 1, 0 (c) -1, 0 (d) 3, 1

24 If $\sin^{-1} \left(\frac{x}{5} \right) + \operatorname{cosec}^{-1} \left(\frac{5}{4} \right) = \frac{\pi}{2}$, then the value of x is

→ AIEEE 2007

- (a) 1 (b) 3 (c) 4 (d) 5

25 If $\operatorname{cosec}^{-1} x + \cos^{-1} y + \sec^{-1} z$

$$\geq \alpha^2 - \sqrt{2}\pi \alpha + 3\pi, \text{ where } \alpha \text{ is a real number,}$$

then

- (a) $x = 1, y = -1$ (b) $x = -1, z = -1$
(c) $x = 2, y = 1$ (d) $x = 1, y = -2$

26 The solution of $\sin^{-1} x \leq \cos^{-1} x$ is

- (a) $\left(-1, \frac{1}{\sqrt{2}} \right)$ (b) $\left[-1, \frac{1}{\sqrt{2}} \right]$
(c) $\left[1, \frac{1}{\sqrt{2}} \right]$ (d) $\left(1, \frac{1}{\sqrt{2}} \right)$

27 If m and M are the least and the greatest value of $(\cos^{-1} x)^2 + (\sin^{-1} x)^2$, then $\frac{M}{m}$ is equal to

- (a) 10 (b) 5 (c) 4 (d) 2

28 The number of real solutions of the equation

$$\sqrt{1 + \cos 2x} = \sqrt{2} \cos^{-1}(\cos x) \text{ in } \left[\frac{\pi}{2}, \pi \right] \text{ is}$$

→ NCERT Exemplar

- (a) 0 (b) 1 (c) 2 (d) infinite

29 The sum of the infinite series

$$\sin^{-1}\left(\frac{1}{\sqrt{2}}\right) + \sin^{-1}\left(\frac{\sqrt{2}-1}{\sqrt{6}}\right) + \sin^{-1}\left(\frac{\sqrt{3}-\sqrt{2}}{\sqrt{12}}\right) + \dots + \sin^{-1}\left(\frac{\sqrt{n}-\sqrt{(n-1)}}{\sqrt{n(n+1)}}\right) \text{ is}$$

(a) $\frac{\pi}{8}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{2}$ (d) π

30 A root of the equation

$$17x^2 + 17x \tan\left[2 \tan^{-1}\left(\frac{1}{5}\right) - \frac{\pi}{4}\right] - 10 = 0 \text{ is}$$

(a) $\frac{10}{17}$ (b) -1 (c) $-\frac{7}{17}$ (d) 1

31 The value of x for which $\sin[\cot^{-1}(1+x)] = \cos(\tan^{-1} x)$, is
 (a) $-\frac{1}{2}$ (b) 1 (c) 0 (d) $\frac{1}{2}$ → JEE Mains 2013

32 If $0 < x < 1$, then $\sqrt{1+x^2} [\{x \cos(\cot^{-1} x) + \sin(\cot^{-1} x)\}^2 - 1]^{1/2}$ is equal to
 (a) $\frac{x}{\sqrt{1+x^2}}$ (b) x (c) $x\sqrt{1+x^2}$ (d) $\sqrt{1+x^2}$

33 If x, y and z are in AP and $\tan^{-1} x, \tan^{-1} y$ and $\tan^{-1} z$ are also in AP, then
 (a) $x = y = z$ (b) $2x = y = 6z$
 (c) $6x = 3y = 2z$ (d) $6x = 4y = 3z$ → AIEEE 2012

DAY PRACTICE SESSION 2

PROGRESSIVE QUESTIONS EXERCISE

1 If $\sin^{-1}\left(x - \frac{x^2}{2} + \frac{x^3}{4} - \dots\right) + \cos^{-1}\left(x^2 - \frac{x^4}{2} + \frac{x^6}{4} - \dots\right) = \frac{\pi}{2}$, where $0 < |x| < \sqrt{2}$, then x is equal to
 (a) $\frac{1}{2}$ (b) 1 (c) $-\frac{1}{2}$ (d) -1

2 If the mapping $f(x) = ax + b, a > 0$ maps $[-1, 1]$ onto $[0, 2]$, then $\cot[\cot^{-1} 7 + \cot^{-1} 8 + \cot^{-1} 18]$ is equal to
 (a) $f(-1)$ (b) $f(0)$ (c) $f(1)$ (d) $f(2)$

3 If $S = \tan^{-1}\left(\frac{1}{n^2 + n + 1}\right) + \tan^{-1}\left(\frac{1}{n^2 + 3n + 3}\right) + \dots + \tan^{-1}\left\{\frac{1}{1 + (n+19)(n+20)}\right\}$, then $\tan S$ is equal to → JEE Mains 2013
 (a) $\frac{20}{401 + 20n}$ (b) $\frac{n}{n^2 + 20n + 1}$
 (c) $\frac{20}{n^2 + 20n + 1}$ (d) $\frac{n}{401 + 20n}$

4 If $f(x) = e^{\cos^{-1} \sin\left(x + \frac{\pi}{3}\right)}$, then
 (a) $f\left(-\frac{7\pi}{4}\right) = e^{\frac{\pi}{11}}$ (b) $f\left(\frac{8\pi}{9}\right) = e^{\frac{13\pi}{18}}$
 (c) $f\left(-\frac{7\pi}{4}\right) = e^{\frac{3\pi}{12}}$ (d) $f\left(-\frac{7\pi}{4}\right) = e^{\frac{11\pi}{13}}$

5 If $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$ and $f(1) = 2$,
 $f(p+q) = f(p) \cdot f(q), \forall p, q \in R$, then
 $x^{f(1)} + y^{f(2)} + z^{f(3)} - \frac{(x+y+z)}{x^{f(1)} + y^{f(2)} + z^{f(3)}}$ is equal to
 (a) 0 (b) 1
 (c) 2 (d) 3

6 If $[\cot^{-1} x] + [\cos^{-1} x] = 0$, where x is a non-negative real number and $[.]$ denotes the greatest integer function, then complete set of values of x is
 (a) $(\cos 1, 1]$ (b) $(\cot 1, 1]$
 (c) $(\cos 1, \cot 1)$ (d) None of these

7 $\cot^{-1}(\sqrt{\cos \alpha}) - \tan^{-1}(\sqrt{\cos \alpha}) = x$, then $\sin x$ is equal to → AIEEE 2002
 (a) $\tan^2\left(\frac{\alpha}{2}\right)$ (b) $\cot^2\left(\frac{\alpha}{2}\right)$ (c) $\tan \alpha$ (d) $\cot\left(\frac{\alpha}{2}\right)$

8 If $(\tan^{-1} x)^2 + (\cot^{-1} x)^2 = \frac{5\pi^2}{8}$, then the value of x is
 (a) -2 (b) -3 (c) -1 (d) 2

9 The solution set of $\tan^2(\sin^{-1} x) > 1$ is
 (a) $\left(-1, -\frac{1}{\sqrt{2}}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$ (b) $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) \sim \{0\}$
 (c) $(-1, 1) \sim \{0\}$ (d) None of these

10 If θ and ϕ are the roots of the equation $8x^2 + 22x + 5 = 0$, then
 (a) both $\sin^{-1} \theta$ and $\sin^{-1} \phi$ are equal
 (b) both $\sec^{-1} \theta$ and $\sec^{-1} \phi$ are real
 (c) both $\tan^{-1} \theta$ and $\tan^{-1} \phi$ are real
 (d) None of the above

11 $2 \tan^{-1}(-2)$ is equal to
 (a) $\cos^{-1}\left(\frac{-3}{5}\right)$ (b) $\pi + \cos^{-1} \frac{3}{5}$
 (c) $-\frac{\pi}{2} + \tan^{-1}\left(-\frac{3}{4}\right)$ (d) $-\pi + \cot^{-1}\left(-\frac{3}{4}\right)$

12 Let $x \in (0, 1)$. The set of all x such that $\sin^{-1} x > \cos^{-1} x$, is the interval → JEE Mains 2013
 (a) $\left(\frac{1}{2}, \frac{1}{\sqrt{2}}\right)$ (b) $\left(\frac{1}{\sqrt{2}}, 1\right)$
 (c) $(0, 1)$ (d) $\left(0, \frac{\sqrt{3}}{2}\right)$

ANSWERS

SESSION 1

1 (d)	2 (a)	3 (a)	4 (d)	5 (c)	6 (b)	7 (a)	8 (c)	9 (a)	10 (a)
11 (b)	12 (b)	13 (c)	14 (b)	15 (d)	16 (d)	17 (b)	18 (b)	19 (c)	20 (d)
21 (c)	22 (b)	23 (c)	24 (b)	25 (a)	26 (b)	27 (a)	28 (a)	29 (c)	30 (d)
31 (a)	32 (c)	33 (a)							

SESSION 2

1 (b)	2 (d)	3 (c)	4 (b)	5 (c)	6 (b)	7 (a)	8 (c)	9 (a)	10 (c)
11 (c)	12 (b)								

Hints and Explanations

SESSION 1

$$\begin{aligned}
 1 \quad \cos\left(\frac{33\pi}{5}\right) &= \cos\left(6\pi + \frac{3\pi}{5}\right) = \cos\frac{3\pi}{5} \\
 &= \sin\left(\frac{\pi}{2} - \frac{3\pi}{5}\right) = \sin\left(-\frac{\pi}{10}\right) \\
 \therefore \sin^{-1}\left(\cos\frac{33\pi}{5}\right) &= \sin^{-1}\left(\sin\left(-\frac{\pi}{10}\right)\right) = -\frac{\pi}{10}
 \end{aligned}$$

$$\begin{aligned}
 2 \quad \text{Since, } -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2} \\
 \therefore \sin^{-1} x_i = \frac{\pi}{2}, \quad 1 \leq i \leq 20 \\
 \Rightarrow x_i = 1, 1 \leq i \leq 20
 \end{aligned}$$

$$\text{Thus, } \sum_{i=1}^{20} x_i = 20$$

$$\begin{aligned}
 3 \quad \text{Given, } f(x) &= \sin^{-1} \sqrt{x-1} \\
 \text{For domain of } f(x) \quad &-1 \leq \sqrt{x-1} \leq 1 \\
 \Rightarrow 0 \leq (x-1) \leq 1 &\Rightarrow 1 \leq x \leq 2 \\
 \therefore x \in [1, 2]
 \end{aligned}$$

$$\begin{aligned}
 4 \quad \cos(2 \cos^{-1} x + \sin^{-1} x) &= \cos[2(\cos^{-1} x + \sin^{-1} x) - \sin^{-1} x] \\
 &= \cos(\pi - \sin^{-1} x) = -\cos(\sin^{-1} x) \\
 &= -\cos\left[\sin^{-1}\left(\frac{1}{5}\right)\right] \quad \left[\because x = \frac{1}{5}\right] \\
 &= -\cos\left[\cos^{-1}\sqrt{1 - \left(\frac{1}{5}\right)^2}\right] \\
 &= -\cos\left(\cos^{-1}\frac{2\sqrt{6}}{5}\right) = -\frac{2\sqrt{6}}{5}
 \end{aligned}$$

$$\begin{aligned}
 5 \quad \text{We have, } \cos[\tan^{-1}\{\sin(\cot^{-1} x)\}] \\
 \text{Let } \cot^{-1} x = \alpha \\
 \Rightarrow \cot \alpha = x \\
 \Rightarrow \operatorname{cosec} \alpha = \sqrt{1+x^2}
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow \sin \alpha &= \frac{1}{\sqrt{1+x^2}} \\
 \Rightarrow \alpha &= \sin^{-1} \frac{1}{\sqrt{1+x^2}} \\
 \text{Hence, } \cos[\tan^{-1}\{\sin(\cot^{-1} x)\}] &= \cos\left[\tan^{-1}\left\{\sin\left(\sin^{-1} \frac{1}{\sqrt{1+x^2}}\right)\right\}\right] \\
 &= \cos\left[\tan^{-1} \frac{1}{\sqrt{1+x^2}}\right] \\
 &= \cos\left[\cos^{-1} \frac{\sqrt{x^2+1}}{\sqrt{x^2+2}}\right] \\
 &= \left[\begin{aligned} &\because \text{let } \tan^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right) = \beta \\ &\tan \beta = \frac{1}{\sqrt{1+x^2}} \\ &\sec \beta = \sqrt{1 + \frac{1}{1+x^2}} = \sqrt{\frac{x^2+2}{x^2+1}} \\ &\cos \beta = \frac{\sqrt{x^2+1}}{\sqrt{x^2+2}} \end{aligned} \right] \\
 &= \frac{\sqrt{x^2+1}}{\sqrt{x^2+2}}
 \end{aligned}$$

$$\begin{aligned}
 6 \quad \text{Given, } \tan^{-1} x - \cot^{-1} x &= \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) \\
 \Rightarrow \tan^{-1} x - \cot^{-1} x &= \frac{\pi}{6} \quad \dots(i) \\
 \text{But } \tan^{-1} x + \cot^{-1} x &= \frac{\pi}{2} \quad \dots(ii) \\
 \text{On adding Eqs. (i) and (ii), we get} \\
 2 \tan^{-1} x &= \frac{\pi}{6} + \frac{\pi}{2} = \frac{2\pi}{3} \\
 \Rightarrow \tan^{-1} x &= \frac{\pi}{3} \\
 \Rightarrow x &= \tan \frac{\pi}{3} \Rightarrow x = \sqrt{3} \\
 \text{It has unique solution.}
 \end{aligned}$$

7 Given,

$$\tan^{-1} y = \tan^{-1} x + \tan^{-1}\left(\frac{2x}{1-x^2}\right),$$

$$\text{where } |x| < \frac{1}{\sqrt{3}}$$

$$\begin{aligned}
 \Rightarrow \tan^{-1} y &= \tan^{-1} \left[\frac{x + \frac{2x}{1-x^2}}{1 - x\left(\frac{2x}{1-x^2}\right)} \right] \\
 \left[\because \tan^{-1} x + \tan^{-1} y &= \tan^{-1} \left(\frac{x+y}{1-xy} \right), \right. \\
 &\quad \left. xy < 1 \right]
 \end{aligned}$$

$$= \tan^{-1} \left(\frac{x - x^3 + 2x}{1 - x^2 - 2x^2} \right)$$

$$\tan^{-1} y = \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right)$$

$$\Rightarrow y = \frac{3x - x^3}{1 - 3x^2}$$

Alternate Method

$$|x| < \frac{1}{\sqrt{3}} \Rightarrow -\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$$

$$\text{Let } x = \tan \theta \Rightarrow -\frac{\pi}{6} < \theta < \frac{\pi}{6}$$

$$\therefore \tan^{-1} y = \theta + \tan^{-1}(\tan 2\theta) = \theta + 2\theta = 3\theta$$

$$\Rightarrow y = \tan 3\theta$$

$$\Rightarrow y = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}$$

$$\Rightarrow y = \frac{3x - x^3}{1 - 3x^2}$$

8 Given that, $\theta = \tan^{-1} a$

$$\text{and } \phi = \tan^{-1} b$$

$$\text{and } ab = -1$$

$$\therefore \tan \theta \tan \phi = ab = -1$$

$$\Rightarrow \tan \theta = -\cot \phi$$

$$\Rightarrow \tan \theta = \tan \left(\frac{\pi}{2} + \phi \right)$$

$$\Rightarrow \theta - \phi = \frac{\pi}{2}$$

9 $f(x) = |3 \tan^{-1} x - \cos^{-1}(0)| - \cos^{-1}(-1)$

$$= \left| 3 \tan^{-1} x - \left(\frac{\pi}{2} \right) \right| - \pi$$

We know that, $-\frac{\pi}{2} < \tan^{-1} x < \frac{\pi}{2}$

$$\Rightarrow \frac{-3\pi}{2} < 3 \tan^{-1} x < \frac{3\pi}{2}$$

$$\Rightarrow -2\pi < 3 \tan^{-1} x - \frac{\pi}{2} < \pi$$

$$\Rightarrow 0 \leq \left| 3 \tan^{-1} x - \frac{\pi}{2} \right| < 2\pi$$

$$\Rightarrow -\pi \leq \left| 3 \tan^{-1} x - \frac{\pi}{2} \right| - \pi < \pi$$

10 $\cos(\cos^{-1} x) = x, \forall x \in [-1, 1]$ and

$$\operatorname{cosec}(\operatorname{cosec}^{-1} x)$$

$$= x, \forall x \in (-\infty, -1] \cup [1, \infty)$$

$$\Rightarrow \cos(\cos^{-1} x) = \operatorname{cosec}(\operatorname{cosec}^{-1} x) \text{ for } x = \pm 1 \text{ only.}$$

Hence, there are two roots.

11 Since, $\operatorname{cosec}^{-1} \left(\frac{5}{3} \right) = \tan^{-1} \left(\frac{3}{4} \right)$

$$\therefore \cot \left(\tan^{-1} \frac{3}{4} + \tan^{-1} \frac{2}{3} \right)$$

$$= \cot \left(\tan^{-1} \left[\frac{\frac{3}{4} + \frac{2}{3}}{1 - \frac{1}{2}} \right] \right)$$

$$= \cot \left(\tan^{-1} \left[\frac{\left(\frac{17}{12} \right)}{\left(\frac{1}{2} \right)} \right] \right)$$

$$= \cot \left[\tan^{-1} \left(\frac{17}{6} \right) \right] = \frac{6}{17}$$

12 Let $\cot^{-1} \frac{1}{2} = \phi \Rightarrow \frac{1}{2} = \cot \phi$

$$\Rightarrow \sin \phi = \frac{1}{\sqrt{1 + \cot^2 \phi}} = \frac{2}{\sqrt{5}}$$

$$\text{Let } \cos^{-1} x = \theta \Rightarrow \sec \theta = \frac{1}{x}$$

$$\Rightarrow \tan \theta = \sqrt{\sec^2 \theta - 1}$$

$$\Rightarrow \tan \theta = \sqrt{\frac{1}{x^2} - 1} \Rightarrow \tan \theta = \frac{\sqrt{1 - x^2}}{x}$$

$$\text{Now, } \tan(\cos^{-1} x) = \sin \left(\cot^{-1} \frac{1}{2} \right)$$

$$\Rightarrow \tan \left(\tan^{-1} \frac{\sqrt{1 - x^2}}{x} \right) = \sin \left(\sin^{-1} \frac{2}{\sqrt{5}} \right)$$

$$\Rightarrow \frac{\sqrt{1 - x^2}}{x} = \frac{2}{\sqrt{5}} \Rightarrow \sqrt{(1 - x^2)5} = 2x$$

On squaring both sides, we get

$$(1 - x^2)5 = 4x^2$$

$$\Rightarrow 9x^2 = 5 \Rightarrow x = \pm \frac{\sqrt{5}}{3}$$

13 $\therefore \tan^{-1} x - \tan^{-1} y = 0 \Rightarrow x = y$

$$\text{Also, } \cos^{-1} x + \cos^{-1} y = \frac{\pi}{2}$$

$$\Rightarrow 2 \cos^{-1} x = \frac{\pi}{2} \Rightarrow \cos^{-1} x = \frac{\pi}{4}$$

$$\Rightarrow x = \frac{1}{\sqrt{2}} \Rightarrow x^2 = \frac{1}{2}$$

$$\text{Hence, } x^2 + xy + y^2 = 3x^2 = \frac{3}{2}$$

14 $\therefore \sin^{-1} \left(\frac{x}{5} \right) + \operatorname{cosec}^{-1} \left(\frac{5}{4} \right) = \frac{\pi}{2}$

$$\Rightarrow \sin^{-1} \left(\frac{x}{5} \right) + \sin^{-1} \left(\frac{4}{5} \right) = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1} \left(\frac{x}{5} \right) = \frac{\pi}{2} - \sin^{-1} \left(\frac{4}{5} \right)$$

$$\Rightarrow \sin^{-1} \left(\frac{x}{5} \right) = \cos^{-1} \left(\frac{4}{5} \right)$$

$$\Rightarrow \sin^{-1} \left(\frac{x}{5} \right) = \sin^{-1} \left(\frac{3}{5} \right)$$

$$\therefore x = 3$$

15 Given, $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$

$$\Rightarrow \sin^{-1} x = \sin^{-1} y = \sin^{-1} z = \frac{\pi}{2}$$

$$\Rightarrow x = y = z = 1$$

$$\therefore \Sigma \frac{(x^{101} + y^{101})(x^{202} + y^{202})}{(x^{303} + y^{303})(x^{404} + y^{404})}$$

$$= \Sigma \frac{(1+1)(1+1)}{(1+1)(1+1)} = \Sigma 1 = 3$$

16 $\tan^{-1} \left[\frac{\frac{x-1}{x+1} + \frac{2x-1}{2x+1}}{1 - \left(\frac{x-1}{x+1} \right) \left(\frac{2x-1}{2x+1} \right)} \right]$

$$= \tan^{-1} \left(\frac{23}{36} \right)$$

$$\Rightarrow \frac{2x^2 - 1}{3x} = \frac{23}{36}$$

$$\Rightarrow 24x^2 - 12 - 23x = 0$$

$$\Rightarrow x = \frac{4}{3}, -\frac{3}{8}$$

But x cannot be negative.

$$\therefore x = \frac{4}{3}$$

17 For existence, $x(x+1) \geq 0$... (i)

$$\text{and } x^2 + x + 1 \leq 1$$

$$\Rightarrow x^2 + x \leq 0$$

$$\Rightarrow x(x+1) \leq 0$$
 ... (ii)

From Eqs. (i) and (ii), we get

$$x(x+1) = 0 \Rightarrow x = 0, -1$$

But $x = -1$ is not satisfied the given equation.

18 Let $I = (\sec^{-1} x)^2 + (\operatorname{cosec}^{-1} x)^2$

$$= (\sec^{-1} x + \operatorname{cosec}^{-1} x)^2$$

$$- 2 \sec^{-1} x \operatorname{cosec}^{-1} x$$

$$= \frac{\pi^2}{4} - 2 \sec^{-1} x \left(\frac{\pi}{2} - \sec^{-1} x \right)$$

$$= \frac{\pi^2}{4} + 2 \left[(\sec^{-1} x)^2 - \frac{\pi}{2} (\sec^{-1} x) + \frac{\pi^2}{16} - \frac{\pi^2}{16} \right]$$

$$= \frac{\pi^2}{8} + 2 \left[\left(\sec^{-1} x - \frac{\pi}{4} \right)^2 \right]$$

$$\therefore I_{\max} = \frac{\pi^2}{8} + 2 \left[\frac{9\pi^2}{16} \right] = \frac{5\pi^2}{4}$$

19 Given that, $\sin^{-1} x = 2 \sin^{-1} a$

$$\therefore -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{2} \leq 2 \sin^{-1} a \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{4} \leq \sin^{-1} a \leq \frac{\pi}{4}$$

$$\Rightarrow \sin \left(-\frac{\pi}{4} \right) \leq a \leq \sin \frac{\pi}{4}$$

$$\Rightarrow -\frac{1}{\sqrt{2}} \leq a \leq \frac{1}{\sqrt{2}}$$

$$\Rightarrow |a| \leq \frac{1}{\sqrt{2}}$$

20 $\tan^{-1} \left(\frac{\tan x}{4} \right) + \tan^{-1} \left(\frac{3 \sin 2x}{5 + 3 \cos 2x} \right)$

$$= \tan^{-1} \left(\frac{\tan x}{4} \right) + \tan^{-1}$$

$$\left(\frac{\frac{6 \tan x}{1 + \tan^2 x}}{5 + \frac{3(1 - \tan^2 x)}{1 + \tan^2 x}} \right)$$

$$= \tan^{-1} \left(\frac{\tan x}{4} \right) + \tan^{-1} \left(\frac{6 \tan x}{8 + 2 \tan^2 x} \right)$$

$$= \tan^{-1} \left(\frac{\tan x}{4} \right) + \tan^{-1} \left(\frac{3 \tan x}{4 + \tan^2 x} \right)$$

$$= \tan^{-1} \left(\frac{\frac{\tan x}{4} + \frac{3 \tan x}{4 + \tan^2 x}}{1 - \frac{3 \tan^2 x}{4(4 + \tan^2 x)}} \right)$$

$$\left(\text{as } \left| \frac{\tan x}{4} \cdot \frac{3 \tan x}{4 \tan^2 x} \right| < 1 \right)$$

$$= \tan^{-1} \left(\frac{16 \tan x + \tan^3 x}{16 + \tan^2 x} \right)$$

$$= \tan^{-1}(\tan x) = x$$

$$\begin{aligned}
21 \quad & \tan \theta + \tan \left(\frac{\pi}{3} + \theta \right) + \tan \left(-\frac{\pi}{3} + \theta \right) \\
&= a \tan 3\theta \\
\Rightarrow & \tan \theta + \frac{\sqrt{3} + \tan \theta}{1 - \sqrt{3} \tan \theta} \\
&+ \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta} = a \tan 3\theta \\
\Rightarrow & \tan \theta + \frac{8 \tan \theta}{1 - 3 \tan^2 \theta} = a \tan 3\theta \\
\Rightarrow & \frac{3(3 \tan \theta - \tan^3 \theta)}{1 - 3 \tan^2 \theta} = a \tan 3\theta \\
\Rightarrow & 3 \tan 3\theta = a \tan 3\theta \\
\Rightarrow & a = 3
\end{aligned}$$

$$\begin{aligned}
22 \quad & \text{Let } \cos^{-1} x = \tan^{-1} x = \theta \\
\Rightarrow & x = \cos \theta = \tan \theta \\
\Rightarrow & \cos \theta = \tan \theta \Rightarrow \cos \theta = \frac{\sin \theta}{\cos \theta} \\
\Rightarrow & \cos^2 \theta = \sin \theta \\
\Rightarrow & \sin^2 \theta + \sin \theta - 1 = 0 \\
\Rightarrow & \sin \theta = \frac{-1 \pm \sqrt{1+4}}{2} \\
\Rightarrow & \sin \theta = \frac{\sqrt{5}-1}{2} \\
\therefore & x^2 = \cos^2 \theta = \sin \theta = \frac{\sqrt{5}-1}{2} \\
& \text{and } \sin(\cos^{-1} x) = \sin \theta \\
&= \frac{\sqrt{5}-1}{2} = x^2
\end{aligned}$$

$$\begin{aligned}
23 \quad & \text{Given, } \tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \\
& \sqrt{x^2 + x + 1} = \frac{\pi}{2} \\
\Rightarrow & \cos^{-1} \frac{1}{\sqrt{1+(x^2+x)}} \\
& + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{2} \\
\Rightarrow & \cos^{-1} \frac{1}{\sqrt{1+(x^2+x)}} = \frac{\pi}{2} \\
& - \sin^{-1} \sqrt{x^2+x+1} \\
\Rightarrow & \cos^{-1} \frac{1}{\sqrt{x^2+x+1}} \\
& = \cos^{-1} \sqrt{x^2+x+1} \\
\Rightarrow & \frac{1}{\sqrt{x^2+x+1}} = \sqrt{x^2+x+1} \\
\Rightarrow & x^2+x+1=1 \Rightarrow x=-1, 0
\end{aligned}$$

$$\begin{aligned}
24 \quad & \text{Since, } \sin^{-1} \left(\frac{x}{5} \right) + \operatorname{cosec}^{-1} \left(\frac{5}{4} \right) = \frac{\pi}{2} \\
\Rightarrow & \sin^{-1} \left(\frac{x}{5} \right) + \sin^{-1} \left(\frac{4}{5} \right) = \frac{\pi}{2} \\
\Rightarrow & \sin^{-1} \left(\frac{x}{5} \right) = \frac{\pi}{2} - \sin^{-1} \left(\frac{4}{5} \right)
\end{aligned}$$

$$\begin{aligned}
\Rightarrow & \sin^{-1} \left(\frac{x}{5} \right) = \cos^{-1} \left(\frac{4}{5} \right) \\
\Rightarrow & \sin^{-1} \left(\frac{x}{5} \right) = \sin^{-1} \left(\frac{3}{5} \right) \\
\therefore & x = 3
\end{aligned}$$

$$\begin{aligned}
25 \quad & \text{Given,} \\
& \operatorname{cosec}^{-1} x + \cos^{-1} y + \sec^{-1} z \geq \alpha^2 \\
& \quad \quad \quad - \sqrt{2\pi} \alpha + 3\pi \\
\text{RHS} &= \alpha^2 - \sqrt{2\pi} \alpha + 3\pi \\
&= \alpha^2 - 2\sqrt{\frac{\pi}{2}} \alpha + \frac{\pi}{2} + 3\pi - \frac{\pi}{2} \\
&= \left(\alpha - \sqrt{\frac{\pi}{2}} \right)^2 + \frac{5\pi}{2} \geq \frac{5\pi}{2} \quad \dots(i) \\
\therefore \text{LHS, } & \operatorname{cosec}^{-1} x \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] - \{0\} \\
& \cos^{-1} y \in [0, \pi] \\
& \text{and } \sec^{-1} z \in [0, \pi] - \left\{ \frac{\pi}{2} \right\} \\
\therefore & \text{LHS} \leq \frac{5\pi}{2} \quad \dots(ii) \\
& \text{From Eqs. (i) and (ii), we get only} \\
& \text{possibility is sign of equality} \\
& x = 1, y = -1, z = -1
\end{aligned}$$

$$\begin{aligned}
26 \quad & \text{Given, } \cos^{-1} x \geq \sin^{-1} x \\
\Rightarrow & \frac{\pi}{2} \geq 2 \sin^{-1} x \\
& \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}, \forall x \in [-1, 1] \right] \\
\Rightarrow & \sin^{-1} x \leq \frac{\pi}{4} \Rightarrow -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2} \\
& \left(\because \text{range of } \sin^{-1} x \text{ is } \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \right) \\
\Rightarrow & -1 \leq x \leq \sin \left(\frac{\pi}{4} \right) \\
\Rightarrow & x \in \left[-1, \frac{1}{\sqrt{2}} \right]
\end{aligned}$$

$$\begin{aligned}
27 \quad & \left(\frac{\pi}{2} - \sin^{-1} x \right)^2 + (\sin^{-1} x)^2 \\
&= \frac{\pi^2}{4} + 2(\sin^{-1} x)^2 - \pi \sin^{-1} x \\
&= \frac{\pi^2}{8} + 2 \left[\sin^{-1} x - \frac{\pi}{4} \right]^2 \\
&\text{Here, } m = \frac{\pi^2}{8}, M = \frac{5\pi^2}{4} \\
\therefore & \frac{M}{m} = 10
\end{aligned}$$

$$\begin{aligned}
28 \quad & \text{Given, } \sqrt{1+\cos 2x} = \sqrt{2} \cos^{-1}(\cos x) \\
\therefore & \sqrt{2 \cos^2 x} = \sqrt{2} x \\
\Rightarrow & \sqrt{2} |\cos x| = \sqrt{2} x \\
& \text{For } x \in \left[\frac{\pi}{2}, \pi \right], |\cos x| = -\cos x \\
& -\sqrt{2} \cos x = \sqrt{2} x \\
\Rightarrow & -\cos x = x
\end{aligned}$$

$$\begin{aligned}
\therefore & \cos x = -x \\
& \text{Hence, no solution exist.}
\end{aligned}$$

$$\begin{aligned}
29 \quad & \because T_r = \sin^{-1} \left\{ \frac{\sqrt{r} - \sqrt{(r-1)}}{\sqrt{r(r+1)}} \right\} \\
&= \tan^{-1} \left\{ \frac{\sqrt{r} - \sqrt{(r-1)}}{1 + \sqrt{r}(\sqrt{r-1})} \right\} \\
\therefore S_n &= \sum_{r=1}^n \tan^{-1} \left(\frac{\sqrt{r} - \sqrt{r-1}}{1 + \sqrt{r} \sqrt{r-1}} \right) \\
&= \sum_{r=1}^n [\tan^{-1} \sqrt{r} - \tan^{-1} \sqrt{(r-1)}] \\
&= \tan^{-1} \sqrt{n} - \tan^{-1} \sqrt{0} \\
&= \tan^{-1} \sqrt{n} - 0 \\
\therefore S_\infty &= \tan^{-1} \infty = \frac{\pi}{2}
\end{aligned}$$

$$30 \quad \text{Now, } \tan \left\{ 2 \tan^{-1} \left(\frac{1}{5} \right) \right\} = \frac{2 \times \frac{1}{5}}{1 - \frac{1}{25}} = \frac{5}{12}$$

Given equation can be rewritten as

$$\begin{aligned}
17x^2 - 17x \tan \left\{ \frac{\pi}{4} - 2 \tan^{-1} \left(\frac{1}{5} \right) \right\} - 10 &= 0 \\
\Rightarrow 17x^2 - 17x \cdot \frac{1 - \frac{5}{12}}{1 + \frac{5}{12}} - 10 &= 0 \\
\Rightarrow 17x^2 - 7x - 10 &= 0 \\
\Rightarrow (x-1)(17x+10) &= 0 \\
&\text{Hence, } x=1 \text{ is a root of the given} \\
&\text{equation.}
\end{aligned}$$

$$\begin{aligned}
31 \quad & \sin \left(\sin^{-1} \frac{1}{\sqrt{(1+x^2)+1}} \right) \\
&= \cos \left(\cos^{-1} \frac{1}{\sqrt{1+x^2}} \right) \\
&= \frac{1}{\sqrt{(1+x^2)+1}} = \frac{1}{\sqrt{1+x^2}} \\
\Rightarrow (1+x^2)+1 &= 1+x^2 \\
\Rightarrow 2x+1 &= 0 \\
\Rightarrow x &= -\frac{1}{2}
\end{aligned}$$

32 We have, $0 < x < 1$

$$\begin{aligned}
& \text{Now, } \sqrt{1+x^2} [\{x \cos(\cot^{-1} x) \\
& \quad + \sin(\cot^{-1} x)\}^2 - 1]^{1/2} \\
&= \sqrt{1+x^2} \\
& \left[\left\{ x \frac{x}{\sqrt{1+x^2}} + \frac{1}{\sqrt{1+x^2}} \right\}^2 - 1 \right]^{1/2} \\
&= \sqrt{1+x^2} \left[\left(\frac{1+x^2}{\sqrt{1+x^2}} \right)^2 - 1 \right]^{1/2} \\
&= \sqrt{1+x^2} [1+x^2-1]^{1/2} \\
&= x \sqrt{1+x^2}
\end{aligned}$$

33 Since, x, y and z are in AP.

$$\therefore 2y = x + z$$

Also, $\tan^{-1} x, \tan^{-1} y$ and $\tan^{-1} z$ are in AP.

$$\therefore 2\tan^{-1} y = \tan^{-1} x + \tan^{-1} z$$

$$\Rightarrow \tan^{-1} \left(\frac{2y}{1-y^2} \right) = \tan^{-1} \left(\frac{x+z}{1-xz} \right)$$

$$\Rightarrow \frac{x+z}{1-y^2} = \frac{x+z}{1-xz}$$

$$\Rightarrow y^2 = xz \quad [\because 2y = x+z]$$

Since x, y and z are in AP as well as in GP.

$$\therefore x = y = z$$

SESSION 2

$$\begin{aligned} \text{1 Now, } x - \frac{x^2}{2} + \frac{x^3}{4} - \dots \\ = \frac{x}{1 + \frac{x}{2}} = \frac{2x}{2+x} \end{aligned}$$

$$\begin{aligned} \text{and } x^2 - \frac{x^4}{2} + \frac{x^6}{4} - \dots \\ = \frac{x^2}{1 + \frac{x^2}{2}} = \frac{2x^2}{2+x^2} \end{aligned}$$

$$\therefore \sin^{-1} \alpha + \cos^{-1} \alpha = \frac{\pi}{2}$$

$$\therefore \frac{2x}{2+x} = \frac{2x^2}{2+x^2}, x \neq 0$$

$$\Rightarrow 2 + x^2 = 2x + x^2$$

$$\therefore x = 1$$

$$\text{2 } \therefore f(x) = ax + b$$

$$\therefore f'(x) = a > 0$$

So, $f(x)$ is an increasing function.

$$\Rightarrow f(-1) = 0 \text{ and } f(1) = 2$$

$$\Rightarrow -a + b = 0$$

$$\text{and } a + b = 2$$

$$\text{Then, } a = b = 1$$

$$\therefore f(x) = x + 1$$

$$\text{Now, } \cot[\cot^{-1} 7 + \cot^{-1} 8 + \cot^{-1} 18]$$

$$= \cot \left[\tan^{-1} \left(\frac{1}{7} \right) + \tan^{-1} \left(\frac{1}{8} \right) + \tan^{-1} \left(\frac{1}{18} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{\frac{1}{7} + \frac{1}{8}}{1 - \frac{1}{7} \cdot \frac{1}{8}} \right) + \tan^{-1} \left(\frac{1}{18} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{15}{55} \right) + \tan^{-1} \left(\frac{1}{18} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{3}{11} \right) + \tan^{-1} \left(\frac{1}{18} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{\frac{3}{11} + \frac{1}{18}}{1 - \frac{3}{11} \cdot \frac{1}{18}} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{65}{195} \right) \right]$$

$$= \cot \left[\tan^{-1} \left(\frac{1}{3} \right) \right] = \cot(\cot^{-1} 3) = 3$$

$$= 1 + 2 = f(2) \quad [\because f(x) = x + 1]$$

$$\text{3 } S = \tan^{-1} \left\{ \frac{(n+1) - (n+0)}{1 + (n+0)(n+1)} \right\}$$

$$+ \tan^{-1} \left\{ \frac{(n+2) - (n+1)}{1 + (n+1)(n+2)} \right\}$$

$$+ \dots + \tan^{-1} \left\{ \frac{(n+20) - (n+19)}{1 + (n+19)(n+20)} \right\}$$

$$= \tan^{-1}(n+1) - \tan^{-1} n$$

$$+ \tan^{-1}(n+2) - \tan^{-1}(n+1)$$

$$+ \dots + \tan^{-1}(n+20) - \tan^{-1}(n+19)$$

$$= \tan^{-1}(n+20) - \tan^{-1} n$$

$$= \tan^{-1} \left\{ \frac{n+20-n}{1+n(n+20)} \right\}$$

$$= \tan^{-1} \left(\frac{20}{n^2 + 20n + 1} \right)$$

$$\Rightarrow \tan S = \tan \left\{ \tan^{-1} \left(\frac{20}{n^2 + 20n + 1} \right) \right\}$$

$$\Rightarrow \tan S = \frac{20}{n^2 + 20n + 1}$$

$$\text{4 Given, } f(x) = e^{\cos^{-1} \sin \left(x + \frac{\pi}{3} \right)}$$

$$\Rightarrow f\left(\frac{8\pi}{9}\right) = e^{\cos^{-1} \sin \left(\frac{8\pi}{9} + \frac{\pi}{3} \right)}$$

$$= e^{\cos^{-1} \sin \left(\frac{11\pi}{9} \right)}$$

$$\Rightarrow f\left(\frac{8\pi}{9}\right) = e^{\cos^{-1} \cos \left(\frac{13\pi}{18} \right)} = e^{\frac{13\pi}{18}}$$

$$\text{Also, } f\left(-\frac{7\pi}{4}\right) = e^{\cos^{-1} \sin \left(-\frac{7\pi}{4} + \frac{\pi}{3} \right)}$$

$$= e^{\cos^{-1} \sin \left(-\frac{17\pi}{12} \right)} = e^{\cos^{-1} \cos \frac{\pi}{12}} = e^{\frac{\pi}{12}}$$

$$\text{5 } \therefore -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}, -\frac{\pi}{2} \leq \sin^{-1} y \leq \frac{\pi}{2}$$

$$\text{and } -\frac{\pi}{2} \leq \sin^{-1} z \leq \frac{\pi}{2}$$

Given that,

$$\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$$

which is possible only when

$$\sin^{-1} x = \sin^{-1} y = \sin^{-1} z = \frac{\pi}{2}$$

$$\Rightarrow x = y = z = 1$$

$$\text{Put } p = q = 1$$

$$\text{Then, } f(2) = f(1)f(1) = 2 \cdot 2 = 4$$

$$\text{and put } p = 1, q = 2$$

$$\text{Then, } f(3) = f(1)f(2) = 2 \cdot 2 = 8$$

$$\therefore x^{f(1)} + y^{f(2)} + z^{f(3)} = \frac{x+y+z}{x^{f(1)} + y^{f(2)} + z^{f(3)}}$$

$$= 1 + 1 + 1 - \frac{3}{1+1+1} = 3 - 1 = 2$$

$$\text{6 } \therefore 0 \leq \cos^{-1} x \leq \pi$$

$$\text{and } 0 < \cot^{-1} x < \pi$$

$$\text{Given, } [\cot^{-1} x] + [\cos^{-1} x] = 0$$

$$\Rightarrow [\cot^{-1} x] = 0 \text{ and } [\cos^{-1} x] = 0$$

$$\Rightarrow 0 < \cot^{-1} x < 1 \text{ and } 0 \leq \cos^{-1} x < 1$$

$$\therefore x \in (\cot 1, \infty)$$

$$\text{and } x \in (\cos 1, 1) \Rightarrow x \in (\cot 1, 1)$$

7 Given that,

$$\cot^{-1}(\sqrt{\cos \alpha}) - \tan^{-1}(\sqrt{\cos \alpha}) = x \quad \dots(i)$$

We know that,

$$\begin{aligned} \cot^{-1}(\sqrt{\cos \alpha}) + \tan^{-1}(\sqrt{\cos \alpha}) &= \frac{\pi}{2} \quad 3 \\ \dots(ii) \\ \left[\because \cot^{-1} x + \tan^{-1} x &= \frac{\pi}{2} \right] \end{aligned}$$

On adding Eqs. (i) and (ii), we get

$$2\cot^{-1}(\sqrt{\cos \alpha}) = \frac{\pi}{2} + x$$

$$\Rightarrow \sqrt{\cos \alpha} = \cot \left(\frac{\pi}{4} + \frac{x}{2} \right)$$

$$\Rightarrow \sqrt{\cos \alpha} = \frac{\cot \frac{x}{2} - 1}{1 + \cot \frac{x}{2}}$$

$$\Rightarrow \sqrt{\cos \alpha} = \frac{\cos \frac{x}{2} - \sin \frac{x}{2}}{\cos \frac{x}{2} + \sin \frac{x}{2}}$$

On squaring both sides, we get

$$\begin{aligned} \Rightarrow \cos \alpha &= \frac{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2}} \\ &\quad - \frac{2\sin \frac{x}{2} \cos \frac{x}{2}}{2} \\ &\quad + \frac{2\sin \frac{x}{2} \cos \frac{x}{2}}{2} \end{aligned}$$

$$\Rightarrow \cos \alpha = \frac{1 - \sin x}{1 + \sin x}$$

$$\Rightarrow \frac{1 - \tan^2 \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}} = \frac{1 - \sin x}{1 + \sin x}$$

On applying componendo and dividendo rule, we get

$$\sin x = \tan^2 \left(\frac{\alpha}{2} \right)$$

8 Given,

$$\begin{aligned}(\tan^{-1} x)^2 + (\cot^{-1} x)^2 &= \frac{5\pi^2}{8} \\ \Rightarrow (\tan^{-1} x + \cot^{-1} x)^2 - 2 \tan^{-1} x \cdot \cot^{-1} x &= \frac{5\pi^2}{8}\end{aligned}$$

$$\begin{aligned}\Rightarrow \left(\frac{\pi}{2}\right)^2 - 2 \tan^{-1} x \left(\frac{\pi}{2} - \tan^{-1} x\right) &= \frac{5\pi^2}{8} \left[\because \tan^{-1} x + \cot^{-1} x = \frac{\pi}{2} \right] \\ \Rightarrow \frac{\pi^2}{4} - 2 \cdot \frac{\pi}{2} \tan^{-1} x &+ 2 (\tan^{-1} x)^2 = \frac{5\pi^2}{8}\end{aligned}$$

$$\Rightarrow 2 (\tan^{-1} x)^2 - \pi \tan^{-1} x - \frac{3\pi^2}{8} = 0$$

$$\Rightarrow \tan^{-1} x = -\frac{\pi}{4}, \frac{3\pi}{4}$$

$$\Rightarrow \tan^{-1} x = -\frac{\pi}{4}$$

$$\Rightarrow x = -1$$

$$\left\{ \begin{array}{l} \text{neglecting } \tan^{-1} x \\ = \frac{3\pi}{4} \text{ as principal value of} \\ \tan^{-1} x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \end{array} \right\}$$

9 $\tan^2 (\sin^{-1} x) > 1$

$$\Rightarrow \frac{\pi}{4} < \sin^{-1} x < \frac{\pi}{2}$$

$$\text{or } -\frac{\pi}{2} < \sin^{-1} x < -\frac{\pi}{4}$$

$$\Rightarrow x \in \left(\frac{1}{\sqrt{2}}, 1\right) \text{ or } x \in \left(-1, -\frac{1}{\sqrt{2}}\right)$$

$$\Rightarrow x \in \left(-1, -\frac{1}{\sqrt{2}}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$$

10 $8x^2 + 22x + 5 = 0$

$$\Rightarrow x = -\frac{1}{4}, -\frac{5}{2}$$

$$\therefore -1 < -\frac{1}{4} < 1 \text{ and } -\frac{5}{2} < -1$$

$$\therefore \sin^{-1} \left(-\frac{1}{4}\right) \text{ exists but } \sin^{-1} \left(-\frac{5}{2}\right)$$

does not exist.

$$\sec^{-1} \left(-\frac{5}{2}\right) \text{ exists but } \sec^{-1} \left(-\frac{1}{4}\right) \text{ does}$$

not exist.

$$\text{So, } \tan^{-1} \left(-\frac{1}{4}\right)$$

$$\text{and } \tan^{-1} \left(-\frac{5}{2}\right) \text{ both exist.}$$

11 Let $\tan^{-1}(-2) = \theta$

$$\Rightarrow \tan \theta = -2$$

$$\Rightarrow \theta \in \left(-\frac{\pi}{2}, 0\right)$$

$$\Rightarrow 2\theta \in (-\pi, 0)$$

$$\text{Now, } \cos(-2\theta) = \cos 2\theta$$

$$= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

$$= \frac{-3}{5}$$

$$\Rightarrow -2\theta = \cos^{-1} \left(\frac{-3}{5}\right)$$

$$= \pi - \cos^{-1} \frac{3}{5}$$

$$\Rightarrow 2\theta = -\pi + \cos^{-1} \frac{3}{5}$$

$$\Rightarrow 2\theta = -\pi + \tan^{-1} \frac{4}{3}$$


$$= -\pi + \cot^{-1} \frac{3}{4}$$

$$= -\pi + \frac{\pi}{2} - \tan^{-1} \frac{3}{4}$$

$$= -\frac{\pi}{2} - \tan^{-1} \frac{3}{4}$$

$$= -\frac{\pi}{2} + \tan^{-1} \left(-\frac{3}{4}\right)$$

12



$$\therefore \sin^{-1} x > \cos^{-1} x, \forall x \in \left(\frac{1}{\sqrt{2}}, 1\right)$$