

Question Paper contains 20 printed pages.
A & Part - B)

3. 0100754

050 (E)
(MARCH, 2020)
SCIENCE STREAM
(CLASS - XII)
(New Course)

A : Time : 1 Hour / Marks : 50
B : Time : 2 Hours / Marks : 50

પ્રશ્ન પેપરનો સેટ નંબર જેની
સામેનું વર્તુળ OMR શીટમાં
ઘટ્ટ કરવાનું રહે છે.

Set No. of Question Paper,
circle against which is to be
darken in OMR sheet.

01

(Part - A)

: 1 Hour]

[Maximum Marks : 50

Instructions :

- 1) There are 50 objective type (M.C.Q.) questions in Part - A and all questions are compulsory.
- 2) The questions are serially numbered from 1 to 50 and each carries 1 mark.
- 3) Read each question carefully, select proper alternative and answer in the O.M.R. sheet.
- 4) The OMR Sheet is given for answering the questions. The answer of each question is represented by (A) O, (B) O, (C) O and (D) O. Darken the circle ● of the correct answer with ball-pen.
- 5) Rough work is to be done in the space provided for this purpose in the Test Booklet only.
- 6) Set No. of Question Paper printed on the upper-most right side of the Question Paper is to be written in the column provided in the OMR sheet.
- 7) Use of simple calculator and log table is allowed, if required.
- 8) Notations used in this question paper have proper meaning.

- 9) Let R be the relation on the set N given by $R = \{(a,b) : a = b - 2, b > 6\}$. Choose the correct answer.

(A) $(2,4) \in R$

(B) $(3,8) \in R$

☒ (C) $(6,8) \in R$

(D) $(8,7) \in R$

Rough Work

2) $a * b = \frac{ab}{10}$ defined on $\mathbb{Q}$. Inverse of 0.001 is _____

(A) 100000

☒ (B) 10000

(C) 1000000

☒ (D) 1000

3) For sets $S = \{\pi, \pi^2, \pi^3\}$ and $T = \{e, e^2, e^3\}$, if $F^{-1} : T \rightarrow S$ is defined as $F^{-1} = \{(e, \pi^3), (e^2, \pi^2), (e^3, \pi)\}$, then function $F =$ _____

(A) $\{(\pi^3, e), (\pi^2, e^2), (\pi, e^3)\}$

(B) $\{(\pi, e^2), (\pi^3, e), (\pi^2, e^3)\}$

(C) $\{(e^2, \pi), (e^3, \pi^2), (e, \pi^3)\}$

☒ (D) $\{(\pi, e), (\pi^2, e^2), (\pi^3, e^3)\}$

4) $\sum_{i=0}^2 \cot^{-1}\{-(i+1)\} =$ _____



☒ (A) $\pi/2$



(B) $-3\pi/2$

(C) $-5\pi/2$

(D) $5\pi/2$

i) Which of the following is a graph of $f(x) = \tan^{-1}x$, ($x \in \mathbb{R}$)?





6) $\sec^{-1}x + \operatorname{cosec}^{-1}x + \cos^{-1}(x^{-1}) + \sin^{-1}(x^{-1}) =$ _____
 (where $|x| \geq 1, x \in \mathbb{R}$).

(A) $\frac{\pi}{2}$

(B) $\frac{3\pi}{2}$

(C) π

(D) 0

7) $\cot \left\{ \frac{2019\pi}{2} - \left(\operatorname{cosec}^{-1} \frac{5}{3} + \tan^{-1} \frac{2}{3} \right) \right\} = \underline{\hspace{2cm}}$

(A) $-17/6$

(B) $19/6$

(C) $17/6$

(D) $-19/6$

8) For a 3×4 matrix, elements are given by $a_{ij} = |-3i + 4j|$, then $\sum_{i=1}^3 (a_{ii})' = \underline{\hspace{2cm}}$

(A) 3^3

(B) 4^3

(C) 2^3

(D) 6^3

9) A is 3×3 matrix and $\det(A) = 7$. If $B = \operatorname{adj} A$ then $\det(AB) = \underline{\hspace{2cm}}$

(A) 7^5

(B) 7^2

(C) 7

(D) 7^3

10) If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$ and $A^2 - 5A = kI$ then $k = \underline{\hspace{2cm}}$

(A) -7

(B) 7

(C) 5

(D) -5

11) Matrices X and Y are inverse of each other then $\underline{\hspace{2cm}}$

(A) $XY = YX = 0$

(B) $XY = YX = -I$

(C) $XY = I, YX = -I$

(D) $X^{-1}Y^{-1} = Y^{-1}X^{-1} = I$

12) If $\Delta = \begin{vmatrix} x+y+z^2 & x^2+y+z & x+y^2+z \\ z^2 & x^2 & y^2 \\ x+y & y+z & x+z \end{vmatrix}$, (where $x \neq y \neq z$)

$x, y, z \in \mathbb{R} - \{0\}$ then $\Delta =$ _____

- (A) $x+y+z$ (B) 1
☒ (C) 0 (D) $x^2 + y^2 + z^2$

13) For $\Delta = \begin{vmatrix} 2019 & \textcircled{2020} & 2021 \\ 2022 & 2023 & 2024 \\ 2025 & 2026 & 2027 \end{vmatrix}$ sum of minor and cofactor of

2020 is _____

- (A) 2020 ☒ (B) 0
 (C) 4040 (D) -2020

14) If area of triangle is 35 sq. units with vertices $(2, -6)$, $(5, 4)$ and $(k, 4)$, then $k =$ _____

- (A) 1.2 (B) -20
☒ (C) -12, -2 (D) 12, -2

15) Let the function f be defined by

$$f(x) = \begin{cases} cx+1, & \text{if } x \leq 3 \\ dx+3, & \text{if } x > 3 \end{cases}$$

If f is continuous at $x=3$, then $d-c =$ _____

- ☒ (A) $-\frac{2}{3}$ (B) $\frac{3}{2}$
 (C) $-\frac{3}{2}$ (D) $\frac{2}{3}$

(16) If $y = (x+3)^2 \cdot (x+4)^3 \cdot (x+5)^4$, then first order derivative of y with respect to x is _____.

(A) $\frac{y}{x} \sum_{i=2}^4 \frac{i}{(x+1-i)}$

(B) $\frac{x}{y} \sum_{i=1}^3 \frac{i+1}{(x+1+i)}$

(C) $\frac{1}{y} \sum_{i=1}^3 \frac{i-1}{(x+1-i)}$

(D) $y \sum_{i=2}^4 \left(\frac{i}{(x+1)+i} \right)$

(17) If $y = \log_e(\log_e x)$, then $\frac{d^2y}{dx^2} =$ _____ (where $x > 1$).

(A) $-\frac{\log_e(ex)}{(x \cdot \log_e x)^2}$

(B) $\frac{\log_e(ex)}{(x \cdot \log_e x)^2}$

(C) $-\frac{(x \cdot \log_e x)^2}{\log_e(ex)}$

(D) $\frac{\log_e\left(\frac{e}{x}\right)}{(x \cdot \log_e x)^2}$

18) At which point the slope of the normal to the curve

$y = \sqrt{4x-3} - 1$ is $\frac{2}{3}$?

(A) (3, 2)

(B) $\left(\frac{43}{36}, \frac{1}{3}\right)$

(C) $\left(\frac{43}{16}, -\frac{7}{8}\right)$

(D) (2, 3)

19) Approximate value of $\sqrt{0.081} =$ _____.

(A) 0.2867

(B) 0.2850

(C) 0.2866

(D) 0.2845

20) Function $f(x) = \left\lfloor \sin x \right\rfloor, x \in \left(-\frac{\pi}{2}, 0\right)$ is :

- ☒ (A) Strictly increasing
 (B) Neither increasing nor decreasing
☒ (C) Only an increasing
 (D) Strictly decreasing

21) Local maximum value of the $f(x) = x + \frac{1}{x}, (x \neq 0)$ is _____.

- (A) 2 (B) -2
 (C) $\frac{1}{2}$ (D) $-\frac{1}{2}$

22) $\int \sqrt{\frac{\cos x - \cos^3 x}{1 - \cos^3 x}} dx = \text{_____} + C.$

$\left(\text{where } x \in \mathbb{R} - \left\{ \frac{k\pi}{2} / k \in \mathbb{Z} \right\} \right)$

- ☒ (A) $-\frac{2}{3} \sin^{-1}(\cos^{3/2} x)$ ☒ (B) $\frac{2}{3} \tan^{-1}(\cos^{3/2} x)$
 (C) $\frac{2}{3} \cos^{-1}(\sin^{3/2} x)$ (D) $\frac{2}{3} \sin^{-1}(\sin^{3/2} x)$

23) If $\int \frac{1}{e^x + 1} dx = px - q \log |1 + e^x| + C$ then

$p + q = \text{_____}.$

- (A) -2 ☒ (B) 2
 (C) 0 (D) 1

24) $\int e^{x^3} \cdot 5^{x^3} \cdot x \cdot [\log 25 + 3x] dx = \underline{\hspace{2cm}} + C.$

(A) $\frac{1}{6} \cdot e^{x^3} \cdot 5^{x^3} \cdot x$

✓(B) $\frac{1}{6} \cdot e^{x^3} \cdot 5^{x^3}$

(C) $e^{x^3} \cdot 5^{x^3} \cdot x$

(D) $e^{x^3} \cdot 5^{x^3}$

25) $\int \frac{dx}{\sqrt{2x-x^2}} = \underline{\hspace{2cm}} + C.$

✓(A) $\sin^{-1}(x-1)$

(B) $\frac{1}{2} \sin^{-1}(x-1)$

(C) $2 \sin^{-1}(x-1)$

(D) $\log|(x-1) + \sqrt{2x-x^2}|$

26) $\int_{-1}^{\sqrt{3}} \frac{dx}{1+x^2} = \underline{\hspace{2cm}}.$

(A) $\frac{\pi}{12}$

(B) $\frac{\pi}{6}$

✓(C) $\frac{7\pi}{12}$

(D) $\frac{5\pi}{12}$

27) $\int_0^{\pi} \cos^3 x \cdot \sin^4 x dx = \underline{\hspace{2cm}}.$

(A) π

✓(B) 0

(C) $-\pi$

(D) 2π

28) $\int_{-\pi/6}^{\pi/6} \sin^3 x \cos^2 x \, dx = \underline{\hspace{2cm}}$

(A) $\left(\frac{\pi}{6}\right)^3 - \left(\frac{\pi}{6}\right)^2$ ☒ (B) 0

(C) $\frac{1}{\sqrt{2}} - 1$ (D) $\left(\frac{\pi}{6}\right)^2 - \left(\frac{\pi}{6}\right)^5$

29) $\int_0^2 f(x) \, dx = \underline{\hspace{2cm}}$; where $f(x) = \max\{x, x^2\}$.

☒ (A) $\frac{17}{6}$ (B) $\frac{13}{6}$

(C) $\frac{8}{3}$ (D) $\frac{19}{6}$

30) Area bounded by curve $y = \tan \pi x$; $x \in \left[-\frac{1}{4}, \frac{1}{4}\right]$ and X-axis is

(A) $\log 2$ (B) $\frac{\log 2}{2}$

(C) $\frac{\log 2}{2\pi}$ ☒ (D) $\frac{\log 2}{\pi}$

31) If the area of the region bounded by two curves $y = x^2$ and $y = x^3$ is $\frac{k}{6}$ then $k = \underline{\hspace{2cm}}$.

☒ (A) $\frac{1}{2}$ (B) $\frac{1}{12}$

(C) $\frac{1}{3}$ (D) $\frac{1}{4}$

32) Area bounded by the ellipse $\frac{x^2}{4} + \frac{y^2}{16} = 4$ is _____.

(A) 8π

☒ (B) 32π

(C) 64π

(D) $\frac{\pi}{64}$

33) The order and degree of the differential equation $(y''')^3 + (y'')^4 + (y')^4 + y = 7$ are _____ respectively.

☒ (A) 3 and 3

(B) 1 and 4

(C) 4 and 1

(D) 2 and 4

34) The number of arbitrary constant in the particular solution of a differential equation of order 4 will be _____.

(A) 4

(B) 2

☒ (C) 0

(D) 1

35) Integrating factor of the differential equation

$y dx - (x + 2y^2) dy = 0$ is _____.

(A) $-\frac{1}{y}$

(B) $-y$

(C) y

☒ (D) $\frac{1}{y}$

36) Measure of the angle between the vectors $\vec{a} = \hat{i} - \hat{j} + \hat{k}$ and

$\vec{b} = \hat{i} + \hat{j} + \hat{k}$ is _____.

(A) $\cos^{-1} \frac{1}{\sqrt{3}}$

(B) $\pi - \cos^{-1} \frac{1}{3}$

☒ (C) $\sin^{-1} \frac{2\sqrt{2}}{3}$

(D) $\sin^{-1} \frac{1}{3}$

37) If $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} - 3\hat{k}$, then $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b})$
 $=$ _____.

(A) 8

~~(B)~~ -8

(C) -2

(D) 2

38) Find the area of a parallelogram whose adjacent sides are given by the vectors $\vec{a} = 3\hat{i} + 5\hat{j} - 2\hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} + 3\hat{k}$.

(A) $\sqrt{507}$

(B) $\sqrt{387}$

~~(C)~~ $\frac{1}{2}\sqrt{507}$

(D) 25

39) Let $|\vec{x}| = |\vec{y}| = |\vec{x} + \vec{y}| = 1$ and if measure of the angle between $\vec{x}$ and $\vec{y}$ is α , then $\sin \alpha =$ _____ $\leftarrow \alpha$

~~(A)~~ $-\frac{\sqrt{3}}{2}$

(B) $\frac{\sqrt{3}}{2}$

(C) $-\frac{1}{2}$

(D) 1

40) $\hat{i} \cdot (\hat{k} \times \hat{j}) + \hat{j} \cdot (\hat{i} \times \hat{k}) + \hat{k} \cdot (\hat{j} \times \hat{i}) + \hat{i} \cdot (\hat{i} \times \hat{j}) + \hat{j} \cdot (\hat{j} \times \hat{k})$
 $=$ _____

(A) 3

(B) 1

(C) -1

~~(D)~~ -3

41) For three vectors $\vec{a}, \vec{b}$ and $\vec{c}$, $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ and $|\vec{a}| = 3, |\vec{b}| = 4, |\vec{c}| = 5$, then evaluate $2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$.

(A) -25

(B) 50

(C) 100

~~(D)~~ -50

- 42) If the lines $\frac{2x-5}{k} = \frac{y+2}{-5} = \frac{z}{1}$ and $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ are perpendicular to each other, then value of k is _____.

(A) 7

(B) 14

(C) -7

(D) 26

- 43) If the plane $2x + 3y + 4z = 1$ intersects X-axis, Y-axis and Z-axis at the points A, B and C respectively, then the centroid of a ΔABC is _____.

(A) $\left(\frac{1}{6}, \frac{1}{9}, \frac{1}{12}\right)$

(B) (6, 9, 12)

(C) $\left(\frac{2}{3}, 1, \frac{4}{3}\right)$ (D) $\left(\frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right)$

- 44) Distance between the two planes $2x - 2y + z = 5$ and $6x - 6y + 3z = 25$ is _____ units.

(A) $\frac{20}{3}$ (B) $\frac{10}{9}$ (C) $\frac{20}{9}$

(D) 10

- 45) The objective function of a linear programming problem is _____.

(A) a constant

(B) a quadratic equation ✗

(C) a function to be optimized ✓

(D) an inequality

- 46) The vertices of the feasible region determined by some linear constraints are $(0,2)$, $(1,1)$, $(3,3)$, $(1,5)$. Let $Z = px + qy$ where $p, q > 0$. The condition on p and q so that the maximum of Z occurs at both the points $(3,3)$ and $(1,5)$ is _____.

- (A) $q = 2p$ (B) $p = 2q$
 ✓(C) $p = q$ (D) $p = 3q$

- 47) If the vertices of a feasible region are $O(0,0)$, $A(10,0)$, $B(0,20)$, $C(15,15)$, then minimum value of a objective function $Z = 10x - 20y + 30$ is _____.

- (A) 30 (B) 130
 (C) -120 ✓(D) -370

- 48) If $P(E) = 0.8$, $P(F) = 0.5$ and $P(F/E) = 0.4$, then $P(E/F) =$ _____.

- ✓(A) 0.64 (B) 0.32
 (C) 0.80 (D) 0.98

- 49) A random variable X has the following probability distribution:

X	0	1	2	3	4
P(X)	0.1	k	$2k$	$2k$	0.15

then $P(X \leq 1) =$ _____

- (A) 0.15 (B) 0.25
 (C) 0.55 (D) 0.75

5

- 50) The probability of obtaining an even prime number on each dice when a pair of dice is rolled is _____.

- (A) 1 (B) 0
 ✓(C) $\frac{1}{36}$ (D) $\frac{35}{36}$

050 (E)

(MARCH, 2020)
 SCIENCE STREAM
 (CLASS - XII)
 (New Course)

(Part - B)**Time : 2 Hours]****[Maximum Marks : 50****Instructions :**

- 1) Write in a clear legible handwriting.
- 2) There are three sections in Part - B of the question paper and total 1 to 18 questions are there.
- 3) All the questions are compulsory. Internal options are given.
- 4) The numbers at right side represent the marks of the question.
- 5) Start new section on new page.
- 6) Maintain sequence.
- 7) Use of simple calculator and log table is allowed, if required.
- 8) Use the graph paper to solve the problem of L.P.

SECTION-A

- Answer the following 1 to 8 questions as directed in the question. (Each question carries 2 marks) [16]

1) Find the value :

$$\tan \frac{1}{2} \left[\sin^{-1} \frac{2x}{1+x^2} + \cos^{-1} \frac{1-y^2}{1+y^2} \right], |x| < 1, y > 0 \text{ and } xy < 1, x \neq y.$$

2) If $y = 50 e^{10x} + 60 e^{-10x}$, prove that $\frac{d^2y}{dx^2} = 100y$.

3) Evaluate $\int_0^1 e^x dx$ as the limit of sum.

- 4) If the area bounded by the parabola $y^2 = 4ax$ and its latus rectum in the first quadrant is 48 units then, using integration find the value of a .

- 5) Find the area between the curves $y = 2x$ and $y = x^2$.

OR

Find the area of the smaller region bounded by the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ and the line $2x + 3y = 6$.

- 6) If the vectors $\vec{a}, \vec{b}, \vec{c}$ are coplanar then, prove that $\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}$ are coplanar.

- 7) Find the equation of the plane passing through the intersection of the planes $x + y + z - 6 = 0$ and $2x + 3y + 4z + 5 = 0$ and the point $(2, 3, 4)$.

- 8) Bag-I contains 3 gold and 4 silver coins while another Bag-II contains 5 gold and 6 silver coins. One coin is drawn at random from one of the bags. Find the probability that a randomly selected coin is of gold.

OR

Find the mean of the number obtained on a throw of an unbiased dice.

SECTION - B

- Answer the following 9 to 14 questions as directed in the question. (Each question carries 3 marks) [18]

- 9) Consider $f : \mathbb{R}^+ \rightarrow [-5, \infty)$ given by $f(x) = 9x^2 + 6x - 5$. Show that f is invertible with $f^{-1}(y) = \left(\frac{(\sqrt{y+6}) - 1}{3} \right)$; where $\mathbb{R}^+$ is the set of all non-negative real numbers.

10) Solve the following system of equations by matrix method.

$$x + y + z = 6, 2y + z = 7, x - y + z = 2$$

OR

Express the matrix $A = \begin{bmatrix} 3 & -2 & 1 \\ 4 & 0 & 6 \\ -1 & 2 & 1 \end{bmatrix}$ as the sum of a symmetric and a skew symmetric matrices.

11) If $x = a(\cos \theta + \theta \sin \theta)$ and $y = a(\sin \theta - \theta \cos \theta)$ then, find $\frac{d^2 y}{dx^2}$.

12) A line makes angles α, β, γ and δ with the diagonals of a cube, prove that

$$\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + \sin^2 \delta = \frac{8}{3}$$

OR

Find the equation of the line passing through $(1, 2, 3)$ and parallel to the planes $x - y + 2z - 5 = 0$ and $3x + y + z - 6 = 0$.

13) Solve the following linear programming problem graphically. Subject to the constraints : $x + y \leq 50, 3x + y \leq 90, x \geq 0, y \geq 0$, obtain the maximum and minimum values of $Z = 5x + 10y$.

14) If a fair coin is tossed 10 times, find the probability of

i) exactly 2 heads

ii) at least 9 heads

SECTION - C

- Answer the following 15 to 18 questions as directed in the question. (Each question carries 4 marks)

15) Using properties of determinants prove :

$$\begin{vmatrix} a & a^2 & 1+pa^3 \\ b & b^2 & 1+pb^3 \\ c & c^2 & 1+pc^3 \end{vmatrix} = (1+abc)(a-b)(b-c)(c-a)$$

- 16) Find the global maximum and minimum values of the function f given by $f(x) = 2x^3 - 15x^2 + 36x + 1, x \in [1, 5]$.

OR

Show that of all the rectangles inscribed in a given fixed circle, the square has the maximum area.

- 17) Find : $\int \sqrt[3]{\tan x} \, dx$; (where $x \neq \frac{k\pi}{2}, k \in \mathbb{Z}$)

18) Obtain the particular solution of the differential equation :

$$\left\{ x \cos\left(\frac{y}{x}\right) + y \sin\left(\frac{y}{x}\right) \right\} y \, dx = \left\{ y \sin\left(\frac{y}{x}\right) - x \cos\left(\frac{y}{x}\right) \right\} x \, dy$$

where : $y' = \frac{\pi}{2}$ when $x = 2$.

