



## PUZZLER

More than 300 years ago, Isaac Newton realized that the same gravitational force that causes apples to fall to the Earth also holds the Moon in its orbit. In recent years, scientists have used the Hubble Space Telescope to collect evidence of the gravitational force acting even farther away, such as at this protoplanetary disk in the constellation Taurus. What properties of an object such as a protoplanet or the Moon determine the strength of its gravitational attraction to another object? (Left, Larry West/FPG International; right, Courtesy of NASA)

### web

For more information about the Hubble, visit the Space Telescope Science Institute at <http://www.stsci.edu/>

## chapter

# 14

## The Law of Gravity

### Chapter Outline

- |  |  |
|--|--|
| <b>14.1</b> Newton's Law of Universal Gravitation              | <b>14.7</b> Gravitational Potential Energy   |
| <b>14.2</b> Measuring the Gravitational Constant               | <b>14.8</b> Energy Considerations in Planetary and Satellite Motion                      |
| <b>14.3</b> Free-Fall Acceleration and the Gravitational Force | <b>14.9</b> (Optional) The Gravitational Force Between an Extended Object and a Particle |
| <b>14.4</b> Kepler's Laws                                      | <b>14.10</b> (Optional) The Gravitational Force Between a Particle and a Spherical Mass  |
| <b>14.5</b> The Law of Gravity and the Motion of Planets       |  |
| <b>14.6</b> The Gravitational Field                            |  |

**B**efore 1687, a large amount of data had been collected on the motions of the Moon and the planets, but a clear understanding of the forces causing these motions was not available. In that year, Isaac Newton provided the key that unlocked the secrets of the heavens. He knew, from his first law, that a net force had to be acting on the Moon because without such a force the Moon would move in a straight-line path rather than in its almost circular orbit. Newton reasoned that this force was the gravitational attraction exerted by the Earth on the Moon. He realized that the forces involved in the Earth–Moon attraction and in the Sun–planet attraction were not something special to those systems, but rather were particular cases of a general and universal attraction between objects. In other words, Newton saw that the same force of attraction that causes the Moon to follow its path around the Earth also causes an apple to fall from a tree. As he put it, “I deduced that the forces which keep the planets in their orbs must be reciprocally as the squares of their distances from the centers about which they revolve; and thereby compared the force requisite to keep the Moon in her orb with the force of gravity at the surface of the Earth; and found them answer pretty nearly.”

In this chapter we study the law of gravity. We place emphasis on describing the motion of the planets because astronomical data provide an important test of the validity of the law of gravity. We show that the laws of planetary motion developed by Johannes Kepler follow from the law of gravity and the concept of conservation of angular momentum. We then derive a general expression for gravitational potential energy and examine the energetics of planetary and satellite motion. We close by showing how the law of gravity is also used to determine the force between a particle and an extended object.

### 14.1 NEWTON'S LAW OF UNIVERSAL GRAVITATION

You may have heard the legend that Newton was struck on the head by a falling apple while napping under a tree. This alleged accident supposedly prompted him to imagine that perhaps all bodies in the Universe were attracted to each other in the same way the apple was attracted to the Earth. Newton analyzed astronomical data on the motion of the Moon around the Earth. From that analysis, he made the bold assertion that the force law governing the motion of planets was the *same* as the force law that attracted a falling apple to the Earth. This was the first time that “earthly” and “heavenly” motions were unified. We shall look at the mathematical details of Newton’s analysis in Section 14.5.

In 1687 Newton published his work on the law of gravity in his treatise *Mathematical Principles of Natural Philosophy*. **Newton’s law of universal gravitation** states that



every particle in the Universe attracts every other particle with a force that is directly proportional to the product of their masses and inversely proportional to the square of the distance between them.

If the particles have masses  $m_1$  and  $m_2$  and are separated by a distance  $r$ , the magnitude of this gravitational force is

$$F_g = G \frac{m_1 m_2}{r^2} \quad (14.1)$$

where  $G$  is a constant, called the *universal gravitational constant*, that has been measured experimentally. As noted in Example 6.6, its value in SI units is

$$G = 6.673 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2 \quad (14.2)$$


The form of the force law given by Equation 14.1 is often referred to as an **inverse-square law** because the magnitude of the force varies as the inverse square of the separation of the particles.<sup>1</sup> We shall see other examples of this type of force law in subsequent chapters. We can express this force in vector form by defining a unit vector  $\hat{\mathbf{r}}_{12}$  (Fig. 14.1). Because this unit vector is directed from particle 1 to particle 2, the force exerted by particle 1 on particle 2 is

$$\mathbf{F}_{12} = -G \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}_{12} \quad (14.3)$$

where the minus sign indicates that particle 2 is attracted to particle 1, and hence the force must be directed toward particle 1. By Newton's third law, the force exerted by particle 2 on particle 1, designated  $\mathbf{F}_{21}$ , is equal in magnitude to  $\mathbf{F}_{12}$  and in the opposite direction. That is, these forces form an action–reaction pair, and  $\mathbf{F}_{21} = -\mathbf{F}_{12}$ .

Several features of Equation 14.3 deserve mention. The gravitational force is a field force that always exists between two particles, regardless of the medium that separates them. Because the force varies as the inverse square of the distance between the particles, it decreases rapidly with increasing separation. We can relate this fact to the geometry of the situation by noting that the intensity of light emanating from a point source drops off in the same  $1/r^2$  manner, as shown in Figure 14.2.

Another important point about Equation 14.3 is that **the gravitational force exerted by a finite-size, spherically symmetric mass distribution on a particle outside the distribution is the same as if the entire mass of the distribution were concentrated at the center**. For example, the force exerted by the




**Figure 14.1** The gravitational force between two particles is attractive. The unit vector  $\hat{\mathbf{r}}_{12}$  is directed from particle 1 to particle 2. Note that  $\mathbf{F}_{21} = -\mathbf{F}_{12}$ .

Properties of the gravitational force

### QuickLab

Inflate a balloon just enough to form a small sphere. Measure its diameter. Use a marker to color in a 1-cm square on its surface. Now continue inflating the balloon until it reaches twice the original diameter. Measure the size of the square you have drawn. Also note how the color of the marked area has changed. Have you verified what is shown in Figure 14.2?



**Figure 14.2** Light radiating from a point source drops off as  $1/r^2$ , a relationship that matches the way the gravitational force depends on distance. When the distance from the light source is doubled, the light has to cover four times the area and thus is one fourth as bright.

<sup>1</sup> An inverse relationship between two quantities  $x$  and  $y$  is one in which  $y = k/x$ , where  $k$  is a constant. A direct proportion between  $x$  and  $y$  exists when  $y = kx$ .

Earth on a particle of mass  $m$  near the Earth's surface has the magnitude


$$F_g = G \frac{M_E m}{R_E^2} \quad (14.4)$$

where  $M_E$  is the Earth's mass and  $R_E$  its radius. This force is directed toward the center of the Earth.

We have evidence of the fact that the gravitational force acting on an object is directly proportional to its mass from our observations of falling objects, discussed in Chapter 2. All objects, regardless of mass, fall in the absence of air resistance at the same acceleration  $g$  near the surface of the Earth. According to Newton's second law, this acceleration is given by  $g = F_g/m$ , where  $m$  is the mass of the falling object. If this ratio is to be the same for all falling objects, then  $F_g$  must be directly proportional to  $m$ , so that the mass cancels in the ratio. If we consider the more general situation of a gravitational force between any two objects with mass, such as two planets, this same argument can be applied to show that the gravitational force is proportional to one of the masses. We can choose *either* of the masses in the argument, however; thus, the gravitational force must be directly proportional to *both* masses, as can be seen in Equation 14.3.

## 14.2 MEASURING THE GRAVITATIONAL CONSTANT

The universal gravitational constant  $G$  was measured in an important experiment by Henry Cavendish (1731–1810) in 1798. The Cavendish apparatus consists of two small spheres, each of mass  $m$ , fixed to the ends of a light horizontal rod suspended by a fine fiber or thin metal wire, as illustrated in Figure 14.3. When two large spheres, each of mass  $M$ , are placed near the smaller ones, the attractive force between smaller and larger spheres causes the rod to rotate and twist the wire suspension to a new equilibrium orientation. The angle of rotation is measured by the deflection of a light beam reflected from a mirror attached to the vertical suspension. The deflection of the light is an effective technique for amplifying the motion. The experiment is carefully repeated with different masses at various separations. In addition to providing a value for  $G$ , the results show experimentally that the force is attractive, proportional to the product  $mM$ , and inversely proportional to the square of the distance  $r$ .



**Figure 14.3** Schematic diagram of the Cavendish apparatus for measuring  $G$ . As the small spheres of mass  $m$  are attracted to the large spheres of mass  $M$ , the rod between the two small spheres rotates through a small angle. A light beam reflected from a mirror on the rotating apparatus measures the angle of rotation. The dashed line represents the original position of the rod.

### EXAMPLE 14.1 Billiards, Anyone?


Three 0.300-kg billiard balls are placed on a table at the corners of a right triangle, as shown in Figure 14.4. Calculate the gravitational force on the cue ball (designated  $m_1$ ) resulting from the other two balls.

**Solution** First we calculate separately the individual forces on the cue ball due to the other two balls, and then we find the vector sum to get the resultant force. We can see graphically that this force should point upward and toward the

right. We locate our coordinate axes as shown in Figure 14.4, placing our origin at the position of the cue ball.

The force exerted by  $m_2$  on the cue ball is directed upward and is given by

$$\mathbf{F}_{21} = G \frac{m_2 m_1}{r_{21}^2} \mathbf{j}$$



**Figure 14.4** The resultant gravitational force acting on the cue ball is the vector sum  $\mathbf{F}_{21} + \mathbf{F}_{31}$ .

$$\begin{aligned} &= \left( 6.67 \times 10^{-11} \frac{\text{N} \cdot \text{m}^2}{\text{kg}^2} \right) \frac{(0.300 \text{ kg})(0.300 \text{ kg})}{(0.400 \text{ m})^2} \mathbf{j} \\ &= 3.75 \times 10^{-11} \mathbf{j} \text{ N} \end{aligned}$$

This result shows that the gravitational forces between everyday objects have extremely small magnitudes. The force exerted by  $m_3$  on the cue ball is directed to the right:

$$\begin{aligned} \mathbf{F}_{31} &= G \frac{m_3 m_1}{r_{31}^2} \mathbf{i} \\ &= \left( 6.67 \times 10^{-11} \frac{\text{N} \cdot \text{m}^2}{\text{kg}^2} \right) \frac{(0.300 \text{ kg})(0.300 \text{ kg})}{(0.300 \text{ m})^2} \mathbf{i} \\ &= 6.67 \times 10^{-11} \mathbf{i} \text{ N} \end{aligned}$$

Therefore, the resultant force on the cue ball is

$$\mathbf{F} = \mathbf{F}_{21} + \mathbf{F}_{31} = (3.75\mathbf{j} + 6.67\mathbf{i}) \times 10^{-11} \text{ N}$$

and the magnitude of this force is

$$\begin{aligned} F &= \sqrt{F_{21}^2 + F_{31}^2} = \sqrt{(3.75)^2 + (6.67)^2} \times 10^{-11} \\ &= 7.65 \times 10^{-11} \text{ N} \end{aligned}$$

**Exercise** Find the direction of  $\mathbf{F}$ .

**Answer**  $29.3^\circ$  counterclockwise from the positive  $x$  axis.

## 14.3 FREE-FALL ACCELERATION AND THE GRAVITATIONAL FORCE

In Chapter 5, when defining  $mg$  as the weight of an object of mass  $m$ , we referred to  $g$  as the magnitude of the free-fall acceleration. Now we are in a position to obtain a more fundamental description of  $g$ . Because the force acting on a freely falling object of mass  $m$  near the Earth's surface is given by Equation 14.4, we can equate  $mg$  to this force to obtain

$$\begin{aligned} mg &= G \frac{M_E m}{R_E^2} \\ g &= G \frac{M_E}{R_E^2} \end{aligned} \quad (14.5)$$

Free-fall acceleration near the Earth's surface

Now consider an object of mass  $m$  located a distance  $h$  above the Earth's surface or a distance  $r$  from the Earth's center, where  $r = R_E + h$ . The magnitude of the gravitational force acting on this object is

$$F_g = G \frac{M_E m}{r^2} = G \frac{M_E m}{(R_E + h)^2}$$

The gravitational force acting on the object at this position is also  $F_g = mg'$ , where  $g'$  is the value of the free-fall acceleration at the altitude  $h$ . Substituting this expres-

sion for  $F_g$  into the last equation shows that  $g'$  is

Variation of  $g$  with altitude

$$g' = \frac{GM_E}{r^2} = \frac{GM_E}{(R_E + h)^2} \quad (14.6)$$

Thus, it follows that  $g'$  decreases with increasing altitude. Because the weight of a body is  $mg'$ , we see that as  $r \rightarrow \infty$ , its weight approaches zero.

### EXAMPLE 14.2 Variation of $g$ with Altitude $h$

The International Space Station is designed to operate at an altitude of 350 km. When completed, it will have a weight (measured at the Earth's surface) of  $4.22 \times 10^6$  N. What is its weight when in orbit?

**Solution** Because the station is above the surface of the Earth, we expect its weight in orbit to be less than its weight on Earth,  $4.22 \times 10^6$  N. Using Equation 14.6 with  $h = 350$  km, we obtain

$$\begin{aligned} g' &= \frac{GM_E}{(R_E + h)^2} \\ &= \frac{(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2)(5.98 \times 10^{24} \text{ kg})}{(6.37 \times 10^6 \text{ m} + 0.350 \times 10^6 \text{ m})^2} \\ &= 8.83 \text{ m/s}^2 \end{aligned}$$

Because  $g'/g = 8.83/9.80 = 0.901$ , we conclude that the weight of the station at an altitude of 350 km is 90.1% of the value at the Earth's surface. So the station's weight in orbit is

$$(0.901)(4.22 \times 10^6 \text{ N}) = 3.80 \times 10^6 \text{ N}$$

Values of  $g'$  at other altitudes are listed in Table 14.1.

**TABLE 14.1** Free-Fall Acceleration  $g'$  at Various Altitudes Above the Earth's Surface

Altitude $h$ (km)	$g'$ (m/s <sup>2</sup> )
1 000	7.33
2 000	5.68
3 000	4.53
4 000	3.70
5 000	3.08
6 000	2.60
7 000	2.23
8 000	1.93
9 000	1.69
10 000	1.49
50 000	0.13
$\infty$	0

#### web

The official web site for the International Space Station is [www.station.nasa.gov](http://www.station.nasa.gov)

### EXAMPLE 14.3 The Density of the Earth


Using the fact that  $g = 9.80$  m/s<sup>2</sup> at the Earth's surface, find the average density of the Earth.

**Solution** Using  $g = 9.80$  m/s<sup>2</sup> and  $R_E = 6.37 \times 10^6$  m, we find from Equation 14.5 that  $M_E = 5.96 \times 10^{24}$  kg. From this result, and using the definition of density from Chapter 1, we obtain

$$\begin{aligned} \rho_E &= \frac{M_E}{V_E} = \frac{M_E}{\frac{4}{3}\pi R_E^3} = \frac{5.96 \times 10^{24} \text{ kg}}{\frac{4}{3}\pi(6.37 \times 10^6 \text{ m})^3} \\ &= 5.50 \times 10^3 \text{ kg/m}^3 \end{aligned}$$

Because this value is about twice the density of most rocks at the Earth's surface, we conclude that the inner core of the Earth has a density much higher than the average value. It is most amazing that the Cavendish experiment, which determines  $G$  (and can be done on a tabletop), combined with simple free-fall measurements of  $g$ , provides information about the core of the Earth.





Astronauts F. Story Musgrave and Jeffrey A. Hoffman, along with the Hubble Space Telescope and the space shuttle *Endeavor*, are all falling around the Earth.

## 14.4 KEPLER'S LAWS

People have observed the movements of the planets, stars, and other celestial bodies for thousands of years. In early history, scientists regarded the Earth as the center of the Universe. This so-called geocentric model was elaborated and formalized by the Greek astronomer Claudius Ptolemy (c. 100–c. 170) in the second century A.D. and was accepted for the next 1 400 years. In 1543 the Polish astronomer Nicolaus Copernicus (1473–1543) suggested that the Earth and the other planets revolved in circular orbits around the Sun (the heliocentric model).

The Danish astronomer Tycho Brahe (1546–1601) wanted to determine how the heavens were constructed, and thus he developed a program to determine the positions of both stars and planets. It is interesting to note that those observations of the planets and 777 stars visible to the naked eye were carried out with only a large sextant and a compass. (The telescope had not yet been invented.)

The German astronomer Johannes Kepler was Brahe's assistant for a short while before Brahe's death, whereupon he acquired his mentor's astronomical data and spent 16 years trying to deduce a mathematical model for the motion of the planets. Such data are difficult to sort out because the Earth is also in motion around the Sun. After many laborious calculations, Kepler found that Brahe's data on the revolution of Mars around the Sun provided the answer.




**Johannes Kepler** German astronomer (1571–1630) The German astronomer Johannes Kepler is best known for developing the laws of planetary motion based on the careful observations of Tycho Brahe. (Art Resource)

For more information about Johannes Kepler, visit our Web site at [www.saunderscollege.com/physics/](http://www.saunderscollege.com/physics/)

Kepler's analysis first showed that the concept of circular orbits around the Sun had to be abandoned. He eventually discovered that the orbit of Mars could be accurately described by an **ellipse**. Figure 14.5 shows the geometric description of an ellipse. The longest dimension is called the major axis and is of length  $2a$ , where  $a$  is the **semimajor axis**. The shortest dimension is the minor axis, of length  $2b$ , where  $b$  is the **semiminor axis**. On either side of the center is a **focal point**, a distance  $c$  from the center, where  $a^2 = b^2 + c^2$ . The Sun is located at one of the focal points of Mars's orbit. Kepler generalized his analysis to include the motions of all planets. The complete analysis is summarized in three statements known as **Kepler's laws**:

#### Kepler's laws

1. All planets move in elliptical orbits with the Sun at one focal point.
2. The radius vector drawn from the Sun to a planet sweeps out equal areas in equal time intervals.
3. The square of the orbital period of any planet is proportional to the cube of the semimajor axis of the elliptical orbit.



**Figure 14.5** Plot of an ellipse. The semimajor axis has a length  $a$ , and the semiminor axis has a length  $b$ . The focal points are located at a distance  $c$  from the center, where  $a^2 = b^2 + c^2$ .


Most of the planetary orbits are close to circular in shape; for example, the semimajor and semiminor axes of the orbit of Mars differ by only 0.4%. Mercury and Pluto have the most elliptical orbits of the nine planets. In addition to the planets, there are many asteroids and comets orbiting the Sun that obey Kepler's laws. Comet Halley is such an object; it becomes visible when it is close to the Sun every 76 years. Its orbit is very elliptical, with a semiminor axis 76% smaller than its semimajor axis.

Although we do not prove it here, Kepler's first law is a direct consequence of the fact that the gravitational force varies as  $1/r^2$ . That is, under an inverse-square gravitational-force law, the orbit of a planet can be shown mathematically to be an ellipse with the Sun at one focal point. Indeed, half a century after Kepler developed his laws, Newton demonstrated that these laws are a consequence of the gravitational force that exists between any two masses. Newton's law of universal gravitation, together with his development of the laws of motion, provides the basis for a full mathematical solution to the motion of planets and satellites.

## 14.5 THE LAW OF GRAVITY AND THE MOTION OF PLANETS

In formulating his law of gravity, Newton used the following reasoning, which supports the assumption that the gravitational force is proportional to the inverse square of the separation between the two interacting bodies. He compared the acceleration of the Moon in its orbit with the acceleration of an object falling near the Earth's surface, such as the legendary apple (Fig. 14.6). Assuming that both accelerations had the same cause—namely, the gravitational attraction of the Earth—Newton used the inverse-square law to reason that the acceleration of the Moon toward the Earth (centripetal acceleration) should be proportional to  $1/r_M^2$ , where  $r_M$  is the distance between the centers of the Earth and the Moon. Furthermore, the acceleration of the apple toward the Earth should be proportional to  $1/R_E^2$ , where  $R_E$  is the radius of the Earth, or the distance between the centers of the Earth and the apple. Using the values  $r_M = 3.84 \times 10^8$  m and





**Figure 14.6** As it revolves around the Earth, the Moon experiences a centripetal acceleration  $\mathbf{a}_M$  directed toward the Earth. An object near the Earth's surface, such as the apple shown here, experiences an acceleration  $\mathbf{g}$ . (Dimensions are not to scale.)

$R_E = 6.37 \times 10^6$  m, Newton predicted that the ratio of the Moon's acceleration  $a_M$  to the apple's acceleration  $g$  would be

$$\frac{a_M}{g} = \frac{(1/r_M)^2}{(1/R_E)^2} = \left(\frac{R_E}{r_M}\right)^2 = \left(\frac{6.37 \times 10^6 \text{ m}}{3.84 \times 10^8 \text{ m}}\right)^2 = 2.75 \times 10^{-4}$$

Therefore, the centripetal acceleration of the Moon is

$$a_M = (2.75 \times 10^{-4})(9.80 \text{ m/s}^2) = 2.70 \times 10^{-3} \text{ m/s}^2$$


Newton also calculated the centripetal acceleration of the Moon from a knowledge of its mean distance from the Earth and its orbital period,  $T = 27.32$  days  $= 2.36 \times 10^6$  s. In a time  $T$ , the Moon travels a distance  $2\pi r_M$ , which equals the circumference of its orbit. Therefore, its orbital speed is  $2\pi r_M/T$  and its centripetal acceleration is

$$\begin{aligned} a_M &= \frac{v^2}{r_M} = \frac{(2\pi r_M/T)^2}{r_M} = \frac{4\pi^2 r_M}{T^2} = \frac{4\pi^2(3.84 \times 10^8 \text{ m})}{(2.36 \times 10^6 \text{ s})^2} \\ &= 2.72 \times 10^{-3} \text{ m/s}^2 \approx \frac{9.80 \text{ m/s}^2}{60^2} \end{aligned}$$

In other words, because the Moon is roughly 60 Earth radii away, the gravitational acceleration at that distance should be about  $1/60^2$  of its value at the Earth's surface. This is just the acceleration needed to account for the circular motion of the Moon around the Earth. The nearly perfect agreement between this value and the value Newton obtained using  $g$  provides strong evidence of the inverse-square nature of the gravitational force law.

Although these results must have been very encouraging to Newton, he was deeply troubled by an assumption he made in the analysis. To evaluate the acceleration of an object at the Earth's surface, Newton treated the Earth as if its mass were all concentrated at its center. That is, he assumed that the Earth acted as a particle as far as its influence on an exterior object was concerned. Several years later, in 1687, on the basis of his pioneering work in the development of calculus, Newton proved that this assumption was valid and was a natural consequence of the law of universal gravitation.

Acceleration of the Moon



**Figure 14.7** A planet of mass  $M_p$  moving in a circular orbit around the Sun. The orbits of all planets except Mercury and Pluto are nearly circular.

Kepler's third law

### Kepler's Third Law

It is informative to show that Kepler's third law can be predicted from the inverse-square law for circular orbits.<sup>2</sup> Consider a planet of mass  $M_p$  moving around the Sun of mass  $M_S$  in a circular orbit, as shown in Figure 14.7. Because the gravitational force exerted by the Sun on the planet is a radially directed force that keeps the planet moving in a circle, we can apply Newton's second law ( $\Sigma F = ma$ ) to the planet:

$$\frac{GM_S M_p}{r^2} = \frac{M_p v^2}{r}$$

Because the orbital speed  $v$  of the planet is simply  $2\pi r/T$ , where  $T$  is its period of revolution, the preceding expression becomes

$$\frac{GM_S}{r^2} = \frac{(2\pi r/T)^2}{r}$$

$$T^2 = \left( \frac{4\pi^2}{GM_S} \right) r^3 = K_S r^3 \quad (14.7)$$

where  $K_S$  is a constant given by

$$K_S = \frac{4\pi^2}{GM_S} = 2.97 \times 10^{-19} \text{ s}^2/\text{m}^3$$

Equation 14.7 is Kepler's third law. It can be shown that the law is also valid for elliptical orbits if we replace  $r$  with the length of the semimajor axis  $a$ . Note that the constant of proportionality  $K_S$  is independent of the mass of the planet. Therefore, Equation 14.7 is valid for *any* planet.<sup>3</sup> Table 14.2 contains a collection of useful planetary data. The last column verifies that  $T^2/r^3$  is a constant. The small variations in the values in this column reflect uncertainties in the measured values of the periods and semimajor axes of the planets.

If we were to consider the orbit around the Earth of a satellite such as the Moon, then the proportionality constant would have a different value, with the Sun's mass replaced by the Earth's mass.

### EXAMPLE 14.4 The Mass of the Sun

Calculate the mass of the Sun using the fact that the period of the Earth's orbit around the Sun is  $3.156 \times 10^7$  s and its distance from the Sun is  $1.496 \times 10^{11}$  m.

**Solution** Using Equation 14.7, we find that

$$M_S = \frac{4\pi^2 r^3}{GT^2} = \frac{4\pi^2 (1.496 \times 10^{11} \text{ m})^3}{(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2) (3.156 \times 10^7 \text{ s})^2}$$

$$= 1.99 \times 10^{30} \text{ kg}$$

In Example 14.3, an understanding of gravitational forces enabled us to find out something about the density of the Earth's core, and now we have used this understanding to determine the mass of the Sun.

<sup>2</sup> The orbits of all planets except Mercury and Pluto are very close to being circular; hence, we do not introduce much error with this assumption. For example, the ratio of the semiminor axis to the semimajor axis for the Earth's orbit is  $b/a = 0.99986$ .

<sup>3</sup> Equation 14.7 is indeed a proportion because the ratio of the two quantities  $T^2$  and  $r^3$  is a constant. The variables in a proportion are not required to be limited to the first power only.

**TABLE 14.2** Useful Planetary Data


Body	Mass (kg)	Mean Radius (m)	Period of Revolution (s)	Mean Distance from Sun (m)	$\frac{T^2}{r^3}$ (s <sup>2</sup> /m <sup>3</sup> )
Mercury	$3.18 \times 10^{23}$	$2.43 \times 10^6$	$7.60 \times 10^6$	$5.79 \times 10^{10}$	$2.97 \times 10^{-19}$
Venus	$4.88 \times 10^{24}$	$6.06 \times 10^6$	$1.94 \times 10^7$	$1.08 \times 10^{11}$	$2.99 \times 10^{-19}$
Earth	$5.98 \times 10^{24}$	$6.37 \times 10^6$	$3.156 \times 10^7$	$1.496 \times 10^{11}$	$2.97 \times 10^{-19}$
Mars	$6.42 \times 10^{23}$	$3.37 \times 10^6$	$5.94 \times 10^7$	$2.28 \times 10^{11}$	$2.98 \times 10^{-19}$
Jupiter	$1.90 \times 10^{27}$	$6.99 \times 10^7$	$3.74 \times 10^8$	$7.78 \times 10^{11}$	$2.97 \times 10^{-19}$
Saturn	$5.68 \times 10^{26}$	$5.85 \times 10^7$	$9.35 \times 10^8$	$1.43 \times 10^{12}$	$2.99 \times 10^{-19}$
Uranus	$8.68 \times 10^{25}$	$2.33 \times 10^7$	$2.64 \times 10^9$	$2.87 \times 10^{12}$	$2.95 \times 10^{-19}$
Neptune	$1.03 \times 10^{26}$	$2.21 \times 10^7$	$5.22 \times 10^9$	$4.50 \times 10^{12}$	$2.99 \times 10^{-19}$
Pluto	$\approx 1.4 \times 10^{22}$	$\approx 1.5 \times 10^6$	$7.82 \times 10^9$	$5.91 \times 10^{12}$	$2.96 \times 10^{-19}$
Moon	$7.36 \times 10^{22}$	$1.74 \times 10^6$	—	—	—
Sun	$1.991 \times 10^{30}$	$6.96 \times 10^8$	—	—	—

### Kepler's Second Law and Conservation of Angular Momentum

Consider a planet of mass  $M_p$  moving around the Sun in an elliptical orbit (Fig. 14.8). The gravitational force acting on the planet is always along the radius vector, directed toward the Sun, as shown in Figure 14.9a. When a force is directed toward or away from a fixed point and is a function of  $r$  only, it is called a **central force**. The torque acting on the planet due to this force is clearly zero; that is, because  $\mathbf{F}$  is parallel to  $\mathbf{r}$ ,

$$\boldsymbol{\tau} = \mathbf{r} \times \mathbf{F} = \mathbf{r} \times F\hat{\mathbf{r}} = 0$$


(You may want to revisit Section 11.2 to refresh your memory on the vector product.) Recall from Equation 11.19, however, that torque equals the time rate of change of angular momentum:  $\boldsymbol{\tau} = d\mathbf{L}/dt$ . Therefore, **because the gravitational**



**Figure 14.8** Kepler's second law is called the law of equal areas. When the time interval required for a planet to travel from  $A$  to  $B$  is equal to the time interval required for it to go from  $C$  to  $D$ , the two areas swept out by the planet's radius vector are equal. Note that in order for this to be true, the planet must be moving faster between  $C$  and  $D$  than between  $A$  and  $B$ .



Separate views of Jupiter and of Periodic Comet Shoemaker–Levy 9—both taken with the Hubble Space Telescope about two months before Jupiter and the comet collided in July 1994—were put together with the use of a computer. Their relative sizes and distances were altered. The black spot on Jupiter is the shadow of its moon Io.



**Figure 14.9** (a) The gravitational force acting on a planet is directed toward the Sun, along the radius vector. (b) As a planet orbits the Sun, the area swept out by the radius vector in a time  $dt$  is equal to one-half the area of the parallelogram formed by the vectors  $\mathbf{r}$  and  $d\mathbf{r} = \mathbf{v}dt$ .

**force exerted by the Sun on a planet results in no torque on the planet, the angular momentum  $\mathbf{L}$  of the planet is constant:**

$$\mathbf{L} = \mathbf{r} \times \mathbf{p} = \mathbf{r} \times M_p \mathbf{v} = M_p \mathbf{r} \times \mathbf{v} = \text{constant} \quad (14.8)$$

Because  $\mathbf{L}$  remains constant, the planet's motion at any instant is restricted to the plane formed by  $\mathbf{r}$  and  $\mathbf{v}$ .

We can relate this result to the following geometric consideration. The radius vector  $\mathbf{r}$  in Figure 14.9b sweeps out an area  $dA$  in a time  $dt$ . This area equals one-half the area  $|\mathbf{r} \times d\mathbf{r}|$  of the parallelogram formed by the vectors  $\mathbf{r}$  and  $d\mathbf{r}$  (see Section 11.2). Because the displacement of the planet in a time  $dt$  is  $d\mathbf{r} = \mathbf{v}dt$ , we can say that

$$dA = \frac{1}{2} |\mathbf{r} \times d\mathbf{r}| = \frac{1}{2} |\mathbf{r} \times \mathbf{v} dt| = \frac{L}{2M_p} dt$$

$$\frac{dA}{dt} = \frac{L}{2M_p} = \text{constant} \quad (14.9)$$


where  $L$  and  $M_p$  are both constants. Thus, we conclude that

the radius vector from the Sun to a planet sweeps out equal areas in equal time intervals.

It is important to recognize that this result, which is Kepler's second law, is a consequence of the fact that the force of gravity is a central force, which in turn implies that angular momentum is constant. Therefore, Kepler's second law applies to *any* situation involving a central force, whether inverse-square or not.

### EXAMPLE 14.5 Motion in an Elliptical Orbit

A satellite of mass  $m$  moves in an elliptical orbit around the Earth (Fig. 14.10). The minimum distance of the satellite from the Earth is called the *perigee* (indicated by  $p$  in Fig.



**Figure 14.10** As a satellite moves around the Earth in an elliptical orbit, its angular momentum is constant. Therefore,  $mv_a r_a = mv_p r_p$ , where the subscripts  $a$  and  $p$  represent apogee and perigee, respectively.

14.10), and the maximum distance is called the *apogee* (indicated by  $a$ ). If the speed of the satellite at  $p$  is  $v_p$ , what is its speed at  $a$ ?

**Solution** As the satellite moves from perigee toward apogee, it is moving farther from the Earth. Thus, a component of the gravitational force exerted by the Earth on the satellite is opposite the velocity vector. Negative work is done on the satellite, which causes it to slow down, according to the work–kinetic energy theorem. As a result, we expect the speed at apogee to be lower than the speed at perigee.

The angular momentum of the satellite relative to the Earth is  $\mathbf{r} \times m\mathbf{v} = m\mathbf{r} \times \mathbf{v}$ . At the points  $a$  and  $p$ ,  $\mathbf{v}$  is perpendicular to  $\mathbf{r}$ . Therefore, the magnitude of the angular momentum at these positions is  $L_a = mv_a r_a$  and  $L_p = mv_p r_p$ . Because angular momentum is constant, we see that

$$mv_a r_a = mv_p r_p$$

$$v_a = \frac{r_p}{r_a} v_p$$

**Quick Quiz 14.1**

How would you explain the fact that Saturn and Jupiter have periods much greater than one year?

**14.6 THE GRAVITATIONAL FIELD**

When Newton published his theory of universal gravitation, it was considered a success because it satisfactorily explained the motion of the planets. Since 1687 the same theory has been used to account for the motions of comets, the deflection of a Cavendish balance, the orbits of binary stars, and the rotation of galaxies. Nevertheless, both Newton's contemporaries and his successors found it difficult to accept the concept of a force that acts through a distance, as mentioned in Section 5.1. They asked how it was possible for two objects to interact when they were not in contact with each other. Newton himself could not answer that question.

An approach to describing interactions between objects that are not in contact came well after Newton's death, and it enables us to look at the gravitational interaction in a different way. As described in Section 5.1, this alternative approach uses the concept of a **gravitational field** that exists at every point in space. When a particle of mass  $m$  is placed at a point where the gravitational field is  $\mathbf{g}$ , the particle experiences a force  $\mathbf{F}_g = m\mathbf{g}$ . In other words, the field exerts a force on the particle. Hence, the gravitational field  $\mathbf{g}$  is defined as

$$\mathbf{g} \equiv \frac{\mathbf{F}_g}{m} \quad (14.10)$$

Gravitational field

That is, the gravitational field at a point in space equals the gravitational force experienced by a *test particle* placed at that point divided by the mass of the test particle. Notice that the presence of the test particle is not necessary for the field to exist—the Earth creates the gravitational field. We call the object creating the field the *source particle* (although the Earth is clearly not a particle; we shall discuss shortly the fact that we can approximate the Earth as a particle for the purpose of finding the gravitational field that it creates). We can detect the presence of the field and measure its strength by placing a test particle in the field and noting the force exerted on it.


Although the gravitational force is inherently an interaction between two objects, the concept of a gravitational field allows us to “factor out” the mass of one of the objects. In essence, we are describing the “effect” that any object (in this case, the Earth) has on the empty space around itself in terms of the force that *would* be present if a second object were somewhere in that space.<sup>4</sup>

As an example of how the field concept works, consider an object of mass  $m$  near the Earth's surface. Because the gravitational force acting on the object has a magnitude  $GM_E m/r^2$  (see Eq. 14.4), the field  $\mathbf{g}$  at a distance  $r$  from the center of the Earth is

$$\mathbf{g} = \frac{\mathbf{F}_g}{m} = -\frac{GM_E}{r^2} \hat{\mathbf{r}} \quad (14.11)$$

where  $\hat{\mathbf{r}}$  is a unit vector pointing radially outward from the Earth and the minus

<sup>4</sup> We shall return to this idea of mass affecting the space around it when we discuss Einstein's theory of gravitation in Chapter 39.



**Figure 14.11** (a) The gravitational field vectors in the vicinity of a uniform spherical mass such as the Earth vary in both direction and magnitude. The vectors point in the direction of the acceleration a particle would experience if it were placed in the field. The magnitude of the field vector at any location is the magnitude of the free-fall acceleration at that location. (b) The gravitational field vectors in a small region near the Earth's surface are uniform in both direction and magnitude.

sign indicates that the field points toward the center of the Earth, as illustrated in Figure 14.11a. Note that the field vectors at different points surrounding the Earth vary in both direction and magnitude. In a small region near the Earth's surface, the downward field  $\mathbf{g}$  is approximately constant and uniform, as indicated in Figure 14.11b. Equation 14.11 is valid at all points *outside* the Earth's surface, assuming that the Earth is spherical. At the Earth's surface, where  $r = R_E$ ,  $\mathbf{g}$  has a magnitude of  $9.80 \text{ N/kg}$ .

## 14.7 GRAVITATIONAL POTENTIAL ENERGY

In Chapter 8 we introduced the concept of gravitational potential energy, which is the energy associated with the position of a particle. We emphasized that the gravitational potential energy function  $U = mgy$  is valid only when the particle is near the Earth's surface, where the gravitational force is constant. Because the gravitational force between two particles varies as  $1/r^2$ , we expect that a more general potential energy function—one that is valid without the restriction of having to be near the Earth's surface—will be significantly different from  $U = mgy$ .

Before we calculate this general form for the gravitational potential energy function, let us first verify that *the gravitational force is conservative*. (Recall from Section 8.2 that a force is conservative if the work it does on an object moving between any two points is independent of the path taken by the object.) To do this, we first note that the gravitational force is a central force. By definition, a central force is any force that is directed along a radial line to a fixed center and has a magnitude that depends only on the radial coordinate  $r$ . Hence, a central force can be represented by  $F(r)\hat{\mathbf{r}}$ , where  $\hat{\mathbf{r}}$  is a unit vector directed from the origin to the particle, as shown in Figure 14.12.

Consider a central force acting on a particle moving along the general path  $P$  to  $Q$  in Figure 14.12. The path from  $P$  to  $Q$  can be approximated by a series of



steps according to the following procedure. In Figure 14.12, we draw several thin wedges, which are shown as dashed lines. The outer boundary of our set of wedges is a path consisting of short radial line segments and arcs (gray in the figure). We select the length of the radial dimension of each wedge such that the short arc at the wedge's wide end intersects the actual path of the particle. Then we can approximate the actual path with a series of zigzag movements that alternate between moving along an arc and moving along a radial line.

By definition, a central force is always directed along one of the radial segments; therefore, the work done by  $\mathbf{F}$  along any radial segment is

$$dW = \mathbf{F} \cdot d\mathbf{r} = F(r) dr$$

You should recall that, by definition, the work done by a force that is perpendicular to the displacement is zero. Hence, the work done in moving along any arc is zero because  $\mathbf{F}$  is perpendicular to the displacement along these segments. Therefore, the total work done by  $\mathbf{F}$  is the sum of the contributions along the radial segments:

$$W = \int_{r_i}^{r_f} F(r) dr$$

where the subscripts  $i$  and  $f$  refer to the initial and final positions. Because the integrand is a function only of the radial position, this integral depends only on the initial and final values of  $r$ . Thus, the work done is the same over *any* path from  $P$  to  $Q$ . Because the work done is independent of the path and depends only on the end points, we conclude that *any central force is conservative*. We are now assured that a potential energy function can be obtained once the form of the central force is specified.


Recall from Equation 8.2 that the change in the gravitational potential energy associated with a given displacement is defined as the negative of the work done by the gravitational force during that displacement:

$$\Delta U = U_f - U_i = - \int_{r_i}^{r_f} F(r) dr \quad (14.12)$$

We can use this result to evaluate the gravitational potential energy function. Consider a particle of mass  $m$  moving between two points  $P$  and  $Q$  above the Earth's surface (Fig. 14.13). The particle is subject to the gravitational force given by Equation 14.1. We can express this force as


$$F(r) = - \frac{GM_E m}{r^2}$$

where the negative sign indicates that the force is attractive. Substituting this expression for  $F(r)$  into Equation 14.12, we can compute the change in the gravita-



**Figure 14.13** As a particle of mass  $m$  moves from  $P$  to  $Q$  above the Earth's surface, the gravitational potential energy changes according to Equation 14.12.


Work done by a central force




**Figure 14.12** A particle moves from  $P$  to  $Q$  while acted on by a central force  $\mathbf{F}$ , which is directed radially. The path is broken into a series of radial segments and arcs. Because the work done along the arcs is zero, the work done is independent of the path and depends only on  $r_f$  and  $r_i$ .

Change in gravitational potential energy

Gravitational potential energy of the Earth–particle system for  $r \geq R_E$



**Figure 14.14** Graph of the gravitational potential energy  $U$  versus  $r$  for a particle above the Earth's surface. The potential energy goes to zero as  $r$  approaches infinity.



**Figure 14.15** Three interacting particles.

tional potential energy function:

$$U_f - U_i = GM_E m \int_{r_i}^{r_f} \frac{dr}{r^2} = GM_E m \left[ -\frac{1}{r} \right]_{r_i}^{r_f}$$

$$U_f - U_i = -GM_E m \left( \frac{1}{r_f} - \frac{1}{r_i} \right) \quad (14.13)$$

As always, the choice of a reference point for the potential energy is completely arbitrary. It is customary to choose the reference point where the force is zero. Taking  $U_i = 0$  at  $r_i = \infty$ , we obtain the important result

$$U = -\frac{GM_E m}{r} \quad (14.14)$$

This expression applies to the Earth–particle system where the two masses are separated by a distance  $r$ , provided that  $r \geq R_E$ . The result is not valid for particles inside the Earth, where  $r < R_E$ . (The situation in which  $r < R_E$  is treated in Section 14.10.) Because of our choice of  $U_i$ , the function  $U$  is always negative (Fig. 14.14).

Although Equation 14.14 was derived for the particle–Earth system, it can be applied to any two particles. That is, the gravitational potential energy associated with any pair of particles of masses  $m_1$  and  $m_2$  separated by a distance  $r$  is

$$U = -\frac{Gm_1 m_2}{r} \quad (14.15)$$

This expression shows that the gravitational potential energy for any pair of particles varies as  $1/r$ , whereas the force between them varies as  $1/r^2$ . Furthermore, the potential energy is negative because the force is attractive and we have taken the potential energy as zero when the particle separation is infinite. Because the force between the particles is attractive, we know that an external agent must do positive work to increase the separation between them. The work done by the external agent produces an increase in the potential energy as the two particles are separated. That is,  $U$  becomes less negative as  $r$  increases.

When two particles are at rest and separated by a distance  $r$ , an external agent has to supply an energy at least equal to  $+Gm_1 m_2 / r$  in order to separate the particles to an infinite distance. It is therefore convenient to think of the absolute value of the potential energy as the *binding energy* of the system. If the external agent supplies an energy greater than the binding energy, the excess energy of the system will be in the form of kinetic energy when the particles are at an infinite separation.

We can extend this concept to three or more particles. In this case, the total potential energy of the system is the sum over all pairs of particles.<sup>5</sup> Each pair contributes a term of the form given by Equation 14.15. For example, if the system contains three particles, as in Figure 14.15, we find that

$$U_{\text{total}} = U_{12} + U_{13} + U_{23} = -G \left( \frac{m_1 m_2}{r_{12}} + \frac{m_1 m_3}{r_{13}} + \frac{m_2 m_3}{r_{23}} \right) \quad (14.16)$$

The absolute value of  $U_{\text{total}}$  represents the work needed to separate the particles by an infinite distance.

<sup>5</sup> The fact that potential energy terms can be added for all pairs of particles stems from the experimental fact that gravitational forces obey the superposition principle.

**EXAMPLE 14.6** The Change in Potential Energy

A particle of mass  $m$  is displaced through a small vertical distance  $\Delta y$  near the Earth's surface. Show that in this situation the general expression for the change in gravitational potential energy given by Equation 14.13 reduces to the familiar relationship  $\Delta U = mg \Delta y$ .

**Solution** We can express Equation 14.13 in the form

$$\Delta U = -GM_E m \left( \frac{1}{r_f} - \frac{1}{r_i} \right) = GM_E m \left( \frac{r_f - r_i}{r_i r_f} \right)$$

If both the initial and final positions of the particle are close to the Earth's surface, then  $r_f - r_i = \Delta y$  and  $r_i r_f \approx R_E^2$ . (Recall that  $r$  is measured from the center of the Earth.) Therefore, the change in potential energy becomes

$$\Delta U \approx \frac{GM_E m}{R_E^2} \Delta y = mg \Delta y$$

where we have used the fact that  $g = GM_E/R_E^2$  (Eq. 14.5). Keep in mind that the reference point is arbitrary because it is the *change* in potential energy that is meaningful.

## 14.8 ENERGY CONSIDERATIONS IN PLANETARY AND SATELLITE MOTION

Consider a body of mass  $m$  moving with a speed  $v$  in the vicinity of a massive body of mass  $M$ , where  $M \gg m$ . The system might be a planet moving around the Sun, a satellite in orbit around the Earth, or a comet making a one-time flyby of the Sun. If we assume that the body of mass  $M$  is at rest in an inertial reference frame, then the total mechanical energy  $E$  of the two-body system when the bodies are separated by a distance  $r$  is the sum of the kinetic energy of the body of mass  $m$  and the potential energy of the system, given by Equation 14.15:<sup>6</sup>

$$E = K + U$$

$$E = \frac{1}{2}mv^2 - \frac{GMm}{r} \quad (14.17)$$

This equation shows that  $E$  may be positive, negative, or zero, depending on the value of  $v$ . However, for a bound system,<sup>7</sup> such as the Earth–Sun system,  $E$  is necessarily *less than zero* because we have chosen the convention that  $U \rightarrow 0$  as  $r \rightarrow \infty$ .

We can easily establish that  $E < 0$  for the system consisting of a body of mass  $m$  moving in a circular orbit about a body of mass  $M \gg m$  (Fig. 14.16). Newton's second law applied to the body of mass  $m$  gives


$$\frac{GMm}{r^2} = ma = \frac{mv^2}{r}$$

<sup>6</sup> You might recognize that we have ignored the acceleration and kinetic energy of the larger body. To see that this simplification is reasonable, consider an object of mass  $m$  falling toward the Earth. Because the center of mass of the object–Earth system is effectively stationary, it follows that  $mv = M_E v_E$ . Thus, the Earth acquires a kinetic energy equal to

$$\frac{1}{2}M_E v_E^2 = \frac{1}{2} \frac{m^2}{M_E} v^2 = \frac{m}{M_E} K$$

where  $K$  is the kinetic energy of the object. Because  $M_E \gg m$ , this result shows that the kinetic energy of the Earth is negligible.

<sup>7</sup> Of the three examples provided at the beginning of this section, the planet moving around the Sun and a satellite in orbit around the Earth are bound systems—the Earth will always stay near the Sun, and the satellite will always stay near the Earth. The one-time comet flyby represents an unbound system—the comet interacts once with the Sun but is not bound to it. Thus, in theory the comet can move infinitely far away from the Sun.



**Figure 14.16** A body of mass  $m$  moving in a circular orbit about a much larger body of mass  $M$ .

Multiplying both sides by  $r$  and dividing by 2 gives

$$\frac{1}{2}mv^2 = \frac{GMm}{2r} \quad (14.18)$$

Substituting this into Equation 14.17, we obtain

$$E = \frac{GMm}{2r} - \frac{GMm}{r}$$

$$E = -\frac{GMm}{2r} \quad (14.19)$$

Total energy for circular orbits

This result clearly shows that **the total mechanical energy is negative in the case of circular orbits**. Note that **the kinetic energy is positive and equal to one-half the absolute value of the potential energy**. The absolute value of  $E$  is also equal to the binding energy of the system, because this amount of energy must be provided to the system to move the two masses infinitely far apart.

The total mechanical energy is also negative in the case of elliptical orbits. The expression for  $E$  for elliptical orbits is the same as Equation 14.19 with  $r$  replaced by the semimajor axis length  $a$ . Furthermore, the total energy is constant if we assume that the system is isolated. Therefore, as the body of mass  $m$  moves from  $P$  to  $Q$  in Figure 14.13, the total energy remains constant and Equation 14.17 gives

$$E = \frac{1}{2}mv_i^2 - \frac{GMm}{r_i} = \frac{1}{2}mv_f^2 - \frac{GMm}{r_f} \quad (14.20)$$

Combining this statement of energy conservation with our earlier discussion of conservation of angular momentum, we see that **both the total energy and the total angular momentum of a gravitationally bound, two-body system are constants of the motion**.

### EXAMPLE 14.7 Changing the Orbit of a Satellite

The space shuttle releases a 470-kg communications satellite while in an orbit that is 280 km above the surface of the Earth. A rocket engine on the satellite boosts it into a geosynchronous orbit, which is an orbit in which the satellite stays directly over a single location on the Earth. How much energy did the engine have to provide?

**Solution** First we must determine the radius of a geosynchronous orbit. Then we can calculate the change in energy needed to boost the satellite into orbit.

The period of the orbit  $T$  must be one day (86 400 s), so that the satellite travels once around the Earth in the same time that the Earth spins once on its axis. Knowing the period, we can then apply Kepler's third law (Eq. 14.7) to find the radius, once we replace  $K_S$  with  $K_E = 4\pi^2/GM_E = 9.89 \times 10^{-14} \text{ s}^2/\text{m}^3$ :

$$T^2 = K_E r^3$$

$$r = \sqrt[3]{\frac{T^2}{K_E}} = \sqrt[3]{\frac{(86\,400 \text{ s})^2}{9.89 \times 10^{-14} \text{ s}^2/\text{m}^3}} = 4.23 \times 10^7 \text{ m} = R_f$$

This is a little more than 26 000 mi above the Earth's surface.

We must also determine the initial radius (not the altitude above the Earth's surface) of the satellite's orbit when it was still in the shuttle's cargo bay. This is simply

$$R_E + 280 \text{ km} = 6.65 \times 10^6 \text{ m} = R_i$$

Now, applying Equation 14.19, we obtain, for the total initial and final energies,

$$E_i = -\frac{GM_E m}{2R_i} \quad E_f = -\frac{GM_E m}{2R_f}$$

The energy required from the engine to boost the satellite is

$$E_{\text{engine}} = E_f - E_i = -\frac{GM_E m}{2} \left( \frac{1}{R_f} - \frac{1}{R_i} \right)$$

$$= -\frac{(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2)(5.98 \times 10^{24} \text{ kg})(470 \text{ kg})}{2}$$

$$\times \left( \frac{1}{4.23 \times 10^7 \text{ m}} - \frac{1}{6.65 \times 10^6 \text{ m}} \right)$$

$$= 1.19 \times 10^{10} \text{ J}$$

This is the energy equivalent of 89 gal of gasoline. NASA engineers must account for the changing mass of the spacecraft as it ejects burned fuel, something we have not done here. Would you expect the calculation that includes the effect of this changing mass to yield a greater or lesser amount of energy required from the engine?

If we wish to determine how the energy is distributed after the engine is fired, we find from Equation 14.18 that the change in kinetic energy is  $\Delta K = (GM_E m/2)(1/R_f - 1/R_i) = -1.19 \times 10^{10} \text{ J}$  (a decrease),

and the corresponding change in potential energy is  $\Delta U = -GM_E m(1/R_f - 1/R_i) = 2.38 \times 10^{10} \text{ J}$  (an increase). Thus, the change in mechanical energy of the system is  $\Delta E = \Delta K + \Delta U = 1.19 \times 10^{10} \text{ J}$ , as we already calculated. The firing of the engine results in an increase in the total mechanical energy of the system. Because an increase in potential energy is accompanied by a decrease in kinetic energy, we conclude that the speed of an orbiting satellite decreases as its altitude increases.

## Escape Speed

Suppose an object of mass  $m$  is projected vertically upward from the Earth's surface with an initial speed  $v_i$ , as illustrated in Figure 14.17. We can use energy considerations to find the minimum value of the initial speed needed to allow the object to escape the Earth's gravitational field. Equation 14.17 gives the total energy of the object at any point. At the surface of the Earth,  $v = v_i$  and  $r = r_i = R_E$ . When the object reaches its maximum altitude,  $v = v_f = 0$  and  $r = r_f = r_{\text{max}}$ . Because the total energy of the system is constant, substituting these conditions into Equation 14.20 gives

$$\frac{1}{2}mv_i^2 - \frac{GM_E m}{R_E} = -\frac{GM_E m}{r_{\text{max}}}$$

Solving for  $v_i^2$  gives

$$v_i^2 = 2GM_E \left( \frac{1}{R_E} - \frac{1}{r_{\text{max}}} \right) \quad (14.21)$$

Therefore, if the initial speed is known, this expression can be used to calculate the maximum altitude  $h$  because we know that


$$h = r_{\text{max}} - R_E$$

We are now in a position to calculate **escape speed**, which is the minimum speed the object must have at the Earth's surface in order to escape from the influence of the Earth's gravitational field. Traveling at this minimum speed, the object continues to move farther and farther away from the Earth as its speed asymptotically approaches zero. Letting  $r_{\text{max}} \rightarrow \infty$  in Equation 14.21 and taking  $v_i = v_{\text{esc}}$ , we obtain

$$v_{\text{esc}} = \sqrt{\frac{2GM_E}{R_E}} \quad (14.22)$$

Note that this expression for  $v_{\text{esc}}$  is independent of the mass of the object. In other words, a spacecraft has the same escape speed as a molecule. Furthermore, the result is independent of the direction of the velocity and ignores air resistance.

If the object is given an initial speed equal to  $v_{\text{esc}}$ , its total energy is equal to zero. This can be seen by noting that when  $r \rightarrow \infty$ , the object's kinetic energy and its potential energy are both zero. If  $v_i$  is greater than  $v_{\text{esc}}$ , the total energy is greater than zero and the object has some residual kinetic energy as  $r \rightarrow \infty$ .



**Figure 14.17** An object of mass  $m$  projected upward from the Earth's surface with an initial speed  $v_i$  reaches a maximum altitude  $h$ .

Escape speed

**EXAMPLE 14.8** Escape Speed of a Rocket

Calculate the escape speed from the Earth for a 5 000-kg spacecraft, and determine the kinetic energy it must have at the Earth's surface in order to escape the Earth's gravitational field.

**Solution** Using Equation 14.22 gives

$$v_{\text{esc}} = \sqrt{\frac{2GM_E}{R_E}}$$

$$= \sqrt{\frac{2(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2)(5.98 \times 10^{24} \text{ kg})}{6.37 \times 10^6 \text{ m}}}$$

$$= 1.12 \times 10^4 \text{ m/s}$$

This corresponds to about 25 000 mi/h.

The kinetic energy of the spacecraft is

$$K = \frac{1}{2}mv_{\text{esc}}^2 = \frac{1}{2}(5.00 \times 10^3 \text{ kg})(1.12 \times 10^4 \text{ m/s})^2$$

$$= 3.14 \times 10^{11} \text{ J}$$

This is equivalent to about 2 300 gal of gasoline.

**TABLE 14.3**  
Escape Speeds from the  
Surfaces of the Planets,  
Moon, and Sun

Body	$v_{\text{esc}}$ (km/s)
Mercury	4.3
Venus	10.3
Earth	11.2
Moon	2.3
Mars	5.0
Jupiter	60
Saturn	36
Uranus	22
Neptune	24
Pluto	1.1
Sun	618

Equations 14.21 and 14.22 can be applied to objects projected from any planet. That is, in general, the escape speed from the surface of any planet of mass  $M$  and radius  $R$  is

$$v_{\text{esc}} = \sqrt{\frac{2GM}{R}}$$

Escape speeds for the planets, the Moon, and the Sun are provided in Table 14.3. Note that the values vary from 1.1 km/s for Pluto to about 618 km/s for the Sun. These results, together with some ideas from the kinetic theory of gases (see Chapter 21), explain why some planets have atmospheres and others do not. As we shall see later, a gas molecule has an average kinetic energy that depends on the temperature of the gas. Hence, lighter molecules, such as hydrogen and helium, have a higher average speed than heavier species at the same temperature. When the average speed of the lighter molecules is not much less than the escape speed of a planet, a significant fraction of them have a chance to escape from the planet.

This mechanism also explains why the Earth does not retain hydrogen molecules and helium atoms in its atmosphere but does retain heavier molecules, such as oxygen and nitrogen. On the other hand, the very large escape speed for Jupiter enables that planet to retain hydrogen, the primary constituent of its atmosphere.


**Quick Quiz 14.2**

If you were a space prospector and discovered gold on an asteroid, it probably would not be a good idea to jump up and down in excitement over your find. Why?

**Quick Quiz 14.3**

Figure 14.18 is a drawing by Newton showing the path of a stone thrown from a mountain-top. He shows the stone landing farther and farther away when thrown at higher and higher speeds (at points  $D$ ,  $E$ ,  $F$ , and  $G$ ), until finally it is thrown all the way around the Earth. Why didn't Newton show the stone landing at  $B$  and  $A$  before it was going fast enough to complete an orbit?





**Figure 14.18** “The greater the velocity . . . with which [a stone] is projected, the farther it goes before it falls to the Earth. We may therefore suppose the velocity to be so increased, that it would describe an arc of 1, 2, 5, 10, 100, 1000 miles before it arrived at the Earth, till at last, exceeding the limits of the Earth, it should pass into space without touching.” Sir Isaac Newton, *System of the World*.

### Optional Section

## 14.9 THE GRAVITATIONAL FORCE BETWEEN AN EXTENDED OBJECT AND A PARTICLE

We have emphasized that the law of universal gravitation given by Equation 14.3 is valid only if the interacting objects are treated as particles. In view of this, how can we calculate the force between a particle and an object having finite dimensions? This is accomplished by treating the extended object as a collection of particles and making use of integral calculus. We first evaluate the potential energy function, and then calculate the gravitational force from that function.

We obtain the potential energy associated with a system consisting of a particle of mass  $m$  and an extended object of mass  $M$  by dividing the object into many elements, each having a mass  $\Delta M_i$  (Fig. 14.19). The potential energy associated with the system consisting of any one element and the particle is  $U = -Gm\Delta M_i/r_i$ , where  $r_i$  is the distance from the particle to the element  $\Delta M_i$ . The total potential energy of the overall system is obtained by taking the sum over all elements as  $\Delta M_i \rightarrow 0$ . In this limit, we can express  $U$  in integral form as


$$U = -Gm \int \frac{dM}{r} \quad (14.23)$$

Once  $U$  has been evaluated, we obtain the force exerted by the extended object on the particle by taking the negative derivative of this scalar function (see Section 8.6). If the extended object has spherical symmetry, the function  $U$  depends only on  $r$ , and the force is given by  $-dU/dr$ . We treat this situation in Section 14.10. In principle, one can evaluate  $U$  for any geometry; however, the integration can be cumbersome.

An alternative approach to evaluating the gravitational force between a particle and an extended object is to perform a vector sum over all mass elements of the object. Using the procedure outlined in evaluating  $U$  and the law of universal gravitation in the form shown in Equation 14.3, we obtain, for the total force exerted on the particle

$$\mathbf{F}_g = -Gm \int \frac{dM}{r^2} \hat{\mathbf{r}} \quad (14.24)$$

where  $\hat{\mathbf{r}}$  is a unit vector directed from the element  $dM$  toward the particle (see Fig. 14.19) and the minus sign indicates that the direction of the force is opposite that of  $\hat{\mathbf{r}}$ . This procedure is not always recommended because working with a vector function is more difficult than working with the scalar potential energy function. However, if the geometry is simple, as in the following example, the evaluation of  $\mathbf{F}$  can be straightforward.




**Figure 14.19** A particle of mass  $m$  interacting with an extended object of mass  $M$ . The total gravitational force exerted by the object on the particle can be obtained by dividing the object into numerous elements, each having a mass  $\Delta M_i$ , and then taking a vector sum over the forces exerted by all elements.

Total force exerted on a particle by an extended object

**EXAMPLE 14.9** Gravitational Force Between a Particle and a Bar

The left end of a homogeneous bar of length  $L$  and mass  $M$  is at a distance  $h$  from a particle of mass  $m$  (Fig. 14.20). Calculate the total gravitational force exerted by the bar on the particle.

**Solution** The arbitrary segment of the bar of length  $dx$  has a mass  $dM$ . Because the mass per unit length is constant, it follows that the ratio of masses  $dM/M$  is equal to the ratio



**Figure 14.20** The gravitational force exerted by the bar on the particle is directed to the right. Note that the bar is *not* equivalent to a particle of mass  $M$  located at the center of mass of the bar.

of lengths  $dx/L$ , and so  $dM = (M/L) dx$ . In this problem, the variable  $r$  in Equation 14.24 is the distance  $x$  shown in Figure 14.20, the unit vector  $\hat{\mathbf{r}}$  is  $\hat{\mathbf{r}} = -\mathbf{i}$ , and the force acting on the particle is to the right; therefore, Equation 14.24 gives us

$$\mathbf{F}_g = -Gm \int_h^{h+L} \frac{Mdx}{L} \frac{1}{x^2} (-\mathbf{i}) = Gm \frac{M}{L} \int_h^{h+L} \frac{dx}{x^2} \mathbf{i}$$

$$\mathbf{F}_g = \frac{GmM}{L} \left[ -\frac{1}{x} \right]_h^{h+L} \mathbf{i} = \frac{GmM}{h(h+L)} \mathbf{i}$$

We see that the force exerted on the particle is in the positive  $x$  direction, which is what we expect because the gravitational force is attractive.

Note that in the limit  $L \rightarrow 0$ , the force varies as  $1/h^2$ , which is what we expect for the force between two point masses. Furthermore, if  $h \gg L$ , the force also varies as  $1/h^2$ . This can be seen by noting that the denominator of the expression for  $\mathbf{F}_g$  can be expressed in the form  $h^2(1 + L/h)$ , which is approximately equal to  $h^2$  when  $h \gg L$ . Thus, when bodies are separated by distances that are great relative to their characteristic dimensions, they behave like particles.

*Optional Section***14.10 THE GRAVITATIONAL FORCE BETWEEN A PARTICLE AND A SPHERICAL MASS**

We have already stated that a large sphere attracts a particle outside it as if the total mass of the sphere were concentrated at its center. We now describe the force acting on a particle when the extended object is either a spherical shell or a solid sphere, and then apply these facts to some interesting systems.

**Spherical Shell**

**Case 1.** If a particle of mass  $m$  is located outside a spherical shell of mass  $M$  at, for instance, point  $P$  in Figure 14.21a, the shell attracts the particle as though the mass of the shell were concentrated at its center. We can show this, as Newton did, with integral calculus. Thus, as far as the gravitational force acting on a particle outside the shell is concerned, a spherical shell acts no differently from the solid spherical distributions of mass we have seen.

**Case 2.** If the particle is located inside the shell (at point  $P$  in Fig. 14.21b), the gravitational force acting on it can be shown to be zero.


We can express these two important results in the following way:

$$\mathbf{F}_g = -\frac{GMm}{r^2} \hat{\mathbf{r}} \quad \text{for } r \geq R \quad (14.25a)$$

$$\mathbf{F}_g = 0 \quad \text{for } r < R \quad (14.25b)$$

The gravitational force as a function of the distance  $r$  is plotted in Figure 14.21c.

Force on a particle due to a spherical shell



**Figure 14.21** (a) The nonradial components of the gravitational forces exerted on a particle of mass  $m$  located at point  $P$  outside a spherical shell of mass  $M$  cancel out. (b) The spherical shell can be broken into rings. Even though point  $P$  is closer to the top ring than to the bottom ring, the bottom ring is larger, and the gravitational forces exerted on the particle at  $P$  by the matter in the two rings cancel each other. Thus, for a particle located at any point  $P$  inside the shell, there is no gravitational force exerted on the particle by the mass  $M$  of the shell. (c) The magnitude of the gravitational force versus the radial distance  $r$  from the center of the shell.

The shell does not act as a gravitational shield, which means that a particle inside a shell may experience forces exerted by bodies outside the shell.

### Solid Sphere

**Case 1.** If a particle of mass  $m$  is located outside a homogeneous solid sphere of mass  $M$  (at point  $P$  in Fig. 14.22), the sphere attracts the particle as though the

mass of the sphere were concentrated at its center. We have used this notion at several places in this chapter already, and we can argue it from Equation 14.25a. A solid sphere can be considered to be a collection of concentric spherical shells. The masses of all of the shells can be interpreted as being concentrated at their common center, and the gravitational force is equivalent to that due to a particle of mass  $M$  located at that center.


**Case 2.** If a particle of mass  $m$  is located inside a homogeneous solid sphere of mass  $M$  (at point  $Q$  in Fig. 14.22), the gravitational force acting on it is due *only* to the mass  $M'$  contained within the sphere of radius  $r < R$ , shown in Figure 14.22. In other words,

$$\mathbf{F}_g = -\frac{GmM}{r^2} \hat{\mathbf{r}} \quad \text{for } r \geq R \quad (14.26a)$$

$$\mathbf{F}_g = -\frac{GmM'}{r^2} \hat{\mathbf{r}} \quad \text{for } r < R \quad (14.26b)$$

This also follows from spherical-shell Case 1 because the part of the sphere that is

Force on a particle due to a solid sphere



**Figure 14.22** The gravitational force acting on a particle when it is outside a uniform solid sphere is  $GMm/r^2$  and is directed toward the center of the sphere. The gravitational force acting on the particle when it is inside such a sphere is proportional to  $r$  and goes to zero at the center.

farther from the center than  $Q$  can be treated as a series of concentric spherical shells that do not exert a net force on the particle because the particle is inside them. Because the sphere is assumed to have a uniform density, it follows that the ratio of masses  $M'/M$  is equal to the ratio of volumes  $V'/V$ , where  $V$  is the total volume of the sphere and  $V'$  is the volume within the sphere of radius  $r$  only:

$$\frac{M'}{M} = \frac{V'}{V} = \frac{\frac{4}{3}\pi r^3}{\frac{4}{3}\pi R^3} = \frac{r^3}{R^3}$$

Solving this equation for  $M'$  and substituting the value obtained into Equation 14.26b, we have

$$\mathbf{F}_g = -\frac{GmM}{R^3} r \hat{\mathbf{r}} \quad \text{for } r < R \quad (14.27)$$

This equation tells us that at the center of the solid sphere, where  $r = 0$ , the gravitational force goes to zero, as we intuitively expect. The force as a function of  $r$  is plotted in Figure 14.22.

**Case 3.** If a particle is located inside a solid sphere having a density  $\rho$  that is spherically symmetric but not uniform, then  $M'$  in Equation 14.26b is given by an integral of the form  $M' = \int \rho dV$ , where the integration is taken over the volume contained within the sphere of radius  $r$  in Figure 14.22. We can evaluate this integral if the radial variation of  $\rho$  is given. In this case, we take the volume element  $dV$  as the volume of a spherical shell of radius  $r$  and thickness  $dr$ , and thus  $dV = 4\pi r^2 dr$ . For example, if  $\rho = Ar$ , where  $A$  is a constant, it is left to a problem (Problem 63) to show that  $M' = \pi Ar^4$ .

Hence, we see from Equation 14.26b that  $F$  is proportional to  $r^2$  in this case and is zero at the center.

### Quick Quiz 14.4

A particle is projected through a small hole into the interior of a spherical shell. Describe

### EXAMPLE 14.10 A Free Ride, Thanks to Gravity

An object of mass  $m$  moves in a smooth, straight tunnel dug between two points on the Earth's surface (Fig. 14.23). Show that the object moves with simple harmonic motion, and find the period of its motion. Assume that the Earth's density is uniform.

**Solution** The gravitational force exerted on the object acts toward the Earth's center and is given by Equation 14.27:

$$\mathbf{F}_g = -\frac{GmM}{R^3} r \hat{\mathbf{r}}$$

We receive our first indication that this force should result in simple harmonic motion by comparing it to Hooke's law, first seen in Section 7.3. Because the gravitational force on the object is linearly proportional to the displacement, the object experiences a Hooke's law force.

The  $y$  component of the gravitational force on the object is balanced by the normal force exerted by the tunnel wall, and the  $x$  component is


$$F_x = -\frac{GmM_E}{R_E^3} r \cos \theta$$

Because the  $x$  coordinate of the object is  $x = r \cos \theta$ , we can write

$$F_x = -\frac{GmM_E}{R_E^3} x$$

Applying Newton's second law to the motion along the  $x$  direction gives

$$F_x = -\frac{GmM_E}{R_E^3} x = ma_x$$



**Figure 14.23** An object moves along a tunnel dug through the Earth. The component of the gravitational force  $\mathbf{F}_g$  along the  $x$  axis is the driving force for the motion. Note that this component always acts toward  $O$ .

Solving for  $a_x$ , we obtain

$$a_x = -\frac{GM_E}{R_E^3} x$$

If we use the symbol  $\omega^2$  for the coefficient of  $x$ — $GM_E/R_E^3 = \omega^2$ —we see that

$$(1) \quad a_x = -\omega^2 x$$

an expression that matches the mathematical form of Equation 13.9, which gives the acceleration of a particle in simple harmonic motion:  $a_x = -\omega^2 x$ . Therefore, Equation (1),

which we have derived for the acceleration of our object in the tunnel, is the acceleration equation for simple harmonic motion at angular speed  $\omega$  with

$$\omega = \sqrt{\frac{GM_E}{R_E^3}}$$

Thus, the object in the tunnel moves in the same way as a block hanging from a spring! The period of oscillation is

$$\begin{aligned} T &= \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{R_E^3}{GM_E}} \\ &= 2\pi \sqrt{\frac{(6.37 \times 10^6 \text{ m})^3}{(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2)(5.98 \times 10^{24} \text{ kg})}} \\ &= 5.06 \times 10^3 \text{ s} = 84.3 \text{ min} \end{aligned}$$

This period is the same as that of a satellite traveling in a circular orbit just above the Earth's surface (ignoring any trees, buildings, or other objects in the way). Note that the result is independent of the length of the tunnel.

A proposal has been made to operate a mass-transit system between any two cities, using the principle described in this example. A one-way trip would take about 42 min. A more precise calculation of the motion must account for the fact that the Earth's density is not uniform. More important, there are many practical problems to consider. For instance, it would be impossible to achieve a frictionless tunnel, and so some auxiliary power source would be required. Can you think of other problems?

the motion of the particle inside the shell.

## SUMMARY

**Newton's law of universal gravitation** states that the gravitational force of attraction between any two particles of masses  $m_1$  and  $m_2$  separated by a distance  $r$  has the magnitude

$$F_g = G \frac{m_1 m_2}{r^2} \quad (14.1)$$

where  $G = 6.673 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2$  is the universal gravitational constant. This equation enables us to calculate the force of attraction between masses under a wide variety of circumstances.

An object at a distance  $h$  above the Earth's surface experiences a gravitational force of magnitude  $mg'$ , where  $g'$  is the free-fall acceleration at that elevation:

$$g' = \frac{GM_E}{r^2} = \frac{GM_E}{(R_E + h)^2} \quad (14.6)$$



In this expression,  $M_E$  is the mass of the Earth and  $R_E$  is its radius. Thus, the weight of an object decreases as the object moves away from the Earth's surface.

**Kepler's laws of planetary motion** state that

1. All planets move in elliptical orbits with the Sun at one focal point.
2. The radius vector drawn from the Sun to a planet sweeps out equal areas in equal time intervals.
3. The square of the orbital period of any planet is proportional to the cube of the semimajor axis of the elliptical orbit.

Kepler's third law can be expressed as

$$T^2 = \left( \frac{4\pi^2}{GM_S} \right) r^3 \quad (14.7)$$

where  $M_S$  is the mass of the Sun and  $r$  is the orbital radius. For elliptical orbits, Equation 14.7 is valid if  $r$  is replaced by the semimajor axis  $a$ . Most planets have nearly circular orbits around the Sun.

The **gravitational field** at a point in space equals the gravitational force experienced by any test particle located at that point divided by the mass of the test particle:

$$\mathbf{g} = \frac{\mathbf{F}_g}{m} \quad (14.10)$$

The gravitational force is conservative, and therefore a potential energy function can be defined. The **gravitational potential energy** associated with two particles separated by a distance  $r$  is

$$U = -\frac{Gm_1m_2}{r} \quad (14.15)$$

where  $U$  is taken to be zero as  $r \rightarrow \infty$ . The total potential energy for a system of particles is the sum of energies for all pairs of particles, with each pair represented by a term of the form given by Equation 14.15.

If an isolated system consists of a particle of mass  $m$  moving with a speed  $v$  in the vicinity of a massive body of mass  $M$ , the total energy  $E$  of the system is the sum of the kinetic and potential energies:

$$E = \frac{1}{2}mv^2 - \frac{GMm}{r} \quad (14.17)$$

The total energy is a constant of the motion. If the particle moves in a circular orbit of radius  $r$  around the massive body and if  $M \gg m$ , the total energy of the system is

$$E = -\frac{GMm}{2r} \quad (14.19)$$

The total energy is negative for any bound system.

The **escape speed** for an object projected from the surface of the Earth is

$$v_{\text{esc}} = \sqrt{\frac{2GM_E}{R_E}} \quad (14.22)$$


## QUESTIONS

1. Use Kepler's second law to convince yourself that the Earth must move faster in its orbit during December, when it is closest to the Sun, than during June, when it is farthest from the Sun.
2. The gravitational force that the Sun exerts on the Moon is about twice as great as the gravitational force that the Earth exerts on the Moon. Why doesn't the Sun pull the Moon away from the Earth during a total eclipse of the Sun?
3. If a system consists of five particles, how many terms appear in the expression for the total potential energy? How many terms appear if the system consists of  $N$  particles?
4. Is it possible to calculate the potential energy function associated with a particle and an extended body without knowing the geometry or mass distribution of the extended body?
5. Does the escape speed of a rocket depend on its mass? Explain.
6. Compare the energies required to reach the Moon for a  $10^5$ -kg spacecraft and a  $10^3$ -kg satellite.
7. Explain why it takes more fuel for a spacecraft to travel from the Earth to the Moon than for the return trip. Estimate the difference.
8. Why don't we put a geosynchronous weather satellite in orbit around the 45th parallel? Wouldn't this be more useful for the United States than such a satellite in orbit around the equator?
9. Is the potential energy associated with the Earth–Moon system greater than, less than, or equal to the kinetic energy of the Moon relative to the Earth?
10. Explain why no work is done on a planet as it moves in a circular orbit around the Sun, even though a gravitational force is acting on the planet. What is the net work done on a planet during each revolution as it moves around the Sun in an elliptical orbit?
11. Explain why the force exerted on a particle by a uniform sphere must be directed toward the center of the sphere. Would this be the case if the mass distribution of the sphere were not spherically symmetric?
12. Neglecting the density variation of the Earth, what would be the period of a particle moving in a smooth hole dug between opposite points on the Earth's surface, passing through its center?
13. At what position in its elliptical orbit is the speed of a planet a maximum? At what position is the speed a minimum?
14. If you were given the mass and radius of planet X, how would you calculate the free-fall acceleration on the surface of this planet?
15. If a hole could be dug to the center of the Earth, do you think that the force on a mass  $m$  would still obey Equation 14.1 there? What do you think the force on  $m$  would be at the center of the Earth?
16. In his 1798 experiment, Cavendish was said to have "weighed the Earth." Explain this statement.
17. The gravitational force exerted on the *Voyager* spacecraft by Jupiter accelerated it toward escape speed from the Sun. How is this possible?
18. How would you find the mass of the Moon?
19. The *Apollo 13* spaceship developed trouble in the oxygen system about halfway to the Moon. Why did the spaceship continue on around the Moon and then return home, rather than immediately turn back to Earth?

## PROBLEMS

1, 2, 3 = straightforward, intermediate, challenging □ = full solution available in the *Student Solutions Manual and Study Guide*

WEB = solution posted at <http://www.saunderscollege.com/physics/>  = Computer useful in solving problem  = Interactive Physics

 = paired numerical/symbolic problems

### Section 14.1 Newton's Law of Universal Gravitation

### Section 14.2 Measuring the Gravitational Constant

### Section 14.3 Free-Fall Acceleration and the Gravitational Force

1. Determine the order of magnitude of the gravitational force that you exert on another person 2 m away. In your solution, state the quantities that you measure or estimate and their values.
2. A 200-kg mass and a 500-kg mass are separated by 0.400 m. (a) Find the net gravitational force exerted by these masses on a 50.0-kg mass placed midway between them. (b) At what position (other than infinitely remote ones) can the 50.0-kg mass be placed so as to experience a net force of zero?
3. Three equal masses are located at three corners of a square of edge length  $\ell$ , as shown in Figure P14.3. Find the gravitational field  $\mathbf{g}$  at the fourth corner due to these masses.
4. Two objects attract each other with a gravitational force of magnitude  $1.00 \times 10^{-8}$  N when separated by 20.0 cm. If the total mass of the two objects is 5.00 kg, what is the mass of each?
5. Three uniform spheres of masses 2.00 kg, 4.00 kg, and 6.00 kg are placed at the corners of a right triangle, as illustrated in Figure P14.5. Calculate the resultant gravitational force on the 4.00 kg sphere.




Figure P14.3




Figure P14.5

tational force on the 4.00-kg mass, assuming that the spheres are isolated from the rest of the Universe.

6. The free-fall acceleration on the surface of the Moon is about one-sixth that on the surface of the Earth. If the radius of the Moon is about  $0.250 R_E$ , find the ratio of their average densities,  $\rho_{\text{Moon}}/\rho_{\text{Earth}}$ .
7. During a solar eclipse, the Moon, Earth, and Sun all lie on the same line, with the Moon between the Earth and the Sun. (a) What force is exerted by the Sun on the Moon? (b) What force is exerted by the Earth on the Moon? (c) What force is exerted by the Sun on the Earth?
8. The center-to-center distance between the Earth and the Moon is 384 400 km. The Moon completes an orbit in 27.3 days. (a) Determine the Moon's orbital speed. (b) If gravity were switched off, the Moon would move along a straight line tangent to its orbit, as described by Newton's first law. In its actual orbit in 1.00 s, how far does the Moon fall below the tangent line and toward the Earth?
- WEB 9. When a falling meteoroid is at a distance above the Earth's surface of 3.00 times the Earth's radius, what is its acceleration due to the Earth's gravity?
10. Two ocean liners, each with a mass of 40 000 metric tons, are moving on parallel courses, 100 m apart. What is the magnitude of the acceleration of one of the liners toward the other due to their mutual gravitational attraction? (Treat the ships as point masses.)

11. A student proposes to measure the gravitational constant  $G$  by suspending two spherical masses from the ceiling of a tall cathedral and measuring the deflection of the cables from the vertical. Draw a free-body diagram of one of the masses. If two 100.0-kg masses are suspended at the end of 45.00-m-long cables, and the cables are attached to the ceiling 1.000 m apart, what is the separation of the masses?
12. On the way to the Moon, the Apollo astronauts reached a point where the Moon's gravitational pull became stronger than the Earth's. (a) Determine the distance of this point from the center of the Earth. (b) What is the acceleration due to the Earth's gravity at this point?

#### Section 14.4 Kepler's Laws

#### Section 14.5 The Law of Gravity and the Motion of Planets

13. A particle of mass  $m$  moves along a straight line with constant speed in the  $x$  direction, a distance  $b$  from the  $x$  axis (Fig. P14.13). Show that Kepler's second law is satisfied by demonstrating that the two shaded triangles in the figure have the same area when  $t_4 - t_3 = t_2 - t_1$ .






Figure P14.13

14. A communications satellite in geosynchronous orbit remains above a single point on the Earth's equator as the planet rotates on its axis. (a) Calculate the radius of its orbit. (b) The satellite relays a radio signal from a transmitter near the north pole to a receiver, also near the north pole. Traveling at the speed of light, how long is the radio wave in transit?
15. Plaskett's binary system consists of two stars that revolve in a circular orbit about a center of mass midway between them. This means that the masses of the two stars are equal (Fig. P14.15). If the orbital velocity of each star is 220 km/s and the orbital period of each is 14.4 days, find the mass  $M$  of each star. (For comparison, the mass of our Sun is  $1.99 \times 10^{30}$  kg.)
16. Plaskett's binary system consists of two stars that revolve in a circular orbit about a center of gravity midway between them. This means that the masses of the two stars are equal (see Fig. P14.15). If the orbital speed of each star is  $v$  and the orbital period of each is  $T$ , find the mass  $M$  of each star.



**Figure P14.15** Problems 15 and 16.


17. The *Explorer VIII* satellite, placed into orbit November 3, 1960, to investigate the ionosphere, had the following orbit parameters: perigee, 459 km; apogee, 2 289 km (both distances above the Earth's surface); and period, 112.7 min. Find the ratio  $v_p/v_a$  of the speed at perigee to that at apogee.
18. Comet Halley (Fig. P14.18) approaches the Sun to within 0.570 AU, and its orbital period is 75.6 years (AU is the symbol for astronomical unit, where  $1 \text{ AU} = 1.50 \times 10^{11} \text{ m}$  is the mean Earth–Sun distance). How far from the Sun will Halley's comet travel before it starts its return journey?



**Figure P14.18**

- WEB 19.** Io, a satellite of Jupiter, has an orbital period of 1.77 days and an orbital radius of  $4.22 \times 10^5 \text{ km}$ . From these data, determine the mass of Jupiter.

20. Two planets, X and Y, travel counterclockwise in circular orbits about a star, as shown in Figure P14.20. The radii of their orbits are in the ratio 3:1. At some time, they are aligned as in Figure P14.20a, making a straight line with the star. During the next five years, the angular displacement of planet X is  $90.0^\circ$ , as shown in Figure P14.20b. Where is planet Y at this time?



**Figure P14.20**

21. A synchronous satellite, which always remains above the same point on a planet's equator, is put in orbit around Jupiter so that scientists can study the famous red spot. Jupiter rotates once every 9.84 h. Use the data in Table 14.2 to find the altitude of the satellite.
22. Neutron stars are extremely dense objects that are formed from the remnants of supernova explosions. Many rotate very rapidly. Suppose that the mass of a certain spherical neutron star is twice the mass of the Sun and that its radius is 10.0 km. Determine the greatest possible angular speed it can have for the matter at the surface of the star on its equator to be just held in orbit by the gravitational force.
23. The Solar and Heliospheric Observatory (SOHO) spacecraft has a special orbit, chosen so that its view of the Sun is never eclipsed and it is always close enough to the Earth to transmit data easily. It moves in a near-circle around the Sun that is smaller than the Earth's circular orbit. Its period, however, is not less than 1 yr but is just equal to 1 yr. It is always located between the Earth and the Sun along the line joining them. Both objects exert gravitational forces on the observatory. Show that the spacecraft's distance from the Earth must be between  $1.47 \times 10^9 \text{ m}$  and  $1.48 \times 10^9 \text{ m}$ . In 1772 Joseph Louis Lagrange determined theoretically the special location that allows this orbit. The SOHO spacecraft took this position on February 14, 1996. (*Hint:* Use data that are precise to four digits. The mass of the Earth is  $5.983 \times 10^{24} \text{ kg}$ .)

### Section 14.6 The Gravitational Field

24. A spacecraft in the shape of a long cylinder has a length of 100 m, and its mass with occupants is 1 000 kg. It has




Figure P14.24

strayed too close to a 1.0-m-radius black hole having a mass 100 times that of the Sun (Fig. P14.24). The nose of the spacecraft is pointing toward the center of the black hole, and the distance between the nose and the black hole is 10.0 km. (a) Determine the total force on the spacecraft. (b) What is the difference in the gravitational fields acting on the occupants in the nose of the ship and on those in the rear of the ship, farthest from the black hole?

25. Compute the magnitude and direction of the gravitational field at a point  $P$  on the perpendicular bisector of two equal masses separated by a distance  $2a$ , as shown in Figure P14.25.




Figure P14.25

26. Find the gravitational field at a distance  $r$  along the axis of a thin ring of mass  $M$  and radius  $a$ .

### Section 14.7 Gravitational Potential Energy

Note: Assume that  $U = 0$  as  $r \rightarrow \infty$ .

27. A satellite of the Earth has a mass of 100 kg and is at an altitude of  $2.00 \times 10^6$  m. (a) What is the potential energy of the satellite–Earth system? (b) What is the magnitude of the gravitational force exerted by the Earth on the satellite? (c) What force does the satellite exert on the Earth?
28. How much energy is required to move a 1 000-kg mass from the Earth's surface to an altitude twice the Earth's radius?
29. After our Sun exhausts its nuclear fuel, its ultimate fate may be to collapse to a *white-dwarf* state, in which it has approximately the same mass it has now but a radius

equal to the radius of the Earth. Calculate (a) the average density of the white dwarf, (b) the acceleration due to gravity at its surface, and (c) the gravitational potential energy associated with a 1.00-kg object at its surface.

30. At the Earth's surface a projectile is launched straight up at a speed of 10.0 km/s. To what height will it rise? Ignore air resistance.
31. A system consists of three particles, each of mass 5.00 g, located at the corners of an equilateral triangle with sides of 30.0 cm. (a) Calculate the potential energy of the system. (b) If the particles are released simultaneously, where will they collide?
32. How much work is done by the Moon's gravitational field as a 1 000-kg meteor comes in from outer space and impacts the Moon's surface?

### Section 14.8 Energy Considerations in Planetary and Satellite Motion

33. A 500-kg satellite is in a circular orbit at an altitude of 500 km above the Earth's surface. Because of air friction, the satellite is eventually brought to the Earth's surface, and it hits the Earth with a speed of 2.00 km/s. How much energy was transformed to internal energy by means of friction?
34. (a) What is the minimum speed, relative to the Sun, that is necessary for a spacecraft to escape the Solar System if it starts at the Earth's orbit? (b) *Voyager 1* achieved a maximum speed of 125 000 km/h on its way to photograph Jupiter. Beyond what distance from the Sun is this speed sufficient for a spacecraft to escape the Solar System?
35. A satellite with a mass of 200 kg is placed in Earth orbit at a height of 200 km above the surface. (a) Assuming a circular orbit, how long does the satellite take to complete one orbit? (b) What is the satellite's speed? (c) What is the minimum energy necessary to place this satellite in orbit (assuming no air friction)?
36. A satellite of mass  $m$  is placed in Earth orbit at an altitude  $h$ . (a) Assuming a circular orbit, how long does the satellite take to complete one orbit? (b) What is the satellite's speed? (c) What is the minimum energy necessary to place this satellite in orbit (assuming no air friction)?

- WEB 37. A spaceship is fired from the Earth's surface with an initial speed of  $2.00 \times 10^4$  m/s. What will its speed be when it is very far from the Earth? (Neglect friction.)
38. A 1 000-kg satellite orbits the Earth at a constant altitude of 100 km. How much energy must be added to the system to move the satellite into a circular orbit at an altitude of 200 km?
39. A "treetop satellite" moves in a circular orbit just above the surface of a planet, which is assumed to offer no air resistance. Show that its orbital speed  $v$  and the escape speed from the planet are related by the expression  $v_{\text{esc}} = \sqrt{2}v$ .
40. The planet Uranus has a mass about 14 times the Earth's mass, and its radius is equal to about 3.7 Earth

radii. (a) By setting up ratios with the corresponding Earth values, find the acceleration due to gravity at the cloud tops of Uranus. (b) Ignoring the rotation of the planet, find the minimum escape speed from Uranus.

41. Determine the escape velocity for a rocket on the far side of Ganymede, the largest of Jupiter's moons. The radius of Ganymede is  $2.64 \times 10^6$  m, and its mass is  $1.495 \times 10^{23}$  kg. The mass of Jupiter is  $1.90 \times 10^{27}$  kg, and the distance between Jupiter and Ganymede is  $1.071 \times 10^9$  m. Be sure to include the gravitational effect due to Jupiter, but you may ignore the motions of Jupiter and Ganymede as they revolve about their center of mass (Fig. P14.41).




Figure P14.41

42. In Robert Heinlein's *The Moon is a Harsh Mistress*, the colonial inhabitants of the Moon threaten to launch rocks down onto the Earth if they are not given independence (or at least representation). Assuming that a rail gun could launch a rock of mass  $m$  at twice the lunar escape speed, calculate the speed of the rock as it enters the Earth's atmosphere. (By *lunar escape speed* we mean the speed required to escape entirely from a stationary Moon alone in the Universe.)
43. Derive an expression for the work required to move an Earth satellite of mass  $m$  from a circular orbit of radius  $2R_E$  to one of radius  $3R_E$ .

(Optional)

#### Section 14.9 The Gravitational Force Between an Extended Object and a Particle

44. Consider two identical uniform rods of length  $L$  and mass  $m$  lying along the same line and having their closest points separated by a distance  $d$  (Fig. P14.44). Show that the mutual gravitational force between these rods has a magnitude

$$F = \frac{Gm^2}{L^2} \ln \left( \frac{(L+d)^2}{d(2L+d)} \right)$$

45. A uniform rod of mass  $M$  is in the shape of a semicircle of radius  $R$  (Fig. P14.45). Calculate the force on a point mass  $m$  placed at the center of the semicircle.




Figure P14.44




Figure P14.45

(Optional)

#### Section 14.10 The Gravitational Force Between a Particle and a Spherical Mass

46. (a) Show that the period calculated in Example 14.10 can be written as

$$T = 2\pi \sqrt{\frac{R_E}{g}}$$

where  $g$  is the free-fall acceleration on the surface of the Earth. (b) What would this period be if tunnels were made through the Moon? (c) What practical problem regarding these tunnels on Earth would be removed if they were built on the Moon?

47. A 500-kg uniform solid sphere has a radius of 0.400 m. Find the magnitude of the gravitational force exerted by the sphere on a 50.0-g particle located (a) 1.50 m from the center of the sphere, (b) at the surface of the sphere, and (c) 0.200 m from the center of the sphere.
48. A uniform solid sphere of mass  $m_1$  and radius  $R_1$  is inside and concentric with a spherical shell of mass  $m_2$  and radius  $R_2$  (Fig. P14.48). Find the gravitational force exerted by the spheres on a particle of mass  $m$  located at (a)  $r = a$ , (b)  $r = b$ , and (c)  $r = c$ , where  $r$  is measured from the center of the spheres.




Figure P14.48



### ADDITIONAL PROBLEMS

49. Let  $\Delta g_M$  represent the difference in the gravitational fields produced by the Moon at the points on the Earth's surface nearest to and farthest from the Moon. Find the fraction  $\Delta g_M/g$ , where  $g$  is the Earth's gravitational field. (This difference is responsible for the occurrence of the *lunar tides* on the Earth.)
50. Two spheres having masses  $M$  and  $2M$  and radii  $R$  and  $3R$ , respectively, are released from rest when the distance between their centers is  $12R$ . How fast will each sphere be moving when they collide? Assume that the two spheres interact only with each other.
51. In Larry Niven's science-fiction novel *Ringworld*, a rigid ring of material rotates about a star (Fig. P14.51). The rotational speed of the ring is  $1.25 \times 10^6$  m/s, and its radius is  $1.53 \times 10^{11}$  m. (a) Show that the centripetal acceleration of the inhabitants is  $10.2$  m/s<sup>2</sup>. (b) The inhabitants of this ring world experience a normal contact force  $\mathbf{n}$ . Acting alone, this normal force would produce an inward acceleration of  $9.90$  m/s<sup>2</sup>. Additionally, the star at the center of the ring exerts a gravitational force on the ring and its inhabitants. The difference between the total acceleration and the acceleration provided by the normal force is due to the gravitational attraction of the central star. Show that the mass of the star is approximately  $10^{32}$  kg.




Figure P14.51

52. (a) Show that the rate of change of the free-fall acceleration with distance above the Earth's surface is

$$\frac{dg}{dr} = -\frac{2GM_E}{R_E^3}$$

This rate of change over distance is called a *gradient*.

(b) If  $h$  is small compared to the radius of the Earth, show that the difference in free-fall acceleration between two points separated by vertical distance  $h$  is

$$|\Delta g| = \frac{2GM_E h}{R_E^3}$$

(c) Evaluate this difference for  $h = 6.00$  m, a typical height for a two-story building.

53. A particle of mass  $m$  is located inside a uniform solid sphere of radius  $R$  and mass  $M$ , at a distance  $r$  from its center. (a) Show that the gravitational potential energy of the system is  $U = (GmM/2R^3)r^2 - 3GmM/2R$ . (b) Write an expression for the amount of work done by the gravitational force in bringing the particle from the surface of the sphere to its center.
54. *Voyagers 1* and *2* surveyed the surface of Jupiter's moon Io and photographed active volcanoes spewing liquid sulfur to heights of 70 km above the surface of this moon. Find the speed with which the liquid sulfur left the volcano. Io's mass is  $8.9 \times 10^{22}$  kg, and its radius is 1 820 km.
55. As an astronaut, you observe a small planet to be spherical. After landing on the planet, you set off, walking always straight ahead, and find yourself returning to your spacecraft from the opposite side after completing a lap of 25.0 km. You hold a hammer and a falcon feather at a height of 1.40 m, release them, and observe that they fall together to the surface in 29.2 s. Determine the mass of the planet.
56. A cylindrical habitat in space, 6.00 km in diameter and 30 km long, was proposed by G. K. O'Neill in 1974. Such a habitat would have cities, land, and lakes on the inside surface and air and clouds in the center. All of these would be held in place by the rotation of the cylinder about its long axis. How fast would the cylinder have to rotate to imitate the Earth's gravitational field at the walls of the cylinder?
- WEB 57. In introductory physics laboratories, a typical Cavendish balance for measuring the gravitational constant  $G$  uses lead spheres with masses of 1.50 kg and 15.0 g whose centers are separated by about 4.50 cm. Calculate the gravitational force between these spheres, treating each as a point mass located at the center of the sphere.
58. Newton's law of universal gravitation is valid for distances covering an enormous range, but it is thought to fail for very small distances, where the structure of space itself is uncertain. The crossover distance, far less than the diameter of an atomic nucleus, is called the *Planck length*. It is determined by a combination of the constants  $G$ ,  $c$ , and  $h$ , where  $c$  is the speed of light in vacuum and  $h$  is Planck's constant (introduced briefly in Chapter 11 and discussed in greater detail in Chapter 40) with units of angular momentum. (a) Use dimensional analysis to find a combination of these three universal constants that has units of length. (b) Determine the order of magnitude of the Planck length. (*Hint:* You will need to consider noninteger powers of the constants.)
59. Show that the escape speed from the surface of a planet of uniform density is directly proportional to the radius of the planet.
60. (a) Suppose that the Earth (or another object) has density  $\rho(r)$ , which can vary with radius but is spherically

symmetric. Show that at any particular radius  $r$  inside the Earth, the gravitational field strength  $g(r)$  will increase as  $r$  increases, if and only if the density there exceeds  $2/3$  the average density of the portion of the Earth inside the radius  $r$ . (b) The Earth as a whole has an average density of  $5.5 \text{ g/cm}^3$ , while the density at the surface is  $1.0 \text{ g/cm}^3$  on the oceans and about  $3 \text{ g/cm}^3$  on land. What can you infer from this?

**WEB 61.** Two hypothetical planets of masses  $m_1$  and  $m_2$  and radii  $r_1$  and  $r_2$ , respectively, are nearly at rest when they are an infinite distance apart. Because of their gravitational attraction, they head toward each other on a collision course. (a) When their center-to-center separation is  $d$ , find expressions for the speed of each planet and their relative velocity. (b) Find the kinetic energy of each planet just before they collide, if  $m_1 = 2.00 \times 10^{24} \text{ kg}$ ,  $m_2 = 8.00 \times 10^{24} \text{ kg}$ ,  $r_1 = 3.00 \times 10^6 \text{ m}$ , and  $r_2 = 5.00 \times 10^6 \text{ m}$ . (Hint: Both energy and momentum are conserved.)

**62.** The maximum distance from the Earth to the Sun (at our aphelion) is  $1.521 \times 10^{11} \text{ m}$ , and the distance of closest approach (at perihelion) is  $1.471 \times 10^{11} \text{ m}$ . If the Earth's orbital speed at perihelion is  $30.27 \text{ km/s}$ , determine (a) the Earth's orbital speed at aphelion, (b) the kinetic and potential energies at perihelion, and (c) the kinetic and potential energies at aphelion. Is the total energy constant? (Neglect the effect of the Moon and other planets.)

**63.** A sphere of mass  $M$  and radius  $R$  has a nonuniform density that varies with  $r$ , the distance from its center, according to the expression  $\rho = Ar$ , for  $0 \leq r \leq R$ . (a) What is the constant  $A$  in terms of  $M$  and  $R$ ? (b) Determine an expression for the force exerted on a particle of mass  $m$  placed outside the sphere. (c) Determine an expression for the force exerted on the particle if it is inside the sphere. (Hint: See Section 14.10 and note that the distribution is spherically symmetric.)

**64.** (a) Determine the amount of work (in joules) that must be done on a 100-kg payload to elevate it to a height of 1 000 km above the Earth's surface. (b) Determine the amount of additional work that is required to put the payload into circular orbit at this elevation.

**65.** X-ray pulses from Cygnus X-1, a celestial x-ray source, have been recorded during high-altitude rocket flights. The signals can be interpreted as originating when a blob of ionized matter orbits a black hole with a period of 5.0 ms. If the blob is in a circular orbit about a black hole whose mass is  $20M_{\text{Sun}}$ , what is the orbital radius?

**66.** Studies of the relationship of the Sun to its galaxy—the Milky Way—have revealed that the Sun is located near the outer edge of the galactic disk, about 30 000 lightyears from the center. Furthermore, it has been found that the Sun has an orbital speed of approximately 250 km/s around the galactic center. (a) What is the period of the Sun's galactic motion? (b) What is the order of magnitude of the mass of the Milky Way galaxy? Suppose that the galaxy is made mostly of stars,

of which the Sun is typical. What is the order of magnitude of the number of stars in the Milky Way?

**67.** The oldest artificial satellite in orbit is *Vanguard I*, launched March 3, 1958. Its mass is 1.60 kg. In its initial orbit, its minimum distance from the center of the Earth was 7.02 Mm, and its speed at this perigee point was 8.23 km/s. (a) Find its total energy. (b) Find the magnitude of its angular momentum. (c) Find its speed at apogee and its maximum (apogee) distance from the center of the Earth. (d) Find the semimajor axis of its orbit. (e) Determine its period.

**68.** A rocket is given an initial speed vertically upward of  $v_i = 2\sqrt{Rg}$  at the surface of the Earth, which has radius  $R$  and surface free-fall acceleration  $g$ . The rocket motors are quickly cut off, and thereafter the rocket coasts under the action of gravitational forces only. (Ignore atmospheric friction and the Earth's rotation.) Derive an expression for the subsequent speed  $v$  as a function of the distance  $r$  from the center of the Earth in terms of  $g$ ,  $R$ , and  $r$ .

**69.** Two stars of masses  $M$  and  $m$ , separated by a distance  $d$ , revolve in circular orbits about their center of mass (Fig. P14.69). Show that each star has a period given by

$$T^2 = \frac{4\pi^2 d^3}{G(M + m)}$$

(Hint: Apply Newton's second law to each star, and note that the center-of-mass condition requires that  $Mr_2 = mr_1$ , where  $r_1 + r_2 = d$ .)




Figure P14.69

**70.** (a) A 5.00-kg mass is released  $1.20 \times 10^7 \text{ m}$  from the center of the Earth. It moves with what acceleration relative to the Earth? (b) A  $2.00 \times 10^{24} \text{ kg}$  mass is released  $1.20 \times 10^7 \text{ m}$  from the center of the Earth. It moves with what acceleration relative to the Earth? Assume that the objects behave as pairs of particles, isolated from the rest of the Universe.

**71.** The acceleration of an object moving in the gravitational field of the Earth is

$$\mathbf{a} = -\frac{GM_E}{r^3} \mathbf{r}$$

where  $\mathbf{r}$  is the position vector directed from the center of the Earth to the object. Choosing the origin at the center of the Earth and assuming that the small object is moving in the  $xy$  plane, we find that the rectangular (cartesian) components of its acceleration are


$$a_x = -\frac{GM_E x}{(x^2 + y^2)^{3/2}} \quad a_y = -\frac{GM_E y}{(x^2 + y^2)^{3/2}}$$

Use a computer to set up and carry out a numerical pre-

diction of the motion of the object, according to Euler's method. Assume that the initial position of the object is  $x = 0$  and  $y = 2R_E$ , where  $R_E$  is the radius of the Earth. Give the object an initial velocity of 5 000 m/s in the  $x$  direction. The time increment should be made as small as practical. Try 5 s. Plot the  $x$  and  $y$  coordinates of the object as time goes on. Does the object hit the Earth? Vary the initial velocity until you find a circular orbit.

## ANSWERS TO QUICK QUIZZES

- 14.1 Kepler's third law (Eq. 14.7), which applies to all the planets, tells us that the period of a planet is proportional to  $r^{3/2}$ . Because Saturn and Jupiter are farther from the Sun than the Earth is, they have longer periods. The Sun's gravitational field is much weaker at Saturn and Jupiter than it is at the Earth. Thus, these planets experience much less centripetal acceleration than the Earth does, and they have correspondingly longer periods.
- 14.2 The mass of the asteroid might be so small that you would be able to exceed escape velocity by leg power alone. You would jump up, but you would never come back down!
- 14.3 Kepler's first law applies not only to planets orbiting the Sun but also to any relatively small object orbiting another under the influence of gravity. Any elliptical path that does not touch the Earth before reaching point  $G$  will continue around the other side to point  $V$  in a complete orbit (see figure in next column).





- 14.4** The gravitational force is zero inside the shell (Eq. 14.25b). Because the force on it is zero, the particle moves with constant velocity in the direction of its original motion outside the shell until it hits the wall opposite the entry hole. Its path thereafter depends on the nature of the collision and on the particle's original direction.

B.C.



5.15



by John Hart

