

Physics and Chemistry


Ve/hysics a/

8. Clear images of soft tissues can be well studied using
 - a) MRI
 - b) X-rays
 - ☒ c) Ultrasonics
 - d) I.R rays
 9. Particles which are not composite and hence truly elementary are
 - a) mesons
 - b) protons
 - c) neutrons
 - ☒ d) leptons
 10. A logic gate whose output will be in logic 0 state only when all inputs are in logic 1 state is called
 - a) AND
 - ☒ b) OR
 - c) NOR
 - d) NAND
 11. n type and p type semiconductors can be obtained by doping pure silicon respectively with
 - a) Arsenic Phosphorous
 - b) Indium Aluminium
 - c) Phosphorous Indium
 - ☒ d) Aluminium Boron
 12. In a CE amplifier $\beta=50$, $R_L=4K\Omega$, $R_i=500\Omega$. Power gain of the amplifier is
 - a) 2×10^4
 - ☒ b) 2×10^3
 - c) 2×10^5
 - d) 2×10^1
 13. Electrons are excited from $n=1$ to $n=4$ state. During downward transitions, possible number of spectral lines observed in Balmer series is
 - a) 4
 - ☒ b) 3
 - c) 2
 - d) 1
 14. IR region lies between
 - a) radio waves and microwave regions
 - b) microwaves and visible
 - c) visible and UV region
 - ☒ d) UV rays and X-ray region.
5. A proton λ_a will be
 - a) 2:1
 - ☒ b) $2\sqrt{2}$
 - c) 4:1
 - d) 1:2
 6. Raman
 - ☒ a) inci
 - b) inci
 - c) reso
 - d) mo
 7. ^{14}C and
 - a) iso
 - b) iso
 - ☒ c) iso
 - d) min
 8. In an int. the ratio
 - ☒ a) 3:1
 - b) 9:1
 - c) 2:1
 - d) 4:1
 19. In Young
 - a) d^2
 - ☒ b) d/λ
 - c) D
 - d) $2d$
 20. Newton
 - a) eq
 - ☒ b) eq
 - c) a
 - d) a
 21. It is di
 - a) lig
 - ☒ b) sp
 - c) lig
 - d) w

Space for calculation / rough work



$R_L = 4K\Omega$, $R_i = 500\Omega$

$R_i = 500\Omega$




5. A proton and an alpha particle are subjected to same potential difference V . Their de-Broglie wavelengths λ_p will be in the ratio
- ☒ 2:1
 - ☐ $2\sqrt{2}:1$
 - ☐ 4:1
 - ☐ 1:2
6. 'Raman Shift' depends on
- ☒ incident wavelength
 - ☐ incident intensity
 - ☐ resolving power of the spectrograph used
 - ☐ molecular energy levels of the scatterer.
7. ${}^6_6\text{C}^{14}$ and ${}^{14}_7\text{N}^{15}$ are the examples of
- ☐ isotopes
 - ☐ isobars
 - ☒ isotones
 - ☐ mirror nuclei
8. In an interference experiment, intensity ratio at the bright to dark fringe is 9:1. Amplitudes of interfering waves are in the ratio
- ☒ 3:1
 - ☐ 9:1
 - ☐ 2:1
 - ☐ 4:1
19. In Young's double slit experiment, 1st dark fringe occurs directly opposite to a slit. Wavelength of light used is
- ☐ d^2/D
 - ☒ d/D
 - ☐ D^2/d
 - ☐ $2d^2/D$
20. Newton's ring pattern in reflected system, viewed under white light consists of
- ☐ equally spaced bright and dark bands with central dark spot
 - ☒ equally spaced bright and dark bands with central white spot
 - ☐ a few coloured rings with central dark spot
 - ☐ a few coloured rings with central white spot
21. It is difficult to observe diffraction in case of light waves, because
- ☐ light waves can travel through vacuum
 - ☒ speed of light is more
 - ☐ light waves are transverse in nature
 - ☐ wavelength of light is small.

Space for calculation / rough work



22. A calcite crystal is placed over a dot on a paper sheet and the crystal is rotated. On viewing through the calcite or sees
- A single stationary dot
 - two stationary dots.
 - two dots rotating about one another
 - one dot rotating about the other stationary dot-sometimes coinciding with it
23. Critical angle of the medium is 45° . Polarising angle of incidence at the surface of the medium is
- 45°
 - 38°
 - 22.5°
 - 54.7°
24. If only 2% of the main current is to be passed through a Galvanometer of resistance G , the resistance of shunt should be
- $G/50$
 - $G/49$
 - $50G$
 - $49G$
25. A small current carrying loop of area A behaves like a tiny magnet of magnetic moment M . Current in the loop is
- MA
 - A/M
 - A^2M
 - M/A
26. Two concentric circular coils, each having 10 turns with radii 0.2m and 0.4m carry currents 0.2A and 0.3A respectively in opposite direction. Magnetic field at the centre is
- $(2/3)\mu_0$
 - $(5/4)\mu_0$
 - $(1/4)\mu_0$
 - $(1/6)\mu_0$
27. Material of permanent magnet has
- high retentivity and high coercivity
 - low retentivity and high coercivity
 - low retentivity and low coercivity
 - high retentivity and low coercivity.
28. Power factor of a series LCR circuit is
- R
 - Z/R
 - R/Z
 - RZ
29. An in
- 0
 - 0
 - 1
 - 1
30. Plane vibrat
- 0
 - 4
 - 2
 - 2
31. A cha
- 0.1 m
 - 0
 - 0
 - 0
32. Diele
- 0
 - 0
 - 0
 - 0
33. Dist
- 0
 - 0
 - 0
 - 0
34. Pot
- 0
 - 0
 - 0
 - 0
35. A n
- 0
 - 0
 - 0
 - 0

Space for calculation / rough work



Ver 9. An inductor 1H is connected across 220V 50Hz supply. Peak value of current is approximately,

- the calcite or
- a) 0.5A
 - b) 0.7A
 - c) 1A
 - d) 1.4A

30. Plane polarised light is passed through an analyser and the intensity of emerging light is reduced by 75%. Optical vibrations make an angle θ with the axis of analyser. Then θ is

- a) 60°
- b) 45°
- c) 30°
- d) 58°

31. A charge 10 nC is situated in a medium of relative permittivity 10. The potential due to this charge at a distance of 0.1 m is

- a) 900V
- b) 90V
- c) 9V
- d) 0.09V

32. Dielectric constant of a metal is

- a) zero
- b) infinite
- c) finite
- d) unpredictable

33. Distance between the two point charges is increased by 20%. Force of interaction between the charges

- a) increases by 10%
- b) decreases by 20%
- c) decreases by 17%
- d) decreases by 31%

34. Potential energy of 2 charges 10 nC each separated by a distance of 0.09m in air is

- a) 10 μ J
- b) 1 mJ
- c) 10 mJ
- d) 10 J

35. A metal plate of thickness $d/2$ is introduced in between the plates of a parallel plate air capacitor with plate separation of d . Capacity


- a) decreases 2 times
- b) increases 2 times
- c) remains same
- d) becomes zero.

Space for calculation / rough work

$Q = 20 \text{ nC}$ 0.09m in air is

$I = \frac{Q}{R}$

$\frac{20}{10} = 2$ (4.5)



$d/2 \times A$


$\frac{d}{2} = 10 \text{ nC}$

36. Specific resistance of a conductor material increases with
- increase with area of cross section
 - decrease in length
 - decrease in area of cross section
 - increases with temperature

37. The resistance of mercury at 4.2K is-
- infinity
 - greater than at lab temperature
 - same as that of lab temperature
 - almost zero.

38. Temperature coefficient of resistance of platinum is $4 \times 10^{-3} / ^\circ\text{C}$ at 20°C . Temperature at which increase in resistance of platinum is 10% its value at 20°C is
- 25°C
 - 70°C
 - 45°C
 - 100°C

39. Ideal voltmeter connected as shown reads



- 16V
 - 12V
 - 4V
 - 8V
40. When a charged particle moves perpendicular to a uniform magnetic field, then
- its momentum changes total energy is same.
 - both momentum and total energy remain the same.
 - both momentum and its total energy will change
 - total energy changes. Momentum remains same.

- 0.04 m/s
them no
a) $5/3$
b) $5/4$
c) $5/2$
d) $4/3$

2. Critical
a) G_1
b) G_2
c) W_1
d) D_1

3. A ray of
index of
a) 1.2
b) 1.4
c) 1.6
d) 1.8

4. In the
a) U_1
b) M_1
c) M_2
d) M_3

5. Conve
Focal l
a) f
b) g
c) k
d) -f


6. Two c
a) 0
b) 0
c) 0
d) -f

17. Eddy
a) b
b) p
c) F
d) F

$7 \times 10^3 \times 10^7$
 100×5

Space for calculation / rough work

41. 0.04 m of glass contains the same number of waves as 0.05 m of water, when monochromatic light passes through them normally. Refractive index of water is $\frac{4}{3}$. Refractive index of glass is
- $\frac{5}{3}$
 - $\frac{5}{4}$
 - $\frac{5}{2}$
 - $\frac{4}{5}$
42. Critical angle will be maximum, when light travels from
- Glass to air
 - Glass to water
 - Water to air
 - Diamond to air
43. A ray of light incident on one face of an equilateral prism at 60° enters and leaves the prism symmetrically. Refractive index of the prism material is
- 1.5
 - 1.62
 - 1.73
 - 1.8
44. In the spectrum of visible light produced by a prism dispersion is
- Uniform throughout the spectrum
 - Maximum in the middle decreases on either sides.
 - Maximum towards yellow
 - Maximum towards violet.
45. Convex lens of focal length f made of glass of Refractive index 1.5 is immersed in water of Refractive index $\frac{4}{3}$. Focal length is
- f
 - greater than f
 - less than f
 - $-f$
46. Two co-axial lenses of power $+4D$ and $-2D$ are placed in contact. The focal length of combination is
- 0.5m
 - 0.25m
 - 0.16m
 - 0.5m
47. Eddy currents are produced in a material when it is
- heated
 - placed in a time varying magnetic field.
 - placed in an electric field
 - placed in a uniform magnetic field.



$$\mu = 1.5 = \frac{3}{2}$$

$$\mu = \frac{4}{3}$$

$$2D$$

$$f = \frac{1}{2}$$

$$= 0.5$$

$$\frac{1}{4} \rightarrow \frac{1}{2}$$

$$\Rightarrow \frac{2-4}{4}$$

$$= \frac{-2}{4} = -\frac{1}{2}$$

Space for calculation / rough work

48. Transformer works on 220V. Its efficiency is 80%. Out put power is 8KW. Primary current is approximately,

- a) 35A
- b) 18A
- c) 22A
- d) 45A

$$\frac{V_1}{V_2} = \frac{I_2}{I_1} \quad I = 10$$

49. Quality factor of a series LCR circuit decreases from 3 to 2. Resonant frequency is 600Hz. Change in band width is

- a) zero
- b) 100Hz increase
- c) 100Hz decrease
- d) 300Hz increase

50. A stone dropped from the top of the tower reaches ground in 4 sec. Height of the tower is ($g=10\text{m/s}^2$)

- a) 20m
- b) 40m
- c) 60m
- d) 80m

$$s = 4$$

$$v = u + at$$

$$v =$$

51. Liquid crystal phase which are more close to the solid than to liquid is

- a) Nematic
- b) Smectic
- c) Lyotropic
- d) Cholesteric

52. If the Earth shrinks in its size (radius) mass remaining the same, the value of g on its surface will

- a) increase
- b) decrease
- c) remains same
- d) is reduced to zero.

53. Two rods of same area of cross section and lengths, and conductivities K_1 and K_2 are connected in series. Then in steady state conductivity of the combination is

- a) $(K_1 + K_2)/(K_1 K_2)$
- b) $2K_1 K_2/(K_1 + K_2)$
- c) $(K_1 + K_2)/2$
- d) $K_1 K_2/(K_1 + K_2)$

54. The square of the resultant of two equal forces acting at a point is equal to three times their product. Angle between them is

- a) 30°
- b) 45°
- c) 60°
- d) 90°

$$F_1 \quad F_2$$

$$(F_1 + F_2)^2 = 3 F_1 F_2$$

$$\vec{F}_1 + \vec{F}_2$$

$$\vec{F}_1 + \vec{F}_2$$

Space for calculation / rough work

24.9 A Curd

55. With the addition of impurities surface tension of a liquid

- a) increases
- b) decreases
- c) remains constant.
- ☒ d) may increase or decrease depending on impurities

56. Viscosity decreases with increase in temperature is the reason for

- (i) hot water moving faster than cold water
- (ii) more viscous oils are used in motor cars during summer than in winter
- a) only (i) is correct
- ☒ b) only (ii) correct
- c) both (i) and (ii) are correct
- d) both are wrong.

57. Moment of momentum of an electron revolving in second Bohr orbit of hydrogen is

- a) $2\pi h$
- b) $h/2\pi$
- ☒ c) h/π
- d) $2h/3\pi$

$$2h/2\pi$$

58. The existence of excitation and ionisation energies in an atom is an evidence for

- a) stability of an atom
- ☒ b) electrical neutrality of an atom
- c) small size of the atom
- d) stationary orbits in an atom.

59. Work function of a photosensitive metal is 3eV. The wavelength of incident radiations which can just eject photo-electrons from the metal is

- a) 600nm
- b) 510nm
- c) 414nm
- ☒ d) 378nm

$$\frac{1.24 \times 10^3}{3} = 413.3 \text{ nm}$$

in

60. Three identical capacitors are first connected in series and then in parallel. The ratio of effective capacitances in the two cases is

- a) 9:1
- b) 3:1
- c) 1:3
- ☒ d) 1:9

$$\frac{1}{C_s} = \frac{1}{C} + \frac{1}{C} + \frac{1}{C} = \frac{3}{C}$$

$$C_s = C/3$$

$$C_p = 3C$$

61. To dry ammonia gas the drying agent used is

- ☒ a) Con. H_2SO_4
- b) P_2O_5
- ☒ c) soda lime
- d) anhydrous CaCl_2



Space for calculation / rough work