

Topics : Method of Differentiation, Complex Number, Continuity & Derivability, Application of Derivatives, Sequence & Series, Function

Type of Questions

Single choice Objective (no negative marking) Q.1,2,3,4

(3 marks, 3 min.)

M.M., Min.

[12, 12]

Multiple choice objective (no negative marking) Q.5

(5 marks, 4 min.)

[5, 4]

Subjective Questions (no negative marking) Q.6,7,8

(4 marks, 5 min.)

[12, 15]

1. Let $y = \tan^{-1} \left(\frac{2\cos(3x^2 - 2) + 5\sin(3x^2 - 2)}{5\cos(3x^2 - 2) - 2\sin(3x^2 - 2)} \right)$, then $\frac{dy}{dx} =$

(A) $6x - 2$

(B) $6x$

(C) $5x$

(D) $\frac{6x}{x^2 + 1}$

2. If $y = at^2 + 2bt + c$ and $t = ax^2 + 2bx + c$, then $\frac{d^3y}{dx^3}$ equals

(A) $24a^2 (at + b)$

(B) $24a (ax + b)^2$

(C) $24a (at + b)^2$

(D) $24a^2 (ax + b)$

3. The complex number $z = x + iy$ for which $\log_{1/2} |z - 2| > \log_{1/2} |z|$, are given by:

(A) $\operatorname{Re}(z) \leq 1$

(B) $\operatorname{Im}(z) \leq 1$

(C) $\operatorname{Re}(z) > 1$

(D) $\operatorname{Im}(z) > 1$

4. If $g(x) = \frac{2h(x) + |h(x)|}{2h(x) - |h(x)|}$ where $h(x) = \sin x - \sin^n x$, $n \in \mathbb{R}^+$, the set of positive real numbers, and

$$f(x) = \begin{cases} [g(x)], & x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right) \\ 3, & x = \frac{\pi}{2} \end{cases} \quad \text{where } [.] \text{ denotes greatest integer function. Then}$$

(A) $f(x)$ is continuous and differentiable at $x = \frac{\pi}{2}$, when $0 < n < 1$

(B) $f(x)$ is continuous and differentiable at $x = \frac{\pi}{2}$, when $n > 1$

(C) $f(x)$ is continuous but not differentiable at $x = \frac{\pi}{2}$, when $0 < n < 1$

(D) $f(x)$ is continuous but not differentiable at $x = \frac{\pi}{2}$, when $n > 1$

5. For the series $S = 1 + \frac{1}{(1+3)} (1+2)^2 + \frac{1}{(1+3+5)} (1+2+3)^2 + \frac{1}{(1+3+5+7)} (1+2+3+4)^2 + \dots$

(A) 7th term is 16

(B) 7th term is 18

(C) sum of first 10th terms is $\frac{505}{4}$

(D) sum of first 10th term is $\frac{405}{4}$

6. Let $f(x) = \frac{1}{1-x}$, $g(x) = f \circ f \circ f \circ f \circ f(x)$ and $h(x) = \tan^{-1}(g(-x^2 - x))$, then find $\lim_{n \rightarrow \infty} \sum_{r=1}^n h(r)$.

7. Prove that in the curve $y = a \ell n(x^2 - a^2)$, sum of the tangent and subtangent varies as the product of the coordinates of the point of contact.

8. If the equation $a_0 x^n + a_1 x^{n-1} + \dots + a_{n-1} x = 0$, (where n is natural number) has a positive root α , prove that the equation $na_0 x^{n-1} + (n-1)a_1 x^{n-2} + \dots + a_{n-1} = 0$ also has a positive root smaller than α .

Answers Key

1. (B) 2. (D) 3. (C) 4. (B)

5. (A)(C) 6. $\frac{\pi}{4}$