

Topic : Atomic Structure
Type of Questions

Single choice Objective ('-1' negative marking) Q.1 to Q.4

(3 marks, 3 min.)

M.M., Min.


[12, 12]

Subjective Questions ('-1' negative marking) Q.5 to Q.8

(4 marks, 5 min.)

[16, 20]

1. In the following electronic configuration, some rules have been violated :



I : Hund

II : Pauli's exclusion

III : Aufbau

(A) I and II

(B) I and III

(C) II and III

(D) I, II and III

2. What is the potential difference through which an electron, with a de Broglie wavelength of $1.5 \text{ \AA}$ should be accelerated, if its de Broglie wavelength has to be reduced to $1 \text{ \AA}$:
- (A) 110 volts (B) 70 volts (C) 83 volts (D) 55 volts
3. X^{2+} is isoelectronic with sulphur and has $(Z + 2)$ neutrons (Z is atomic no. of element X). Hence, mass number of X^{2+} is :
- (A) 34 (B) 36 (C) 38 (D) 40
4. Which of the following compounds is isoelectronic with $[\text{NH}_3 \rightarrow \text{BH}_3]$:
- (A) B_2H_6 (B) C_2H_6 (C) C_2H_4 (D) C_3H_6
5. A neutral atom of an element has 2K, 8L, 9M and 2N electrons. Find out the following :
- (a) Atomic number of element (b) Total number of s electrons
 (c) Total number of p electrons (d) Total number of d electrons
 (e) Number of unpaired electrons in element
6. Calculate :
- (a) the value of spin only magnetic moment of Co^{3+} ion (in BM).
 (b) the number of radial nodes in a 3p-orbital.
 (c) the number of electrons with $(m = 0)$ in Mn^{2+} ion.
 (d) the orbital angular momentum for the unpaired electron in V^{4+} .
7. An element undergoes a reaction as shown :
- $$\text{X} + \text{e}^- \rightarrow \text{X}^- \quad \text{Energy released} = 30.876 \text{ eV}$$
- The energy released, is used to dissociate 8 g of H_2 molecules equally into H^+ and H^* , where H^* is in an excited state, in which the electron travels a path length equal to four times its de Broglie wavelength.
- (a) Determine the least amount (moles) of 'X' that would be required.
 Given: I.E. of H = 13.6 eV/atom
 Bond energy of H_2 = 4.526 eV/molecule.
 (b) Why is the amount of X calculated in the above question 'least'?
8. A compound of Vanadium has a spin magnetic moment 1.73 BM. Work out the electronic configuration of the Vanadium ion in the compound.

Answer Key

DPP No. # 21

1. (C) 2. (C) 3. (C) 4. (B)
5. (a) 21 (b) 8 (c) 12 (d) 1 (e) 1. 6. (a) $\sqrt{24}$ or 4.9 BM (b) 1 (c) 11 (d) $\frac{\sqrt{6} h}{2\pi}$
7. (a) $a = 4$ moles. 8. ${}_{23}\text{V}^{4+} : 1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^1$

Hints & Solutions

DPP No. # 21

1. Definition
2. $\lambda = \frac{12.3}{\sqrt{V}}$
3. ${}_{16}\text{S}^{32} = e^- = 16$
 $\text{X}^{+2} = e^- = 16$
 $(\because A = Z + N)$
4. (B) has same number of electrons i.e., 18.
 $[\text{NH}_3 \rightarrow \text{BH}_3] = 10 + 8 = 18$.
5. E.C. $\rightarrow 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 3d^1, 4s^2$
6. (a) $\text{Co}^{3+} : 1s^2 2s^2 2p^6 3s^2 3p^6 3d^6 \therefore 4$ unpaired electrons $\therefore \mu = \sqrt{4(4+2)} = \sqrt{24} = 4.9 \text{ BM}$
 (b) Number of radial nodes $= n - \ell - 1$
 Number of radial nodes in 3p orbital $= 3 - 1 - 1 = 1$
 (c) Number of electrons with $(m = 0)$ in $\text{Mn}^{2+} (1s^2 2s^2 2p^6 3s^2 3p^6 3d^5)$ ion $= 1s (2) + 2s (2) + 2p (2) + 3s (2) + 3p (2) + 3d (1) = 11$
 (d) Orbital angular momentum for the unpaired electron in V^{4+} lies in 3d orbital. $\therefore \ell = 2$

$$\therefore \text{Orbital angular momentum} = \sqrt{\ell(\ell+1)} \frac{h}{2\pi} = \frac{\sqrt{6} h}{2\pi}$$

7. (a) $x + e^- \rightarrow x^-$
 energy released = $E.A_1 = 30.87$ eV/atom
 Let no. of moles of X be a
 $\therefore a \times N_A \times 30.87 = 4 \times N_A \times 4.526 + 4 \times N_A \times 13.6 + 4 \times N_A \times 12.75 \Rightarrow a = 4$ moles.

8. Number of unpaired electron are given by

$$\text{Magnetic moment} = \sqrt{n(n+2)} \text{ B.M.}$$

where n is number of unpaired electrons

$$\text{or } 1.73 = \sqrt{n(n+2)} \quad \text{or } 1.73 \times 1.73 = n^2 + 2n \quad \therefore n = 1$$

Now Vanadium atom must have one unpaired electron and thus its configuration is

